

## REFERENCES

- Adams, R. D. and Victor, M., Ropper, A. H. Epilepsy and other seizure disorders. In; Principles of Neurology. 6<sup>th</sup>. 313-343. USA:McGraw-Hill. 1997.
- Ali, D. W. and Salter, M. W. NMDA receptor regulation by Src kinase signaling in excitatory synaptic transmission and plasticity. Current opinion Neurobiology 11 (2001): 336-342.
- Allredge, B. K. Seizure Disorder. Herfindal E. T. and Gourley, D. R. In; Textbook of Therapeutics: Drug and Disease Management 7<sup>th</sup>. 1107-1137. USA: Lipincott Williams & Wilkins. 2000.
- Anson, L. C., Chen, P. E., Wyllie, D. J. A., Colquhoun, D. and Schoepfer, R. Identification of amino acid residues of the NR2A subunit that control glutamate potency in recombinant NRI/NR2A NMDA receptors. Journal of Neuroscience. 18(1998): 581-589.
- Asher, P. Zinc, Src and NMDA receptors--a transmembrane connection. Nature Neuroscience. 1(3)(1998):173-175.
- Avanzini, G. and Franceschetti, S. Cellular biology of epileptogenesis. The Lancet 2(2003): 33-42.
- Avanzini, G., de Curtis, M. Panzica, F., Speafico, R. Intrinsic properties of nucleus reticularis thalamic neurones of the rat studied in vitro. Journal of Physiology 416 (1993): 111-122.
- Barchi, L., Ion channel mutations affecting muscle and brain. Currents Opinion in Neurology 11(1998): 461-468.
- Behe, P., Stern, P., Wyllie, D. J. A., Nassar, M., Schoepfer, R. and Colquhoun, D. Determination of NMDA NR1 subunit copy number in recombinant NMDA receptors. Proc. R. Soc. Lond. B. Biol Sci 262 (1995):202-213.
- Binnie, C. D. Vagus nerve stimulation for epilepsy: a review. Seizure 9 (2000): 161-169.
- Buller, A. L. and Monaghan, D. T. Pharmacological heterogeneity of NMDA receptors: characterization of NR1a/NR2D heteromers expressed in *Xenopus* oocytes. European Journal of Pharmacology 320 (1997): 87-94.

- Cartmell, J. and Schoepp, D. D. Review: Regulation of neurotransmitter release by metabotropic glutamate receptors. Journal of Neurochemistry 75 (2000): 889-907.
- Catterall, W. A. Structure and regulation of voltage-gate  $\text{Ca}^{2+}$  channels. Annu. Rev. Cell Dev. Biol. 16 (2000): 211-225.
- Chao, T. I., Alzheimer, C., Effect of phenytoin on the persistent  $\text{Na}^+$  current of mammalian CNS neurons. Neuroreport 6 (1995): 1778-1780.
- Chatterton, J. E. Awobuloyi, M., Premkumar, L. S. et al. Excitatory glycine receptors containing the NR3 family of NMDA receptor subunits. Nature 415 (2002): 793-798.
- Chazot, P. L. and Stephenson, F. A. Molecular dissection of native mammalian forebrain NMDA receptors containing the NRI C2 exon: direct demonstration of NMDA receptors comprising NRI, NR2A, and NR2B subunit within the same complex. Journal of Neurochemistry 69 (1997): 2138-2144.
- Chunngam, T. Preliminary studies of anticonvulsion mechanisms of N-(2-propylpentanoyl) urea. Master's Thesis. Chulalongkorn University, Thailand. 1996.
- Collingridge, G. L. and Singer, W. Excitatory amino acid receptors and synaptic plasticity. Trends in Pharmacological Sciences 11(1990), 290-296.
- Commission on Classification and Terminology of the International League Against Epilepsy. Proposal for revised clinical and electroencephalographic classification of epileptic seizures. Epilepsia. 22 (1981): 489-501.
- Commission on Classification and Terminology of the International League Against Epilepsy. Proposal for revised clinical and electroencephalographic classification of epilepsies and epileptic syndromes. Epilepsia. 30 (1989): 389-399.
- Cooper, E. C., Harrington, E., Jan, Y. N., Jan, Y. Y. M. Channel KCNQ2 subunits are localized to key sites for control of neuronal network oscillation and synchronization in mouse brain. Journal of Neuroscience 21 (2001): 9529-9540.
- Cosford ND, Meinke PT, Stauderman KA, Hess SD. Recent advances in the modulation of voltage-gated ion channels for the treatment of epilepsy. Curr Drug Target CNS Neurol Disord. 1(1) (2002): 81-104.

- Coulter, D. A., Huguenard, J. R., Prince, D. A. Calcium currents in rat thalamocortical relay neurones: kinetic properties of the transient low-threshold current. Journal of Physiology 414 (1989): 587-604.
- Cramer, J. A., Menachem, E. B., and French, J., Review of treatment options for refractory epilepsy: new medications and vagal nerve stimulation. Epilepsy Research 47 (2001): 17-25.
- Cull-Candy, S., Brickley, S. and Farrant, M. NMDA receptor subunits: diversity, development and disease. Current opinion in neurobiology 11(2001): 327-335.
- Daikin, Y. and Yudkoff, M. Compartmentation of brain glutamate metabolism in neurons and glia. Journal of Nutrition 130 (2000): 1026S-1031S.
- Das, S., Sasaki, Y. F., Rothe, T. et al. Increased NMDA current and spine density in mice lacking the NMDA receptor subunit NR2A. Nature 393 (1998): 377-381.
- Deckers, C. L. P., Genton, P., Sills, G. J., Schmidt, D., Current limitations of antiepileptic drug therapy: a conference review. Epilepsy Research 53 (2003): 1-17.
- Deutsch SI, Rosse RB, Billingslea EN, Bellack AS, Mastropaolo J. Modulation of MK-801-elicited mouse popping behavior by galantamine is complex and dose-dependent Life Sci. 73(18) (2003):2355-61
- Dingledine, R., Borges, K., Bowie, D., and Traynelis, S. F. The glutamate Receptor ion channels. Pharmacological Reviews 50 (1999): 7-61.
- Doherty, J. and Dingledine, R. The role of metabotropic glutamate receptors in seizures and epilepsy. Current Drug Target CNS Neurol Disord. 3 (2002): 251-260.
- Doi T, Ueda Y, Tokumaru J, Mitsuyama Y, Willmore LJ. Sequential changes in AMPA and NMDA protein levels during Fe(3+)-induced epileptogenesis. Brain Res Mol Brain Res. 92(1-2) (2001):107-14.
- Donevan, S. D. and McCabe, R. T. Conantokin G is an NR2B- selective competitive Antagonist of N-methyl-D-aspartate receptors. Molecular Pharmacology 58 (2000): 614-623.

- Dreifuss, F. E. and Fountain, N. B. Classification of epileptic seizures and the epilepsies and drugs of choice for their treatment. Eadie, M. J. and Vajda, F. J. E. In; Antiepileptic drugs: Pharmacology and therapeutics. 1-17. Germany: Springer-Verlag. 1999.
- Dumont, J. N. Oogenesis in *Xenopus laevis* (dandin) I. stages of oocyte development in laboratory maintained animals. J. Morphol. 136 (1972): 153-157.
- Durand, G. M., Gregor, P., Zheng, X., Uhl, G. R. and Zukin, R. S. Cloning of apparent splice variant of the rat N-methyl-D-aspartate NMDAR1 with altered sensitivity to polyamines and activators of protein kinase C. Proc. Natl. Acad. Sci. USA. 89 (1992): 9359.
- Ferrer-Montiel, A.V., Montal, M. Pentameric subunit stoichiometry of a neuronal glutamate receptor. Proc Natl Acad Sci U S A. 93(7) (1996): 2741-4.
- Foucaud B, Laube B, Schemm R, Kreimeyer A, Goeldner M, Betz H. Structural model of the N-methyl-D-aspartate receptor glycine site probed by site-directed chemical coupling. Journal of Biological Chemistry 278(26) (2003):24011-7.
- Franceschetti, S., Hamon B., and Heineman, U. The action of valproate on spontaneous epileptiform activity in the absence of synaptic transmission and on evoked changes in  $[Ca^{2+}]_o$  and  $[K^+]_o$  in the hippocampal slice. Brain Research. 386(1-2) (1986) :1-11.
- Franck, J., Pokorny, J., Kunkel, D., Schwartzkroin, P. Physiologic and morphologic characteristics of granule cell circuitry in human epileptic hippocampus. Epilepsia 36 (1995): 543-558.
- Fraser, S. P., Djamgoz, M. B. A. Current Aspects of the Neurosciences. Osborne, N. N. 267-315. USA:Macmillan Press.1992.
- Gandolfo, G., Romettino, S., Gottesman, C., Van Luijtelaer, G. Coenen, A., Bidard, J. N., and Lazdunski, M.,  $K^+$  channel openers decrease seizures in genetically epileptic rats. European Journal of Pharmacology 167 (1989): 181-183.
- Gean, P. W., Huang, C. C., Hung, C.R. and Tsai, J. J. Valproic acid suppresses the synaptic response mediated by the NMDA receptors in rat amygdala slices. Brain Res Bull. 33(3) (1994): 333-6.

- Green, T., Rogers, C. A., Contractor, A. and Heinemann, S. F. NMDA receptors formed by NR1 in *Xenopus laevis* oocytes do not contain the endogenous subunit XenU1. Molecular Pharmacology 61(2) (2002): 326-333.
- Griggs, R. C. Epilepsy. Carpenter, C. J., Griggs, R. C. and Loscalzo, J. In; CECIL Essentials of medicine 5<sup>th</sup>. 956-964. USA: W. B. Sanders Company. 2001.
- Gurdon, J. B., Lane, C. D., Woodland, H. R. and Maraix, G. Use of frog oocytes for the study of messenger RNA and its translation in living cells. Nature (Lond) 233 (1971): 177-182.
- Guzikowski, A. P., Cai, S. X., Espitia, S. A., Hawkinson, J. E. et al. Analogs of 3-hydroxy-1H-1-benzazepine-2,5-dione: Structure-activity relationship at N-methyl-D-aspartate receptor glycine site. Journal of Medicinal Chemistry 39 (1996): 4643-4653.
- Hayashi, Y., Hattori, Y., Moriwaki, A., Lu, Y. F. and Hori, Y. Increases in brain polyamine concentrations in chemical kindling and single convulsion induced by pentylenetetrazol in rats. Neurosci Lett. 149(1) (1993): 63-6.
- Hopkins, A., Epilepsy. In; Clinical Neurology: A modern approach. 1<sup>st</sup>. 186-210. USA: Oxford University Press. 1993.
- Huttner, J.E. and Bean, B.P., Block of N-methyl-D-aspartate-activated current by the anticonvulsant MK-801: Selective binding to open channels. Proc. Natl. Acad. Sci. 85 (1988): 1307-1311.
- Igarashi, K., Kashiwagi, K. Polyamines: mysterious modulators of cellular functions. Biochem Biophys Res Commun. 271(3) (2000): 559-64.
- Jacobs, M. P. et al., Future Directions for epilepsy research. Neurology 57, (2001): 1536-1542.
- Jahr, C. E. and Steven, C. F. Calcium permeability of the N-methyl-D-aspartate receptor channel in hippocampal neurons in culture. Proc. Natl. Acad. Sci. USA. 90 (1993): 11573-11577.
- Jenthet, S. Effects of N-(2-propylpentanoyl) urea on GABA<sub>A</sub> receptor in acutely dissociated reat hippocampal neurones. Master's Thesis. Chulaongkorn University, Thailand. 2002.
- Johanessen, C. U. and Johannessen, S. I. Valproate: past, present, and future. CNS Drug Rev. 9(2) (2003):199-216.

- Johnson, E. W., deLanorelle, N. C., Kim, J. H. et al. Central and peripheral benzodiazepine receptors: opposite changes in human epileptogenic tissue. Neurology 42 (1992): 811-815.
- Johnson, J. W. and Ascher, P. Glycine potentiates the NMDA response in cultured mouse brain neurons. Nature (Lond.) 325 (1987): 529-531.
- Johnson, T. D. Polyamines and cerebral ischemia. Prog Drug Res 50 (1998):193-258.
- Jones, O. T. Ca<sup>2+</sup> channels and epilepsy. European Journal of Pharmacology 447 (2002): 211-225.
- Kandel, E. R., Schwartz, J. H. and Jessell T. M. Seizures and epilepsy. In; Principles of neural sciences. 4<sup>th</sup>. 910-935. USA: McGraw-Hill. 2000.
- Katsuwada, T., Kashiwabuchi, N., Mori, H., Sakimura, E. Kushiya, K. Araki, H. Meguro, H. Masaki, T. Kumanishi, Arakawa M. and Mishina, M. Molecular diversity of the NMDA receptor channel. Nature (Lond.) 358 (1992): 36-41.
- Kelso, S. R., Nelson, Leonard J. P. Protein kinase C-mediated enhancement of NMDA currents by metabotropic glutamate receptors in *Xenopus* oocytes. Journal of Physiology. 449 (1992):705-18.
- Kemp, J. A., Leeson PD. The glycine site of the NMDA receptor--five years on. Trends Pharmacol Sci. 14(1) (1993): 20-5.
- Kemp, J. A. and McKernan, R., M. NMDA receptor pathways as drug target. Nature (Neuroscience Supplement 5) (2002): 1039-1042.
- Ko, G. Y., Brown-Croyts, L. M. and Teyler, T. J. The effects of anticonvulsant drugs on NMDA-EPSP, AMPA-EPSP, and GABA-IPSP in the rat hippocampus. Brain Res Bull. 42(4) (1997): 297-302.
- Kojima, N., Ishibashi, H., Obata, K. and Kandel, E. R. Higher seizure susceptibility and enhanced tyrosine phosphorylation of N-methyl-D-aspartate receptor subunit 2B in fyn transgenic mice. Learn Memory 5 (1998): 429-445.
- Khongsombat, O. Effect of N-(2'-propylpentanoyl-2-pyrrolidinone) and N-(2-propylpentanoyl) urea on neurons of cerebellar Purkinje cell in rats. Master's Thesis. Chulalongkorn University, Thailand. 1997.
- Kwan, P., Brodie, M. J., The effectiveness of first antiepileptic drug. Epilepsia 42, (2001): 1255-1260.

- Kwan, P., Sills, G. J., Brodie, M. J. The mechanisms of action of commonly used antiepileptic drugs. Pharmacol Ther. 90(1) (2001): 21-34.
- Lafaire, A. V. and Schwarz, W. J. Voltage dependence of the rheogenic Na<sup>+</sup>/K<sup>+</sup> ATPase in the membrane of oocytes of *Xenopus laevis* Membrane Biology 91(1986): 43-51.
- Laube, B., Kuhse, J. and Betz, H. Evidence for a tetrameric structure of recombinant NMDA receptors. Journal of Neuroscience 18 (1998): 2954-2961.
- Laurie, D. J. and Seeburg, P. H. Regional and developmental heterogeneity in splicing of the rat brain NMDAR1 mRNA. Journal of Neuroscience 14 (1994): 3180-3194.
- Leewanich, P. Neuropharmacological studies on extracts of medicinal plants, *Hunteria zeylanica*, and its main component, corymine. Doctoral's Thesis. Toyama Medical and Pharmaceutical University, Japan. 1998.
- Leonard, J. P. and Kelso, S. R. Apparent desensitization of NMDA responses in *Xenopus* oocytes involves calcium-dependent chloride current. Neuron 2 (1990): 52-60.
- Li-Smerin, Y. and Johnson, J. W. Kinetics of the block by intracellular Mg<sup>2+</sup> of the NMDA-activated channel in cultured rat neurons. Journal of Physiology (Lond.) 491 (1996): 121-135.
- Liu, H., Wang, H., Sheng, M., Jan, L. Y., Jan, Y. N. and Basbaum, A. I. Evidence for presynaptic N-methyl-D-aspartate autoreceptors in the spinal cord dorsal horn. Proc. Natl. Acad. Sci. USA 91 (1994): 8383-8387.
- Lotfis, J. M. and Janowsky, A. The N-methyl-D-aspartate receptor subunit NR2B: localization, functional properties, regulation, and clinical implications. Pharmacology and Therapeutics 97 (2003): 55-85.
- Lombardo, A. J., Kuzneick, R., Powers, R. E., Brown, G. B., Altered brain sodium channel transcript level in human epilepsy. Molecular Brain Research 35 (1996): 84-90.
- Loscher W. Basic pharmacology of valproate: a review after 35 years of clinical use for the treatment of epilepsy. CNS Drugs. 16(10) (2002) :669-94.

- Loscher, W. and Potschka, H. Role of multipledrug transporter in pharmacoresistance to antiepileptic drugs. The Journal of Pharmacology and Experimental Therapeutics 301 (2002): 7-14.
- Loscher, W. New vision in the pharmacology of anticonvulsion. European Journal of Pharmacology 342 (1998): 1-13.
- Loscher, W. and Schmidt, D. New horizons in the development of antiepileptic drugs. Epilepsy Research 50 (2002): 3-16.
- Lott, R. S. and McAuley, J. W. Seizure Disorder. Koda-kimble, M. A., Young, L. Y. In; Applied Therapeutic: The Clinical Uses of Drugs 7<sup>th</sup>. 52-1-52-41. USA: Lippincott Williams&Wikings. 2001.
- Lowenstein, D. H. Seizures and Epilepsy. Fanoi, A. S., Braunwald, E., Isselbacher. K. J., Wilson, J. D., Martin, J. B., Kasper, D. L., Hauser, S. L., Longo, D, L. In; Harrison's Principles of Internal Medicines 14<sup>th</sup>. Singapore: McGraw-Hill.1998.
- Luo, J., Wang, Y., Yasuda, R. P., Dunah, A. W., and Wolfe, B. B. The majority of N-methyl-D-aspartate receptor complexes in adult rat cerebral cortex contain at least three different subunits (NR1/NR2A/NR2B). Molecular Pharmacology 51 (1997): 79-86.
- Lynch, D. R., Guttman, R. P. NMDA receptor pharmacology: perspectives from molecular biology. Curr Drug Targets. 2(3) (2001):215-31
- Lynch., D. R. and Guttman, R. P. Excitotoxicity Perspectives based on N-methyl-D-Aspartate Receptor Subtypes. The Journal of Pharmacology and Experimental Therapeutics 300 (2002): 717-723.
- Madden, D., R. The structure and function of glutamate receptor ion channels. Nature Review Neuroscience 3 (2002): 91-101.
- Marino MJ, Conn PJ. Direct and indirect modulation of the N-methyl D-aspartate receptor. Curr Drug Target CNS Neurol Disord. 1(1) (2002):1-16.
- Matsumoto, H. and Ajmone-Marsan, C. Cortical cellular phenomena in experimental epilepsy: Interictal manifestations. Exp Neurol 9 (1964): 286-304.
- Matute, C., Alberdi, E., Ibarretxe, G. and Sonchez-Gomez, M. Excitotoxicity in glial cells. European Journal Pharmacology 447 (2002): 239-246.



- Mayer, M. L. and Westbrook, G. L. The physiology of excitatory amino acids in the vertebrate central nervous system. Prog Neurobiol. 28(3) (1987):197-276.
- McDonald, Garofalo, E., Hood, T. et al. Altered excitatory inhibitory aminoacid receptor binding in hippocampus of patients with temporal lobe epilepsy. Ann Neurol 29 (1991): 520-541.
- McLean, M. J. and Macdonald, R. L. Carbamazepine and 10,11-epoxycarbamazepine produce use- and voltage-dependent limitation of rapidly firing action potentials of mouse central neurons in cell culture. J Pharmacol Exp Ther. 238(2) (1986): 727-38.
- McNamara, J. O. Drug effective in the therapy of the epilepsies. In; Goodman and Gilman's the pharmacological basis therapeutics 10<sup>th</sup>. 521-547. USA: The McGraw-Hill. 2001.
- Medvedev AV. Epileptiform spikes desynchronize and diminish fast (gamma) activity of the brain. An "anti-binding" mechanism? Brain Res Bull. 58 (1) (2002): 115-28.
- Meesomboon, R. et al. Investigation of embryotoxicity of N-2-(propylpentanoyl) urea in developing rat embryos in vitro. Research Report. Faculty of Pharmaceutical Sciences, Chulalongkorn University, Thailand, 1997
- Meldrum, B. S. Glutamate as a neurotransmitter in the brain: Review of physiology and pathology. Journal of Nutrition 130 (2000): 1007S-1015S.
- Misra, C., Brickley, S. G., Farrant, M. and Cull-candy, S. G. Identification of subunits contributing to synaptic and extrasynaptic NMDA receptors in Golgi cells of the rat cerebellum. Journal of Physiology (Lond) 524 (2000): 147-162.
- Monyer, H., Sprengel, R., Schoepfer, R. et al. Heteromeric NMDA receptor: Molecular and functional distinction of subtypes. Science (Wach DC) 256 (1992): 1217-1221.
- Mori, H. and Mishina, M. Review: Neurotransmitter receptors VIII, structure and function of the NMDA receptor channel. Neuropharmacology 34 (1995): 1219-1237.
- Mori, H. and Mishina, M. Structure and function of the NMDA receptor channel. Neuropharmacology 34 (1995): 1219-1237.

- Moriyoshi, K., Masu, M., Ishii, T., Shigemoto, R., Mizuno, N., Nakanishi, S. Molecular cloning and characterization of the rat NMDA receptor. Nature 354 (1991): 31-37.
- Murashima, Y. L., Yoshii, M. Suzuki, J., Ictogenesis and epileptogenesis in EL mice. Epilepsia 43 (2002): 130-135.
- Najm, I. Ying, Z., Janigro, D., Mechanisms of Epileptogenesis. Neurologic Clinics for North America 19(2) (2001): 237-250.
- Najm, I., Ying, Z., Babb, T. et al. NMDA receptor ZA/B subtype differential expression in human cortical dysplasia: Correlation with in vivo epileptogenicity. Epilepsia 41 (2000): 971-976.
- Narita, M., Aoki, T. and Suzuki, T. Molecular evidence for the involvement of NR2B subunit containing N-ethyl-D-aspartate receptors in the development of morphine induced place preference. Neuroscience 101 (2000): 601-606.
- Nishi, M., Hinds, H., Lu, H., Kawata, M. and Hayashi, Y. Motoneuron-specific expression of NR3B, a novel NMDA-type glutamate receptor subunit that works in a dominant-negative manner. The Journal of Neuroscience 21 (2001):1-6.
- Ozawa, S., Kamiya, H. and Tsuzuki, K. Glutamate receptors in the mammalian central nervous system. Progress in Neurobiology 54 (1998): 581-618.
- Parsons, C. G., Danysz, W., Quack, G. et al. Novel systemically active antagonist of the glycine site of the N-methyl-D-aspartate receptor: Electrophysiological, biochemical and behavioral characterization. The Journal of Pharmacology and Experimental Therapeutics 283 (1997): 1264-1275.
- Patchamart, W. Hepatotoxicity of N-(2-propylpentanoyl) urea in rats. Master's Thesis. Chulalongkorn University, Thailand, 1996.
- Perucca, E. Clinical Pharmacology. Epilepsy. In; McGraw-Hill Clinical Medicines series. 97-113. UK: McGraw-Hill International.2000.
- Perucca E. Pharmacological and therapeutic properties of valproate: a summary after 35 years of clinical experience. CNS Drugs. 16(10) (2002) :695-714.
- Perucca, E. Current trends in antiepileptic drug therapy. Epilepsia 44, suppl 4 (2003): 41-47.

- Pongs, S. O., Voltage-gated potassium channels: from by hyperexcitability to excitement. FEBS Letters 452 (1999): 31-35.
- Porter, R. J. and Meldrum, B. S. Antiseizure drug. Kazung, B. G. In; Basic & Clinical Pharmacology. 8<sup>th</sup>. 395-418. USA: McGraw-Hill. 2001.
- Porter, R. J. and Rogawski, M. A., New antiepileptic drugs from serendipity to rational discovery. Epilepsia 33 (suppl. 1) (1992): S1-S6.
- Preistley, H. K., Loughton, P., Meyer, J., Le Bourdelles, B., Kerby, J. and Whiting, P. J. Pharmacological properties of recombinant human N-methyl-D-aspartate receptor comprising NR1a/NR2A and NR1a/NR2B subunit assemblies expressed in permanently transfected mouse fibroblast cells Molecular Pharmacology 48 (1995): 841-848.
- Premkumar, L. S. and Auerbach., A. Stoichiometry of recombinant N-methyl-D-aspartate receptor channels inferred from single-channel current patterns. Journal of Gen Physio 110 (1997): 485-502.
- Prince, D. A., Physiological mechanisms of focal epileptogenesis. Epilepsia 25 (suppl.) (1985): S3-S14.
- Princivalle AP, Duncan JS, Thom M, Bowery NG. Studies of GABA(B) receptors labelled with [(3)H]-CGP62349 in hippocampus resected from patients with temporal lobe epilepsy. British Journal Pharmacology 136 (8) (2000):1099-106.
- Qian, A. and Johnson, J. W. Channel gating of NMDA receptors Physiol Behav 77(4-5) (2002):577-82.
- Ragsdale, D. S. and Avoli, M., Sodium channels as molecular targets for antiepileptic drugs. Brain Research Review 26 (1998): 16-28.
- Raol, Y. H., Lynch, D. R. and Brooks-Kayal, A. R. Role of excitatory amino acids in developmental epilepsies. Ment Retard Dev Disabil Res Rev. 7(4) (2000): 254-60.
- Regesta, G. and Tangenelli, P. Clinical aspects and biological bases of drug-resistant epilepsies. Epilepsy Research 34 (1999): 109-122.
- Rogawski, M. A. KCNQ2/KCNQ3 K<sup>+</sup> channels and the molecular pathogenesis of epilepsy: implications for therapy. Trends Neuroscience 23 (2000): 393-398.
- Rosenmund, C., Stern-Bach, Y. and Stevens, C. F. The tetrameric structure of a glutamate receptor channel. Science (Wash DC) 280 (1998): 1596-1599.

- Rostock, A. Tober, C., Rundfeldt, C., Bartsch, R., Engel, J., Polymeropoulos, E. E., Kutscher, B., Loscher, W., Honack, D., White, H. S., and Wolf, H. H. D-23129: a new anticonvulsant with a broad spectrum activity in animal models of epileptic seizures. Epilepsy Research 23 (1996): 211-223.
- Ruppersberg, J. P., Mosbacher, J., Gunther, W., Schoepfer, R. and Fakler, B. Studying block in cloned N-methyl-D-Aspartate (NMDA) receptors. Biochem. Pharmac. 46 (1993): 1877-1885.
- Saisorn, B., Patarapanich, C. and Janwittayanuchit, W. Synthesis of monoureide analogs of valproic acid. Thai Journal of Pharmaceutical Sciences 16(2) (1992): 145-50.
- Smart, T. G., Xie, X., Krishek, B. J. Modulation of inhibitory and excitatory amino acid receptor ion channels by zinc. Progress in Neurobiology 42 (1994): 393-441.
- Schachter, SC. Epilepsy: Quality of Life and Cost of Care Epilepsy Behav. 1(2) (2000):120-127.
- Sigel, E. Use of *Xenopus* oocytes for the functional expression of plasma membrane proteins. J Membr Biol. 117(3) (1990):201-21.
- Sills, G. J., Butler, E., Thompson, G. G. and Brodie, M. J. Vigabatrin and tiagabine a pharmacologically different drug. A pre-clinical study. Seizure 8 (1999): 404-411.
- Soloviev, M. M. and Barnard, E. A. *Xenopus* oocytes express a unitary glutamate receptor endogenously. Journal of molecular biology 273 (1997): 14-18.
- Sooksawate, T. Anticonvulsant effects of n-(2-propylpentanoyl) Urea. Master's Thesis. Chulalongkorn University, Thailand.1995
- Sperk, G., Schwarzer, C., Tsunashima, K., Kandlhofer, S. Expression of GABA<sub>A</sub> receptor subunits in the hippocampus of the rat after kainic acid-induced seizures. Epilepsy Research 32 (1998): 129-139.
- Stamler, J. S., Toone, E. J., Lipton, S. A. and Sucher, N. J. (S)NO signal: Translocation, regulation and a consensus motif. Neuron 18 (1997): 691-696.
- Standaert, D.G., Testa, C.M., Young A.B., and Penney, J.B. Jr. Organization of N-methyl-D-aspartate glutamate receptor gene expression in the basal ganglia of the rat. J. Comp. Neurol. 343 (1994): 1-16.

- Stefani, A., Spadoni, F. and Bernardi, G. Voltage-activated calcium channels: targets of antiepileptic drug therapy? Epilepsia 38(1997): 959-965.
- Steinhoff, B. J., Hirsh, E., Mutani, R., Nakken, K. O. The ideal characteristics of antiepileptic therapy.: an overview of old and new AEDs. Acta Neurologica Scandinavica 107, (2003):87-85.
- Steinhouser, C. and seifert, G. Clial membrane channels and receptors in epilepsy: impact for generation and spread of seizure activity. European Journal of Pharmacology 477 (2002): 227-237.
- Stephenson, F. A. Subunit characterization of NMDA receptors. Curr Drug Targets. 2(3) (2001):233-9.
- Sucher, N. J., Awobuluyi, M., Choi, Y. B. and Lipton, S. A. NMDA receptors: from genes to channels. Trends in Pharmacological Science 17 (1996): 348-355.
- Sullivan, J. M., Traynelis, S. F., Chen, H. S., Escobar, W., Heinemann, S. F. and Lipton, S. A. Identification of two cysteine residues that are required for redox modulation of the NMDA subtype of glutamate receptor. Neuron 13 (1994): 929-936.
- Tantisira, B., Tantisira, M. H., Patarapanich, C., Sooksawate, T. and Chungam, T. Preliminary evaluation of the anticonvulsant activity of valproic acid analog: N-(2-propylpentanoyl)urea. Research and Community in Molecular Pathology and Pharmacology 97(2)(1997): 151-163.
- Taverna, S., Sancini, G., Magtegazza, M., Freanceschetti, S. and Avanzini, G. Inhibition of transient and persistent Na<sup>+</sup> current fractions by the new anticonsulant? topiramate. The Journal of Pharmacology and Experimental Therapeutics 288 (1999): 960-968.
- Thavendiranathan, P., Chow, C., Cunnane, S. and McIntyre Burnham, W. The effect of the 'classic' ketogenic diet on animal seizure models. Brain Research 959(2)(2003):206-13.
- Traynelis SF, Burgess MF, Zheng F, Lyuboslavsky P, Powers JL. Control of voltage-independent zinc inhibition of NMDA receptors by the NR1 subunit. J Neurosci. 18(16) (1998):6163-75.
- Trist, D. G. Excitatory amino acid agonists and antagonist: pharmacology and therapeutic application. Pharm Acta Helv 74 (2000):221-229.

- Vissel, B., Krupp, J. J., Heinemann, S. F., Westbrook, G. L. Intracellular domains of NR2 alter calcium-dependent inactivation of N-methyl-D-aspartate receptors. Molecular Pharmacology 61(3) (2002):595-605
- Vreugdenhil, M. Faas, G. O., Wadman, W. H., Sodium currents in isolated rat CAI neurons after kindling. Neuroscience 86 (1999): 99-107.
- Wagner, C. A., Friedrich, B., Setiawan, I., Lang, F. and Bruer, S. The use of *Xenopus laevis* oocytes for the functional characterization of heterologously expressed membrane proteins. Cellular Physiological Biochemistry 10 (2000): 1-12.
- Wang, H-C., Beer, B., Sassano, D., Blume, A. J. and Ziai, M. R. Minireview: Gene expression in *Xenopus* oocytes. Int. J. Biochem. 23 (1991)271-276.
- Weber, W. M. Topical Review: Endogenous ion channels in oocytes of *Xenopus laevis*: recent development. The Journal of Membrane Biology. 170 (1999a): 1-12.
- Weber, W. M. Review: Ion currents of *Xenopus Laevis* oocytes: state of the art. Biochemica et Biophysica Acta. 1421 (1999b): 213-233.
- Wickenden, A. D. Review: Potassium channels as anti-epileptic drug targets. Neuropharmacology 43 (2002): 1055-1060.
- Wiebe, S., Nicolle, M. W. Recent developments in neurology. BMJ. 16 (2002):324(7338):656-60
- Williams, K. Separating dual effects of zinc at recombinant N-methyl-D-Aspartate receptors. Neuroscience Letter 215 (1996): 9-12.
- Williams, K. Interactions of polyamines with ion channels. Biochem J. 325 (Pt 2) (1997): 289-97.
- Wollmuth, L. P., Kuner, T. and Sakmann, B. Intracellular Mg<sup>2+</sup> interacts with structural determinants of the narrow constriction contributed by the NR1-subunit in the NMDA receptor for channel. Journal of Physiology (Lond.) 506 (1998): 33-52.
- Wollmuth, L. P., Kuner, T., Seeburg, P. H., and Sokmann, B. Differential contribution of the NR1- and NR2A- subunits to the selectivity filter of recombinant NMDA receptor channels. Journal of Physiology (Lond) 491 (1996): 779-797.


- Wood, M. W., VanDongen, H. M. A. and VanDongen, A. M. J. Structural conservation of ion conduction pathways in K channels and glutamate receptor. Proc. Nat. Acad. Sci. USA 92 (1995): 4882-4886.
- Yamaguchi, S. and Rogawski, M. A. Effects of anticonvulsant drugs on 4-aminopyridine-induced seizures in mice. Epilepsy Research 11 (1992): 9-16.
- Yamakura, T. and Shimoji, K. Subunit and site-specific pharmacology of the NMDA receptor channel. Progress in Neurobiology 59 (1999): 279-298.
- Ying, Z., Babb, T., Camair, Y., et al. Induced expression of NMDAR2 Proteins and differential expression of NMDAR1 splice variants in dysplastic neurons of human epileptic neocortex. J. Neuropathol Exp. Neurol 57 (1998): 46-47.
- Zeise, M L., Kasparaow, S. and Zeiglgansberger W. Valproate suppresses N-methyl-D-aspartate-evoked, transient depolarizations in the rat neocortex in vitro. Brain Research. 544(2) (1991): 345-8.



ศูนย์วิทยทรัพยากร  
จุฬาลงกรณ์มหาวิทยาลัย

## VITAE

Miss Penphimon Ponsup was born on June 19, 1977 in Chumphon, Thailand. She received her Bachelor of Science in Pharmacy with first class honors in 1999 from the Faculty of Pharmaceutical Sciences, Chulalongkorn University, Thailand. Since her graduation, she has become a faculty member of the Department of Physiology, Faculty of Pharmaceutical Sciences, Chulalongkorn University.



ศูนย์วิทยทรัพยากร  
จุฬาลงกรณ์มหาวิทยาลัย