

Reference

ภาษาไทย

- กมล สวัสดิ์มงคล และ คณะ การประชุมวิชาการเรื่อง ฟั้หะลายโจร
กรมวิทยาศาสตร์การแพทย์, 2535, นนทบุรี
- กรมวิทยาศาสตร์การแพทย์. ข้อมูลฟั้หะลายโจร Andrographis
paniculata (Burm.f.) Nees. กรุงเทพมหานคร : สำนักงาน
ปลัดกระทรวงสาธารณสุข, 2529
- กองวิจัยและพัฒนามสมุนไพร กรมวิทยาศาสตร์การแพทย์ คู่มือสมุนไพร
สาธารณสุขมูลฐาน กระทรวงสาธารณสุข, 2533
- คณิต สุวรรณบริรักษ์ และ ชัยโย ชัยชาญพิพุทธ. น้ำลายฟั้หะ,
ฟั้หะลายโจร. วารสารสมาคมสมุนไพรแห่งประเทศไทย.
7 (มกราคม-มีนาคม 2534) : 3-9.
- ธิดารัตน์ ปลื้มใจ และ นางฤดี สิทธิสมวงศ์ การประชุมวิชาการ เรื่อง
ฟั้หะลายโจร กรมวิทยาศาสตร์การแพทย์, 2533, นนทบุรี
- นันทกาญจน์ มหาวีรวัฒน์. การศึกษาปริมาณสารโคเทออร์ป็นอยด์จากใบ
ฟั้หะลายโจรที่เก็บแต่ละเดือน : วิทยานิพนธ์ปริญญาโทมหาบัณฑิต
ภาควิชาเภสัชเคมี บัณฑิตวิทยาลัย จุฬาลงกรณ์มหาวิทยาลัย, 2533.
- ปัญญาวงศ์ ธนังกุล และ ชัยโย ชัยชาญพิพุทธ. การศึกษาทางคลินิกของ
สมุนไพรฟั้หะลายโจรในโรคอุจจาระร่วงและบิดแบคทีเรีย
รามาธิบดีเวชสาร 8(2528) : 57-61.
- ผจงศิลป์ เฟื่องมาก. ผลของ 14 คืออกซี-11-12-16คีไฮดรแอนโดร-
กราฟโพลีคต่อการหดตัวของกล้ามเนื้อลำไส้เล็กนอกร่างกาย
วิทยานิพนธ์ปริญญาโทมหาบัณฑิต ภาควิชาเภสัชวิทยา บัณฑิตวิทยาลัย
จุฬาลงกรณ์มหาวิทยาลัย, 2531.

เพชรรัตน์ พงศ์จรรยากุล. กฤทธิ์ของแอนโดรกราฟโฟไลด์ต่อกล้ามเนื้อเรียบของ
ลาเล้ง วิทยานิพนธ์ปริญญาโทมหาบัณฑิต ภาควิชาเภสัชวิทยา บัณฑิตวิทยาลัย
จุฬาลงกรณ์มหาวิทยาลัย, 2530.

ศิริมา พรสุวัฒนา, ประสาน ชธรรมอุภกรณ และ อума กิติยานี ทดสอบฤทธิ์การ
ป้องกันและรักษาแผลกระเพาะอาหารของสมุนไพรรักษาหลายโรคและ
เบส้าน้อย ไทยเภสัชสาร 14(2532) : 34-43.

ภาษาอังกฤษ

- Abeyssekera, A.M., Desilva, K.T.D., Silva, W.S.J. and
Ratnayake, S. ^1H and ^{13}C -NMR spectral analysis
of Andrographolide. Fitoterapia 59(1988) : 501-504.
- Allison, A.J., Butcher, D.N., Connolly, J.D. and Overton,
K.H. Paniculides A, B and C Bisabolenoid Lactose
from Tissue Cultures of Andrographis paniculata.
Chemical Communication (1968) 1943.
- Balmain A. and Connolly J.D. Minor diterpenoid constituents
of Andrographis paniculata Nees. J. Chem. Soc.
Perkin J. (1973) : 1247-1251.
- Barua, A.K., Biswas, S., Partra, A., Basu, K., Panda, S.K.
and Ghosh, A. Phloganthoside, A diterpene lactose
glucoside from Phlogacanthus thyrsiflorus.
Phytochemistry (1987) : 491-492.

- Bhatt, R.K. and Sabata B.K. Structure of tinosporide : It's identity in all respects with jateorin, (15, 25, 3R, 4R, 5R, 8R, 10R, 12S)-4-hydroxy-2,3 : 15, 16-diepoxycleroda-13(16), 14-dieno-17,12:18, 1-biscarbolactone. Indian Journal of Chemistry (June 1990) : 521-524.
- Biswas, K.M., Ali, M.E. and Choudhury S.A. Investigations on Andrographis paniculata Nees, Part VI Root flavones and their structure. Pak. J. Sci. Ind. Res. 15(1972) : 33-36.
- Cambie, R.C. a Norditerpene lactone from Podocarpus halli. Phytochemistry (1973) : 2057.
- Cava, M.P. Chan, W.P., Haynes, L.J. and Johnson, L.F. The structure of Andrographolide. Tetrahedron 18(1962) : 397-403.
- Cava, M.P., Chan, W.R., Stein, R.P. and Willis, C.R. Andrographolide; further transformation and stereochemical evidence, the structure of isoandrographolide. Tetrahedron (1965) : 1617-2631.
- Chan, W.R., Taylor, D.R. and Willis, C.R. The structure of Neoandrographolide : A diterpene glucoside from Andrographis paniculata Nees. Tetrahedral Letter. 46(1968) : 4803-4806.
- Chan, W.R., Taylor, D.R., Willis, C.R., Bodden, R.L. The Structure and stereochemistry of Neoandrographolide, A diterpene glucoside from Andrographis paniculata Nees. Tetrahedron 27(1971) : 5081-5090.

- Changqui H., Bingnan, Z., Nan C.P. Isolation and Structure of two new diterpenoid glucosides from Andrographis paniculata Nees. Acta Pharmaceutica Sinica (1982) : 435-440.
- Chakravarty, A.K. and Pal, B.C. Structure of a new diterpene lactone from Gelonium multiflorum by 2-D NMR. Indian Journal of Chemistry (January 1991) : 3-6.
- Chakravarti D. and Chakravarti R.N. Andrographolide Part I. J. Chem. Soc. (1952) : 1697-1700.
- Chaudhuri, S.K. Influence of Andrographis paniculata Nees. on bile flow and Hexabarbitone sleeping in Experimental Animals. Indian J. Exp. Biol. 16(1978) : 830-832.
- Choudhury B.R. and Poddar M.K. Andrographolide and Kalmegh (Andrographis paniculata) Extract : Effect on Intestinal Brush Border Membrane - Bound Hydrolases. Meth and Find Exptl Clin Pharmacol 7(1985) : 617-621.
- _____. Andrographolide and Kalmegh (Andrographis paniculata) Extract : in Vivo and in Vitro Effect on Hepatic Lipid Peroxidation. Meth and Find Exptl Clin Pharmacol 6(1984) : 481-485.
- Cuevas, G., Collera, O., Garcia, F., Cardenas, J., Maldonado E. and Ortega A. Diterpenes from Salvia breviflora. Phytochemistry (1987) : 2019-2021.

- Dasgupta, B., Burke, B.A. and Stuart, K.L. Biflavonoids, Norditerpenes and a Nortriterpines from Podocarpus urbanii. Phytochemistry (1981) : 153-156.
- Delgado, G., Alvaree, L. and Vivar, A.R.D. 15 - oxo-zoapatlin , A diterpene lactone from Viguiera maculata. Phytochemistry (1984) : 2674-2675.
- Duddeck, H. and Dietrich, W. Structure Elucidation by Modern NMR. New York : Springer, 1989.
- Dultta, A. and Sakul, N.C. Filaricidal Properties of a wild herb Andrographis paniculata. J of Helminthology 56(1982) : 81-84.
- Fujita, T, Fujitani R., Takeda Y., Takaishi Y., Yamada T., Kido M. and Miura J. On the diterpenoids of Andrographis paniculata : X-ray Crystallographic Analysis of, Andrographolide. and structure determination of new minor diterpenoid. Chem. Pharm. Bull. 32(1984) : 2117-2125.
- Garcia, L.L., Kintanar, Q.L., Fojas, F.R., Sison, F.M., chua, N.B. and Villanniva B.A. Pharmacological Studies on the leaves of Andrographis paniculata Nees, Plant Growth in the Philippines . Med Aromat. Plants Abstr. 6(1981) : 24-25.
- Gupta S., Choudhury MA. and Yadava J.N.S. Antidiarrheal Activity of Diterpenes of Andrographis paniculata (Kalmegh) against Escherichia coli Enterotoxin in in vivo Models. Int J. Crude Drug Res. 28(1990) : 273-283.

- Gupta, K.K., Taneja, S.C., Dhar, K.L. and Atal C.K. Flavonoids of Andrographis paniculata. Phytochemistry 2(1983) : 314-315.
- Govin, T.R., Pai B.R., Srinivasan M. and Kalyanaraman P.S. Chemical Investigation of Andrographis paniculata. Indian J. Chem 3(1969) : 306.
- Govindachari, T.R. and Parthasarathy, P.C. Chemical investigation of Andrographis serpyllifolia : isolation and structure of serpyllin, a new flavone. Tetrahedron (1968) : 7027-7031.
- Hunaman J.B., Bhatt R.K., Sabata B.K. A diterpenoid Furanolactone from Tihospora cordifolia. Phytochemistry 25(1986) : 1677-1680.
- Hooker, J.D., The flora of British India. Vol 4. : Bissen Sing Mahendra Pal Singh Delhi, 1973.
- Hu Chang-qi and Zhou, B.N. Isolation and structure of two new diterpenoid glucosides from Andrographis paniculata Nees. Acta Pharmaceutica Sinica Vol XVII (June 1982) : 435-440.
- Jajuddin, Snanid A. and Tarig M. Antiinflammatory Activity of Andrographis paniculata Nees. Med Aromat. Plants Abstr. 6(1987) : 481.
- Jalal MA.F., Overton, K.H. and Rycroft, D.S. Formation of three new flavones by differentiating callus cultures of Andrographis paniculata. Phytochemistry 18(1989) : 149-151.

- Kizu H., Imoto, Y., Tomimon T., Tsubono K., Kadota S. and Kikuchi T. Structure of scuterivulactone D Determined by two dimensional NMR spectroscopy. A new diterpenoid from a chinese crude drug "Ban zhi lian". (Scutellaria rivularis wall) Chem. Pharm. Bull. 35(1987) : 1656-1659.
- Kuroyanagi M., Sato, M., Ueno A. and Nishi K. Flavonoid from Andrographis paniculata. Chem Pharm. Bull. 35(1987) : 4427-4435.
- Lin, Y.L., Kuo, Y.H., Ching, M.C. and Wang, Y. Structure of scutellones D and E Determined From X-ray diffraction, spectral and Chemical Evidence. Neochlerodane Type Diterpenoid from Scutellaria rivularis wall. Chem. Pharm Bull (1988) : 2642-2646.
- Labres J.M., Viladomat, F., Bastida, J., Codina, C., Serrano, M., Rubiralta, M. and Feliz, M. Two alkaloids from Narcissus requienii. Phytochemistry (1986) : 1453-1459.
- Marston A. and Hostettmann K. Modern Separation Methods Natural Product Report (1991) : 391-413.
- Martin, G.E. and Zektzer, A.S. Two dimensional NMR Methods for Establishing Molecular connectivity New York : VCH, 1988.
- Matlin, S.A., Bittner M. and Silva M. Lignan and Norditerpene dilactone constituents of Podocarpus saligna. Phytochemistry (1984) : 2867-2870.

- Matlin, S.A. and Prazeres M.A. Isolation and Structure Determination of Norditerpene Dilactones from Podocarpus saligna D. Don J. Chem. Soc. Perkin trans I (1982) : 2589-2593.
- Matsuda, T., Kuroyanagi M.; Sugiyama, S., Ueno, A. Studies on the cell differentiation inducers of Andrographis paniculata Proceeding of the 33rd symposium on the Chemistry of Natural Product OSAKA, 1991.
- McMichael, K.D., Overton, K.H. , and Picken, D.J. Tranesterification of Farnesol Mediated by a lipase from Andrographis paniculata tissue cultures. Phytochemistry (1977) 1290-1292.
- Medforth, C.J., Chang, R.S. , Chen, G.Q. , Olmstead, M.M. and Smith, K.M. A conformation study of Diterpenoid lactones Isolated from the Chinese Medicinal herb Andrographis paniculata. J. Chem. Soc. Perkin. trans 2(1991) : 1011-1016.
- Moktader A. and Guha-Sircar. On the bitter principle from Andrographis paniculata Nees Part I. J. Indian Chem. Soc. (1932) 333-338.
- Ohmoto, T., Koike, K., Fukuda, H., Mitsunaga, K., Kagai, K., Kawai, T. and Sato, T. Studies on the constituents of Picrodendron baccatum Growing in Indonesia : Structure of a Norditerpene lactone, Picrodendrin A. Chem Pharm Bull 37(1989) : 1805-1807.

- Overton, K.H. and Roberts, F.M. Biosynthesis of trans; trans and cis, trans-Farnesols by soluble enzyme for Tissue cultures of Andrographis paniculata Biochem Journal 144(1974) : 585-592.
- Rahman, A.U., Lavi, K.A., Sultana, A.T. and shameel, M. A. diterpenoid lactone from Aplysia juliana. Journal of Natural product (May-June 1991) : 886-888.
- Sanders, J.K.M. and Hunter, B.K. Modern NMR Spectroscopy A Guide for Chemists. 2nd ed Oxford : Oxford University, 1988.
- Satyanarayana, D., Mythirayee, C., Krishna murthy-V Polyphenols of Andrographis paniculata Nees. Leather Sci. 25(1978) : 250-251.
- Still, W.C., Kahhn, M. and Mitra, A. Rapid Chromatographic Technique for Preparative Separations with Moderate Resolution J. Org. Chem (1978) : 2923-2925.
- Suri O.P., Dhar, K.L., Altal, C.K., Kawasaki, T., Miyahara, K. and Kavano, S. High resolution NMR and X-ray Crystallography data of caudicifolin from Euphorbia acaulis. Phytochemistry 25(1986) : 1411-1413.
- Talapatra, S.K., Karmacharya B., De, S.C. and Talapatra, B. Chemical investigation of some medicinal plants of Nepal Indian Journal of Chemistry. (April 1989) : 356-357.

- Teresa, J.D.P., Urones, J.G., Basabe, P., Carrillo, H., Hunoz, M.A.G. and Marcos, I.S. Diterpenoid of Halimium viscosum. Phytochemistry (1985) : 791-794.
- Toyota, M., Nagashima, F and Asakawa, Y. Striatane and pinguisane-type sesquiterpenoids and phytane type diterpenoid from the liverwort Porella navicularis. Phytochemistry (1989) : 1661-1665.
- Weiming, C. and Xiaotian L. "Deoxyandrographolide-19 β -D-glucoside from the leaves of Andrographis paniculata. Planta Medica 45(1982) : 245-246.
- Zhengmu M. Isolation and identification of a new glucose : bicyclic diterpenoid lactone from Andrographis paniculata Nees. Nanging Yaoxupvuan Xupbau (1981) : 15-20.

ศูนย์วิทยทรัพยากร
จุฬาลงกรณ์มหาวิทยาลัย



ศูนย์วิทยทรัพยากร
จุฬาลงกรณ์มหาวิทยาลัย




Figure 1 *Andrographis paniculata* Nees

Thai name : Nam Laai Pangphon, Fah Talai Joan.




Figure 18 UV spectrum of compound C-2 in Chloroform




Figure 19 IR spectrum of compound C-2

ศูนย์วิทยทรัพยากร
จุฬาลงกรณ์มหาวิทยาลัย




Figure 20 Mass spectrum of compound C-2

ศูนย์เภสัชวิทยา
จุฬาลงกรณ์มหาวิทยาลัย




Figure 21 Proton spectrum of compound C-2 in CDCl₃




Figure 22 H,H COSY spectrum of compound C-2 in CDCl₃

ศูนย์วิทยทรัพยากร
จุฬาลงกรณ์มหาวิทยาลัย 3




Figure 23 H,H COSY Expanded scale of compound C-2 in CDCl_3




Figure 24 Carbon-13 decoupling spectrum of compound C-2 in CDCl_3




Figure 25 DEPT-135 spectrum of compound C-2 in CDCl₃

ศูนย์วิจัยเภสัชกร
จุฬาลงกรณ์มหาวิทยาลัย




Figure 27 UV spectrum of compound C-3 in Methanol




Figure 28 IR spectrum of compound C-3




Figure 29 Mass spectrum of compound C-3

ศูนย์วิทยาศาสตร์
จุฬาลงกรณ์มหาวิทยาลัย




Figure 30 Proton spectrum of compound C-3 in DMSO-d₆




Figure 31 ^1H , ^1H COSY spectrum of compound C-3 in DMSO-d_6




Figure 32 H,H COSY expanded scale of compound C-3 in DMSO-d6




Figure 33 Carbon-13 decoupling spectrum of compound C-3 in DMSO-d6




Figure 34 DEPT-135 spectrum of compound C-3 in DMSO-d6




Figure 35 DEPT-90 spectrum of compound C-3 in DMSO-d6




Figure 36 C,H COSY spectrum of compound C-3 in DMSO-d₆

ศูนย์วิจัยทรัพยากร
จุฬาลงกรณ์มหาวิทยาลัย




Figure 37 Proton spectrum of glucose in DMSO-d₆



Figure 38 H,H COSY spectrum of glucose in DMSO-d₆




Figure 39 Carbon-13 decoupling spectrum of glucose in DMSO-d6




Figure 40 DEPT 90 spectrum of glucose in DMSO-d6

ศูนย์วิทยทรัพยากร
จุฬาลงกรณ์มหาวิทยาลัย




Figure 42 UV spectrum of compound C-4 in Methanol




Figure 43 IR spectrum of compound C-4

ศูนย์วิทยทรัพยากร
จุฬาลงกรณ์มหาวิทยาลัย




Figure 44 Mass spectrum of compound C-4




Figure 45 Proton spectrum of compound C-4 in DMSO-d₆




Figure 46 H,H COSY spectrum of compound C-4 in DMSO-d₆




Figure 47 H,H COSY expanded scale spectrum at 2.8-5.2 ppm of compound C-4




Figure 48 H,H COSY expanded scale spectrum at 0.4-2.7 ppm of compound C-4




Figure 49 Carbon-13 decoupling spectrum of compound C-4 in DMSO-d₆




Figure 50 DEPT 135 spectrum of compound C-4 in DMSO-d6




Figure 51 DEPT 90 spectrum of compound C-4 in DMSO-d6




Figure 52 C,H COSY spectrum of compound C-4 in DMSO-d6




Figure 54 UV spectrum of compound C-5 in Methanol




Figure 55 IR spectrum of compound C-5




Figure 56 Mass spectrum of compound C-5




Figure 58 ¹H, ¹H COSY spectrum of compound C-5 in DMSO-d₆




Figure 59 Carbon-13 decoupling spectrum of compound C-5 in DMSO-d6




Figure 60 DEPT 90 spectrum of compound C-5 in DMSO-d6




Figure 62 UV spectrum of compound C-6 in Methanol




Figure 63 IR spectrum of compound C-6




Figure 64 Mass spectrum of compound C-6




Figure 65 Proton spectrum of compound C-6 in DMSO-d₆




Figure 66 H,H COSY spectrum of compound C-6 in DMSO-d₆




Figure 67 Carbon-13 decoupling spectrum of compound C-6 in DMSO-d₆




Figure 68 DEPT 135 spectrum of compound C-6 in DMSO-d₆




Figure 69 .DEPT 90 spectrum of compound C-6 in DMSO-d6



VITA

Miss Srisomporn Preeprame was born on April 21, 1966 in Ubonrajchathani. She graduated with Bachelor Degree of Science in Pharmacy with the second class honors in 1989 from Faculty of Pharmacy, Khon Khaen University.



ศูนย์วิทยทรัพยากร
จุฬาลงกรณ์มหาวิทยาลัย