

เอกสารอ้างอิง

มีชัย ศรีส. 2530. ประสาทกายวิภาคศาสตร์. 1500 เล่ม. พิมพ์ครั้งที่ 3. กรุงเทพมหานคร :
สำนักพิมพ์สินประสิทธิการพิมพ์.

ราตรี สุคทรง. 2532. ประสาทสรีรวิทยา. 1000 เล่ม พิมพ์ครั้งที่ 1. กรุงเทพมหานคร :
สำนักพิมพ์จุฬาลงกรณ์มหาวิทยาลัย.

อนันต์ ศรีเกียรติชจร และ กัมมังส์ พันธุ์มจินดา. 2533. ภาวะกล้ามเนื้อหดเกร็ง.
จุฬาลงกรณ์เวชสาร 34 : 311-322.

Aoyama, M., Hongo, T., Kudo, N. 1973. An uncrossed ascending tract
originating from below Clarke's column and conveying group I
impulses from the hindlimb muscles of the cat. Brain Res.
62 : 237-241.

Baldisserra, F., Hultborn, H., Illert, M. 1981. Integration in spinal
neuronal systems. In V.B. Brooks (ed.), Hand book of Physiology,
509-595. Washington D.C.

Brink, E., Harrison, P.J., Jankowska, E., McCrea, D.A. and Skoog, B.
1983a. Postsynaptic potential in a population of motoneurons
following activity of single interneurons. J Physiol. 343 :
341-359.

_____. 1983b. Inhibit interactions between interneurons in reflex pathways from group Ia and Ib afferent in the cat. J Physiol. 343 : 361-373.

Cooper, S. and Sherrington, C.S. 1940. Gower's tract and spinal border cells. Brain. 63 : 123-134.

Cumming, J.F. and Petras, J.M. 1977. The origin of spinocerebellar pathways. I. The nucleus cervicalis centralis of the cranial cervical spinal cord. J Comp Neurol. 173 : 655-692.

Curtis, D.R., and Eeles, J.C. and Lundburg, A. 1958. Intracellular recording from cells in Clarke's column. Acta Physiol Scand. 43 : 303-314.

Eccles, J.C. Hubbard, J.I. and Oscarsson, O. 1961. Intracellular recording from cells of the ventral spinocerebellar tract. J Physiol. 158 : 486-516.

Eccles, R.M. and Lundberg, A. 1958. Integrative pattern of Ia synaptic on motoneurons of hip and knee muscles. J Physiol. 144 : 271-298.

Edgley, S.A., and Jankowska, E. 1988. Information processed by dorsal horn spinocerebellar tract neurons in the cat. J Physiol. 397 : 81-97.

Fetz, F.E., Jankowska, E., Johannisson, T. and Lipski, J. 1979.
Autogenetic inhibition of motoneurons by impulses in group
Ia muscle spindle afferent. J Physiol. 293 : 173-195.

Gurd, J.W. 1977. Synaptic plasma membrane glycoprotein : molecular
identification of lectin receptors. Biochemistry. 16 :
369-374.

Gonatas, N.K., Harper, C., Mizutani, T. and Gonatas, J.O. 1979.
Superior sensitivity of conjugates of Horseradish Peroxidase
with Wheat Germ Agglutinin for studies of retrograde axonal
transport. J Histochem Cytochem. 27 : 728-734.

Gerfen, C.R., O'Leary, D.D.M. and Cowan, W.M. 1982. A note on the
transneuronal transport of Wheat Germ Agglutinin conjugated
Horseradish Peroxidase in the avian and rodent visual system.
Exp Brain Res. 48 : 443-448.

Grant, G., Wiksten, B., Berkley, K.J. and Aldskogius, H. 1982. The
location of cerebellar projecting neurons within the lumbosacral
spinal cord in the cat. An anatomical study with HRP and
retrograde chromatolysis. J Comp Neurol. 204 : 336-348.

Grant, G. and Xu, Q. 1988. Routes of entry into the cerebellum from
the lower part of the spinal cord. An experiment anatomical
study in cat. Exp Brain Res. 72 : 543-561.

Harrison, P.J., Hultborn, H., Jankowska, E., Katz, R., Storai, B. and Zytnicki, D. 1984. Labelling of interneurons by retrograde transsynaptic transport of Horseradish Peroxidase from motoneurons in rats and cats. Neurosci Lett. 45 : 15-19.

Harrison, P.J., Jankowska, E. and Johannisson, T. 1983. Shared reflexed pathways of group I afferent of different cat hindlimb muscles. J Physiol. 338 : 113-127.

Harrison, P.J., Jankowska, E. and Zytnicki, D. 1984. Labelling of interneurons mediating reciprocal inhibition of feline motoneurons by retrograde transsynaptic transport of WGA-HRP. Acta Physiol Scand. 120 : 14A.

_____. 1986. Lamina VIII interneurons interposed in crossed reflex pathways in the cat. J Physiol. 371 : 147-166.

Hongo, T., Jankowska, E., Ohno, E., Sasaki, T., Yamashita, M. and Yoshida, K. 1983a. Inhibition of dorsal spinocerebellar tract cells by interneurons in upper and lower lumbar segments in the cat. J. Physiol. 342 : 145-159.

_____. 1983b. The same interneurons inhibit dorsal spinocerebellar tract cells and lumbar motoneurons in the cat. J. Physiol. 342 : 161-180.

- Hultborn, H., Illert, M. and Santini, M. 1976a. Convergence on interneurons mediating the reciprocal Ia inhibit of motoneurons. I. Disynaptics Ia inhibition of Ia inhibitory interneurons. Acta Physiol Scand. 96 : 351-367.
- _____. 1976b. Convergence on interneurons mediating the reciprocal Ia inhibit of motoneurons. II. Effects from segmental flexor reflex pathways. Acta Physiol Scand. 96 : 359-367.
- _____. 1976c. Convergence on interneurons mediating the reciprocal Ia inhibit of motoneurons. III. Effects from supraspinal pathways. Acta Physiol Scand. 96 : 359-367.
- Itaya, S.K. and Van Hoesen. 1982. WGA-HRP as a transneuronal marker in the visual pathways of monkey and rat. Brain Res. 236 : 199-204.
- Jankowska, E. 1983. On the multiplicity of neuronal pathways from muscle spindles and tendon organs and their selection in the cat. J Physiol. 78 : 772-774.
- _____. 1985. Further indications for enhancement of retrograde transneuronal transport of WGA-HRP by synaptic activity. Brain Res. 341 : 403-408.
- Jankowska, E., Johannsson, T. and Lipski, J. 1981a. Common interneurons in reflex pathways from group Ia and Ib afferents of ankle extensors in the cat. J Physiol. 310 : 381-402.

Jankowska, E. and Lindstrom, S. 1970. Morphological identification of physiological defined neurons in the cat spinal cord.
Brain Res. 20 : 323-326.

_____. 1972. Morphology of interneurons mediating Ia reciprocal inhibition of motoneurons in spinal cord of the cat. J Physiol. 226 : 805-823.

Jankowska, E. and McCrea, D.A. 1983. Shared reflex pathways from Ib tendon organ afferents and Ia muscle spindle afferents in the cat. J Physiol. 338 : 99-111.

Jankowska, E. and McCrea, D.A. and Mackel, R. 1981b. Pattern of non-reciprocal inhibition of motoneurons by impulses in group Ia muscle spindle afferents in the cat. J Physiol. 316 : 393-409.

Jankowska, E. and Skoog, B. 1986. Labelling of midlumbar neurons projecting to cat hindlimb motoneurons by transneuronal transport of a Horseradish Peroxidase conjugate. Neurosci Lett. 71 : 163-168.

Kristensson, K. and Olsson, Y. 1971. Retrograde axonal transport of protein. Brain Res. 29 : 363-365.

Lafleur, J., Delean, J. and Poirier, L.J. 1974. Physiopathology of the cerebellum in the monkey. I. Origin of cerebellar afferent nervous fibers from the spinal cord and brain stem. J Neurol Sci. 22 : 471-490.

La Vail, J.H. and La Vail, M.M. 1972. Retrograde axonal transport in the central nervous system. Science. 176 : 1416-1417.

Lunberg, A. 1979. Multisensory control of spinal reflex pathways. In R. Granit and O. Pompeiano (eds.), Reflex Control of Posture and Movement, 11-28. Amsterdam : Elsevier.

Matsushita, M. and Hosoya, Y. 1979. Cells of origin of the spinocerebellar tract in the rat. Studied with the method of retrograde transport of Horseradish Peroxidase. Brain Res. 173 : 185-200.

Matsushita, M., Hosoya, Y. and Ikeda. M. 1979. Anatomical organization of the spinocerebellar system in the cat as studied by retrograde transport of Horseradish Peroxidase. J Comp Neurol. 184 : 81-106.

Matsushita, M. and Ikeda, M. 1980. Spinocerebellar projections to the vermis of the posterior lobe and the paramedian lobule in the cat as studied by retrograde transport of Horseradish Peroxidase. J comp Neurol. 192 : 143-162.

Mesulam, M.M. 1978. Tetramethylbenzidine for Horseradish Peroxidase neurohistochemistry : a noncarcinogenic blue reaction produce with superior sensitivity for visualing neuronal afferents and efferents. J Histochem Cytochem. 26 : 106-117.

Oscarsson, O. 1973. Functional organization of spinocerebellar paths. In A. Iggo (ed.), Hand book of Sensory Physiology, vol II. Somatosensory System, 338-380. Springer : Berlin Heidelberg New York.

Petras, J.M. 1977. Spinocerebellar neurons in the rhesus monkey. Brain Res. 130 : 146-151.

Petras, J.M. and Cumming, J.F. 1977. The origin of spinocerebellar pathways. II. The nucleus centrobasalis of the cervical enlargement and the nucleus dorsalis of the thoracolumbar spinal cord. J Comp Neurol. 173 : 693-716.

Rexed, B. 1954. A cytoarchitectonic atlas of the spinal cord in the cat. J Comp Neurol. 100 : 297-379.

Ryall, R.W. 1981. Patterns of recurrent excitation and mutual inhibition of cat Renshaw cells. J Physiol. 316 : 439-452.

Schwab, M.E., Javory-Agid, F. and Agid, Y. 1978. Labelled Wheat Germ Agglutinin (WGA) as a new, highly sensitive retrograde tracer in the rat brain hippocampal system. Brain Res. 152 : 145-150.

Snyder, R.L., Fall, R.L.M. and Mehler, W.R. 1978. A comparative study of the neurons of origin of the spinocerebellar afferents in the rat, cat and squirrel monkey based on the retrograde transport of Horseradish Peroxidase. J Comp Neurol. 181 : 833-852.

Tapper, D.N., Mann, M.D., Brown, P.B. and Cogdell, B. 1975. Cells of origin of the cutaneous division of the dorsal spinocerebellar tract. Brain Res. 85 : 59-63.

Wiksten, B. 1979. The central cervical nucleus in the cat. II. The cerebellar connections studied with retrograde transport of Horseradish Peroxidase. Exp Brain Res. 36 : 155-173.

Xu, Q. and Grant, G. 1988a. Course of spinocerebellar axons in the ventral and lateral funiculi of the spinal cord with projections to the anterior lobe. An experimental anatomical study in the cat using retrograde tracing technique. J Comp Neurol. in press.

_____. 1988b. Collateral projections of neurons from the lower part of the spinal cord to anterior and posterior cerebellar termination areas. A retrograde fluorescent double labelling study in the cat. Exp Brain Res. 72 : 562-576.

_____. 1988c. Do certain spinocerebellar neurons in lamina IX at
lumbrosacral levels send collaterals to peripheral nerves ?
A retrograde fluorescent double labelling study in cat.
Arch. Ital. Biol. 126 : 179-192.

ภาคผนวก

ກາຄະນາດ ດ.

HRP - perfusion (Rat)

Rinse - solution (34°C)

1000 ml NSS

0.5 ml (2500U) HEPARIN

Fix - solution (room temp.) 1000 ml

1% PARAFORMALDEHYDE in 320 m Osm phosphate buffer (PH 7.4)

1.5% GLUTARALDEHYDE

4% SUCROSE

Postfix - solution (4°C) 500 ml

500 ml Phosphate buffer PH 7.4

50 gm Sucrose

Storing solution

Same as postfix (or add 4% Glutaraldehyde if fixation is bad.)

ກາລົມວົກ ພ.

Peroxidase TMB

For 125 ml of solution

1. Rinse in distilled water 3x1 minute

2. Pre incubate B to A for 20 minutes

(A = distilled water 116.6 ml + sodium nitroprusside
0.125 gm + acetate buffer pH 3.3 6.25 ml)

(B = 6.25 mg TMB dissolved in 1.8 ml absolute alcohol
on 40 degree centigrade on heating plate.)

3. Enzymatic reaction for 20 minutes

Add 2.5 ml of 0.3% H₂O₂

4. Rinse in TMB rinsing buffer 3x1 minute

5. Mount on gelatin slides

6. Dry in room temperature

7. Counterstain with 1% Neutral Red for 1-1.5 minutes.

(Butanol staining procedure)

Preparation of TMB rinsing Buffer

5 ml of acetate buffer pH 3.3 + 95 ml of distilled water

Acetate buffer pH 3.3

27 gm CH₃COONa.3H₂O + 190 ml 1 M HCl + 810 ml distilled
water

ກາຈົກນາດ ຮ.

Staining with Neutral Red

1. 1% Neutral Red 1.15 min
2. Acetate buffer 2 times
3. Butanol 3x1 min
4. Butanol + Xylene 1:1 3 min
5. Xylene 1 2 min
- Xylene 2 1 min
- Xylene 3
6. Mount with permount

Acetate buffer

Distilled water 2 litres + Acetate buffer pH 4.8 80 ml

Acetate buffer pH 4.8

500 ml of 0.1 N acetic acid + 750 ml of 0.1 N sodium acetate
filter before using

1% buffer neutral red solution

Add 40 ml of pH 4.8 acetate buffer to each litre of a 1% aqueous neutral red solution. Filter before using.

ประวัติสูเขียน

นางสาวจันทรารณ แสงแข เกิดเมื่อวันที่ 16 ตุลาคม พ.ศ.2506 สำเร็จการ
ศึกษาวิทยาศาสตรบัณฑิต (พยาบาล) จากมหาวิทยาลัยมหิดล วิทยาเขตธิรราช เมื่อปี พ.ศ.
2529