



REFERENCES

- Abraham, K., Schuett, K., Muller, I., and Hoffmeister, H. 1970. Kontinuierliche Polyacrylamid-Elektrophorese I. Untersuchungen an Normalseren. Z. Klin. Chem. u. Klin. Biochem. 8: 92.
- Allen, R.C., and Moore, D.J. 1966. A vertical flat-bed discontinuous electrophoresis system in polyacrylamide gel. Anal. Biochem. 16: 457.
- Alper, C.A., Peters, J.H., Britch, A.G., and Gardner, F.H. 1965. Haptoglobin synthesis. Invivo studies of the production of haptoglobin, fibrinogen and α -globulin by the canine liver. J. Clin. Invest. 44: 574.
- Aly, F.W., and Braun, H.J. 1974. Die quantitative Bestimmung einiger spezieller plasmaproteine und ihre diagnostische Bedeutung, in serumproteine. A Engelhardt. u. H. Lommel ed. Verlag Chensie, Weinheim, Bergstr. W. Germany.
- Anastassea Vlachou, C., Kattamis, C., Lagos, P., Konstantoulakis, M., and Matsaniotis, N. 1972. Haptoglobin and α_2 -macroglobulin levels in children with rheumatoid arthritis. Clin. Chim. Acta. 44: 259.
- Ancell, H. 1839 - 1940. Course of lectures on the physiology and pathology of the blood and the other animal fluids. Lancent. 1: 222.

Aronson, K.F., Ekelund, G., Klindmark, C.O., and Laurell, C.B. 1972.

Sequential changes of plasma proteins after surgical trauma.

Scand. J. Clin. Lab. Invest. 29 (suppl.124): 127 - 236.

Barret, A.J., and Starky, P.M. 1973. Inhibition by α_2 -macroglobulin and other serum protein. Biochem. J. 131: 823.

Bellabarba, D., Inada, M., Versano Aharon, N., and Sterling, K. 1968.

Thyroxine transport and turnover in major non-thyroidal illness. J. Clin. Endocr. 23: 1023.

Berson, S.A., and Yallow, R.S. 1954. The distribution of ^{131}I -labelled human serum albumin introduced into ascitic fluid. Analysis of the kinetics of a three compartment catenary transfer system in man and speculations on possible sites of degradation. J. Clin. Invest. 33 : 377.

Bowman, B.H. 1968. Serum Groups. Ser Haemat. 1: 97.

Braun, H.J., and Aly, F.W. 1971. Die Klinische Bedeutung der quantitative Serum-Hemopexinbestimmung im Verleich zum Haptoglobin. Klin. Wochr. 49: 451.

Brown, R.K., Baker, W.H., Peterskofsky, A., and Kauffman, D.L. 1954. Crystallization and properties of glycoprotein isolated from human plasma. J. Am. Chem. Soc. 76: 4244.

Cherftel, R.I., Cloarec, L., Moretti, J., Rafelson, M., and Jayle, M.F. 1960. Etude de l'haptoglobine III: Composition clinique de la partic glucidique. Bull. Soc. Chim. Biol. 42 : 993.

Clarke, H.G.M., Freeman, T., and Pryse-Phillips, W. 1971. Serum protein changes after injury. Clin. Sci. 40: 337.

Connell, G.E., and Shaw, R.W. 1961. The purification of haptoglobin. Canad. J. Biochem. 39: 1013.

Cooke, G.C. Mulligan, R., and Scherlock, S. 1971. Controlled prospective trial of corticosteroid therapy in active chronic hepatitis. Quart. J. Med. 40: 159.

Cox, D.W. 1966. Factors influencing serum ceruloplasmin levels in normal individuals. J. Lab. Clin. Med. 68: 893.

Dobryszycha, W., Zeineh, R., Ebron, E., and Kukral, J.C. 1969. Metabolism of Plasma Proteins in Injury States. Clin. Sci. 36: 231.

Dykes, P.W. 1968. The rates of distribution and in patients with cirrhosis of the liver. Clin. Sci. 34: 161.

Felgenhauser, K. 1970. Quantitative and specific determination methods after disc-electrophoresis of serum protein. Clin. Chim Acta. 27: 305.

Ganrot, P.O., and Schersten, B. 1967. Serum γ -macroglobulin concentration and its variations with age and sex. Clin. Chim. Acta. 15: 113.

Gardon, S., and Bearn, A.G. 1966. Hemoglobin binding capacity of isolated haptoglobin polypeptide chains. Proc. Soc. Exp. Biol. 121: 846.

Giblett, E.R. 1969. Genetic Markers in Human Blood. ed. Blackwell, Oxford.

Giblett, E.R. 1974. Haptoglobin: In Structure and Function of Plasma Protein. ed. Allison, A.C. pp. 55-72. Plenum Press. London and New York.

Gitlin, D., and Biasucci, A. 1969. Development of γ G, γ A, γ M, β 1G/ β 1A, C'1 Esterase Inhibitor, Ceruloplasmin, transferrin, Hemopexin, Haptoglobin, Fibrinogen, Plasminogen, α_1 -Antitrypsin, Orosomucoid, β -Lipoprotein, α_2 -Macroglobulin and Prealbumin in the human conceptus. J. Clin. Invest. 48: 1433.

Haellen, J., Laurell, C.B. 1972. Plasma protein pattern in cirrhosis of the liver. Scand. J. Clin. Lab. Invest. 29 (suppl. 124): 97-103.

Hasch, E., and Tygstrup, M. 1967. Albumin synthesis rate as a measure of liver function in patients with cirrhosis. Acta. Med. Scand. 182: 1.

Heimburger, N., Schultze, H.E., Heidl, K., Haupt, H., Storiko, K., and Schwich, H.G. 1963. Proceeding of the ninth Congress European Society Haematology. Lisbon. (Karger, Basel, 1963). pp.1315.

- Hochwald, G.M., Thorbecke, G.J. and Asofsky, R. 1961. Sites formation of immune globulin and a component of C'3. I. A new technique for the demonstration of the synthesis of individual serum proteins by tissues in vitro. J. Exp. Med. 114: 459.
- Hoffmeister, H., and Schuett, K.H. 1972. Diagnostische Verwendbarkeit der Polyacrylamid-Electrophoresis. Dtsch. Med. Wochensch. 97: 1464.
- Hoffmeister, H., 1974. Potential Applications of Polyacrylamide Gel Electrophoresis of Clinical Diagnosis: In Electrophoresis and Isoelectric Focussing in Polyacrylamide Gel. eds. Allen, A.C., and Maurer, H.R. Berlin and New York Walter der Gruyter.
- Houchin, O.B. 1958. A rapid colorimetric method for the quantitative determination of copper oxidase activity (ceruloplasmin). Clin. Chem. 4: 519.
- Hrkal, Z., and Muller, E.U. 1971. Partial characterisation of the heme-binding serum glycoproteins rabbit and human hemopexin. Biochemistry. 10: 1746.
- Inada, M., and Sterling, K. 1967. Thyroxine turnover and transport in Laennec's cirrhosis of the liver. J. Clin. Invest. 46: 1275.
- James, K., Taylor, F.B., and Feidenberg, H.H. 1966. Trypsin stabilizer in human serum. The role of α_2 -macroglobulin. Clin. Chim. Acta. 13: 357.

- Javid, J. 1967. Human Serum Haptoglobin. Sem. Haemat. 4: 35.
- Jayle, M.F., and Moretti, J. 1962. Haptoglobin: biochemical genetic and Physiological aspects. eds. Moore, C.V. and Brown, E.B. Progress in Haematology Vol. 3 p.342 (Grune and Stratton, New York).
- Jeanloz, R.W. 1966. Immuno-electrophoretic analysis in Method of Biochemical Analysis. ed. Glick, D. Vol. 7 pp. 1 Interscience Publishers, New York.
- Kashiwagi, N., Groth, C.G., and Starzl, T.E. 1968. Changes in serum haptoglobin and group specific component after orthotopic liver homotransplantation in human. Proc. Soc. Exp. Biol. Med. 128: 247.
- Koj, A. 1974. Acute-phase reactants. In Structure and Function of Plasma Protein. ed. Allison, A.C. Vol. 1 pp. 73 Plenum Press. London and New York.
- Krauss, S., and Sarcione, E.J. 1964. Synthesis of serum haptoglobin by the isolated perfused rat liver. Biochem. Biophys. Acta. 90: 301.
- Krauss, S., Schrott, M., and Sarcione, E.J. 1966. Haptoglobin Metabolism in Hodgkin's Disease. Am. J. Med. Sci. 252: 184.
- Kukral, J.C., Kerth, J.C., Pancer, R.J., Gromer, D.W., and Henegar, G.C. 1961. Plasma protein synthesis in the normal dog and after total hepatectomy. Surg. Gynec. Obstet. 113: 360.

- Kushner, I., Edgington, T.S., Trimber, C., Liem, H.H., and Muller, E.K. 1972. Plasma hemopexin homeostasis during the acute phase response. J. Lab. Clin. Med. 80: 18.
- Lane, R.S. 1966. Changes in plasma transferrin levels following the administration of iron. Brit. J. Haemat. 12: 249.
- Lane, R.S. 1968. Transferrin synthesis in the rat: A study using the fluorescent antibody technique. Brit. J. Haemat. 15: 355.
- Laurell, C.B. 1972. Electroimmuno Assay. Scand. J. Clin. Lab. Invest. 29 (suppl. 124): 21.
- Laurell, C.B., Kullander, B., and Thorell, J. 1970. Rate of Plasma Protein Normalization after Parturition and Withdrawal of Oral Contraceptives. Scand. J. Clin. Lab. Invest. 26: 345.
- Lombart, C.B., Moretti, J., and Jayle, M.F. 1965. Preparation et proprietes physiques des haptoglobin de lapin et de rat. Biochem. Biophys. Acta. 97: 262.
- Mandel, E.E. 1959. Serum iron and iron-binding capacity in clinical diagnosis. Clin. Chem. 5: 1.
- Manuel, Y., De Fontaine, M.C., Bourgoin, J.J., Pargent, M., and Soueck, J.M. 1971. Serum hemopexin levels in patients with malignant melanomas. Clin. Chim. Acta. 31: 485.

- McFalane, H., Okubadejo, M., and Reddy, S. 1972. Transferrin and Staphylococcus aureus in Kwashiorkor. Am. J. Clin. Path. 57: 587.
- Meier, H., and Macpike, A.D. 1968. Levels and heritability of serum ceruloplasmin activity in Inbred. Strains of Mice. Proc. Soc. Exp. Biol. Med. 128: 1185.
- Miller, L.L., Bly, C.G., Watson, M.L., and Bale, W.F. 1951. The dominant role of the liver in plasma protein synthesis. J. Exp. Med. 94: 431.
- Miller, L.L., and Bale, W.F. 1954. Synthesis of all plasma protein fractions except gamma globulin by the liver. J. Exp. Med. 99: 125.
- Morgan, W.T., and Muller, E.U. 1972. Introductions of porphyrins with rabbit hemopexin. J. Biol. Chem. 247: 7181.
- Mouray, H., Moretti, J., and Jayle, M.F. 1964. Biosynthese de l'haptoglobine per perfusion du foie isole de lapine. C.R. Acad. Sci. Paris. 259: 2721.
- Mouray, H. 1966. Biosynthese de l'haptoglobine chez le lapin. eds. Fonlon, R. Paris.
- Neale, F.C., Aber, G.M., and Northam, B.E. 1958. The demonstration of intravascular haemolysis by means of serum paper electrophoresis and a modification of Schumm's reaction. J. Clin. Path. 11: 206.

- Nilprabhassorn, P., Vootiprux, V., and Suvonge, C. 1976. An analysis of the cancer statistic of The Cancer Institute Siriraj Hospital 1969 - 1972. J. Med. Ass. Thai. 59: 162.
- O'Hara, H., Watanabe, K., and Wada, T. 1968. Turnover rate of serum haptoglobin in malignant neoplastic diseases. Clin. Chim. Acta. 19: 41.
- Ohlsson, K. 1971. Elimination of ^{125}I -trypsin α_2 -macroglobulin complex from blood by reticuloendothelial cells in dog. Acta. Physiol. Scand. 81: 269.
- Ohlsson, K. 1974. International between endogeneous proteases and plasma inhibitors. In proteinase inhibitors. Proceedings of the 2th. International Research Conference. eds. Fritz, H., Tschesche, H., Greene, L.J. and Truscheit, E. pp. 96-105. Springer Verlag Berlin. Heidelberg, New York.
- Oppenheimer, J.H., Surks, M.I., Bernstein, G., and Smith J.C. 1965. Metabolism of ^{131}I -labelled Thyroxine-binding Prealbumin in man. Science. 149: 748.
- Oppenheimer, J.H., 1968. Role of plasma proteins in the binding, distribution, and metabolism of the thyroid hormones. New Engl. J. Med. 278: 1153.
- Owen, J.A., Smith, R., Padanyl, R., and Martin, J. 1964. Serum haptoglobin in disease. J. Martin. Clin. Sci. 26: 1.

- Peters, J.H., and Alper, C.A. 1965. Haptoglobin Synthesis II. Cellular Localization Studies. J. Clin. Invest. 45: 341.
- Peters, T. Jr. 1970. Advances in Clinical Chemistry. eds. Sobotka, H., Stewart, C.R. Academic Press, New York, Vol. 13 pp. 37.
- Pinida, E.P., Ravin, H.H., and Rutenburg, A.M. 1962. Serum ceruloplasmin. Observation in patients with cancer, obstructive jaundice and other diseases. Gastroenterology 43: 266.
- Pongpaew, P., Migasena, P., and Schelp, F.P. 1975. Electrophoresis of human serum with polyacrylamide gradient gel slabs. J. Med. Ass. Thai. 58: 15.
- Ravin, H.A. 1961. An improved colorimetric enzymatic assay of ceruloplasmin. J. Lab. Clin. Med. 58: 161.
- Reutsch, I., Quoted by Aly, F.W., and Braun, H.J. 1974. Die Quantitative Bestimmung Einiger Spezieller Plasmaproteine Serumproteine, Verlag Chemic. Weinheim/Bergstr. W. Germany. pp. 9.
- Rice, E.W. 1960. Correlation between serum copper, ceruloplasmin activity and C-reactive protein. Clin. Chim. Acta. 5: 632.
- Rice, E.W. 1961. A study of correlation between C-reactive protein and certain other acute-phase reactants. Clin. Chim. Acta. 6: 170.
- Robert, R.C., Reisen, W.A., and Hall, P.K. 1974. Studies in the quaternary structure of human serum α_2 -macroglobulin. In

- Proteinase Inhibitors. Proceeding of the 2th International Research Conference, 1974. eds. Fritz, H., Tshesche, H., Greene, L.J., and Truscheit, E. pp. 63 Springer Verlag. Berlin, Heidelberg, New York.
- Rosenoer, V.M. 1976. Plasma protein turnover. eds. Bianchi, R., Mariani, G., and McFalane, A.S. Macmillans, London, pp. 103.
- Rothschild, M.A., Oratz, M., and Schreiber, S.S. 1969. Serum Albumin, Amer. J. Digest. Dis. 14: 711.
- Rothschild, M.A., Oratz, M., and Schreiber, S.S. 1970. Changing Concept of Alumin Metabolism and Distribution in cirrhosis of the liver. Scand. J. Gastroenterol. (suppl. 17) 5: 17.
- Rossi, T., Hirvonen, T., and Toivanen, P. 1970. The concentration of prealbumin in human fetal and infant sera. Scand. J. Clin. Lab. Invest. 26: 35.
- Sass-Kortsak, A. 1965. Copper metabolism. Adv. Clin. Chim. 8: 1.
- Schaffner, F., Singer, E.J., Anthony, D.D., and Popper, F.J. 1962. Serum electrophoresis on polyacrylamide gel in liver disease. J. Mount. Sinai. Hosp. 29: 420.
- Schmid, K., and Tokita, K. 1967. The α_1 -acid glycoprotein variants of certain Caucasian patients. Clin. Chim. Acta. 17: 39.
- Schmid, K., Feild, R.A., and Yoshizaki, 1968. Constancy of α_1 -acid glycoprotein variants of Caucasian Patients under conditions of severe stress. J. Med. Genet. 5: 36.

Schortridge, K.F., Robinson, J.D., and Ng. M.H. 1973. Serum α_2 -macroglobulin levels in normal Chinese. Clin. Chim. Acta. 48: 221.

Schuett, K.H., and Hoffmeister, H. 1971. Kontinuierliche Polyacrylamid-electrophorese II. Möglichkeiten zur Diagnose von malignen und entzündlichen Erkrankungen der Lunge. Z. Klin. Chem. u. klin. Biochem. 9: 206.

Schultze, H.E., Heide, K., and Haupt, H. 1961. Über die Einwirkung von konzentrierter Salzsäure auf Lavoglucosan. Naturwissenschaften. 48: 696.

Schultze, H.E., Biel, H., Haupt, H., and Heide, K. 1962. Über die Gc-Komponenten von Hirschfeld II. Darstellung und Eigenschaften. Naturwissenschaften. 49: 108.

Schultze, H.E., and Heremans, J.F. 1966. Molecular Biology of Human Proteins. eds. Schultze, H.E., and Heremans, J.F. Elseveir Publ. Co. Amsterdam. London., New York.

Schultze, H.E., and Heremans, J.F. 1970. Molecular Biology of Human Proteins. eds. Schultze, H.E., and Heremans, J.F. Elseveir Publ. Co. Amsterdam, London, New York. pp.34, 48.

Seal, U. 1964. Vertebrate Distribution of Serum Ceruloplasmin and Sialic acid and the effects of pregnancy. Comp. Biochem. Physiol. 13: 143.

- Seery, V.L., Hathway, G., and Muller, E.U. 1972. Hemopexin of human and rabbit: Molecular Weight and Extinction Coefficient. Arch. Biochem. Biophys. 150: 269.
- Seery, V.L., Muller, E.U. 1973. Binding of porphyrins to rabbit hemopexin and albumin. J. Biol. Chem. 248: 3796.
- Shim, B.S., and Bearn, A.G. 1964. Immunological and biological studies on serum haptoglobin. J. Exp. Med. 120: 611.
- Smithies, O. 1959. An Improved Procedure for Starch gel Electrophoresis: Further variations in the serum Proteins of Normal Individual. Biochem. J. 71: 585.
- Smithies, O., Connell, G.E., and Dixon, G.H. 1962. Chromosomal rearrangements and the evolution of haptoglobin genes. Nature. 196: 232.
- Socolow, E.L., Woeber, K.A., Purdy, R.H., Holloway, M.T., and Ingbar, S.H. 1965. Preparation of ¹³¹I-labelled human serum prealbumin and its metabolism in normal and sick patients. J. Clin. Invest. 44: 1600.
- Stabilini, R., Vergani, C., Agostini, A., Pugno Vanoni Agostini, R. 1968. Influence of age and sex on prealbumin levels. Clin. Chim. Acta. 20: 358.
- Sterling, K. 1951. The turnover rate of serum albumin in man as measured by ¹³¹I-tagged albumin. J. Clin. Invest., 30: 1228.

Sunderman, F.W., and Namoto, S. 1970. Measurement of human serum ceruloplasmin by its p-phenylenediamine oxidase activity.

Clin. Chem. 16: 903.

Surks, M.I., and Oppenheimer, J.H. 1964. Postoperative changes in the concentration of thyroxine-binding prealbumin and serum free thyroxine. J. Clin. Endocr. 24: 794.

Synder, S., and Ashwell, G. 1971. Quantitation of specific serum glycoproteins in malignancy. Clin. Chim. Acta. 34: 449.

Talamo, R.C., Langley, C.E., and Twarog, F. 1975. Protease inhibitors in cord and maternal serum at birth. Behring. Institute Mitteilungen. 58: 46.

Tavill, A.S., Craigie, A., and Rosenoer, V.M. 1968. The measurement of the synthetic rate of albumin in man. Clin. Sci. 34: 1.

Van den Scherieck, H.G. 1969. Les proteins protenses de thyroxine. ed. Louvein.

Weinfeld, A. 1965. Proceeding of International Congress Society of Haematology Stockholm of the 10th. Series Haematologica, Munksgaard, Copenhagen.

Weisman, S., Goldsmith, B., Winzler, R.J., and Lepper, M.H. 1961. Turnover of plasma orosomucoid in man. J. Lab. Clin. Med. 57: 7.

Werner, M. 1969. Serum protein changes during the acute-phase reaction.

Clin. Chim. Acta. 25: 299.

Wewalka, F. 1969. Untersuchung der plasmaproteine und der protein-abbauprodukte. In E. Deutsch and G. Geyer ed. Laboratoriumsdianostik. Normalwerte und Interpretation. Verlag August Steinkopf, Berlin, W. Germany.

Wilkinson, P.W., and Mandenhall, C.L. 1963. Serum albumin turnover in normal subjects and patients with cirrhosis measured by ¹³¹I-labelled human albumin. Clin. Sci. 25: 281.

Wintrobe, M.M., Markowitz, H., Gubler, C.J., Mahony, J.P., and Cartwright, G.E. 1955. Studies of copper metabolism XIV. Copper ceruloplasmin and oxidase activity in sera of normal human subjects, pregnant women, and patients with infection, hepatolenticular degeneration and the nephrotic syndrome. J. Clin. Invest. 34: 1498.

Zeineh, R.A., and Kukral, J.C. 1970. The turnover rate of orosomucoid in burned patients. J. Trauma. 10: 493.

VITA

Name: Sub-Lt. Pimathorn Juttijudata

Education: Bachelor of Science in Pharmacy in 1971,
Mahidol University, Bangkok, Thailand.

Site of the Employer's Office:

Territorial Defense Department
Ministry of Defense.

