



REFERENCES

ภาษาไทย

บำรุง ตับติเสว, ตราดล กราไดย และกัลยา กราไดย. มาศชเวท เล่ม 1. คณะเภสัชศาสตร์ จุฬาลงกรณ์มหาวิทยาลัย, 2526.

มหาวิทยาลัยเชียงใหม่. หน่วยงานตีกษาดัมกิริบลาน. ตำราสมุนไพรล้านนา. เชียงใหม่ : โครงการร่วมระหว่างพิพิธภัณฑ์ชาติพันธุ์ไทยแห่งชาติโอชาガและสถาบันวิจัยสังคม หน่วยงานตีกษาวิจัย ตัมกิริบลาน มหาวิทยาลัยเชียงใหม่, 2525.

เดิม สมิตินันทน์. ชื่อพรรณไม้แห่งประเทศไทย (ชื่อพุกศาสตร์, ชื่อพื้บเมือง). กรุงเทพมหานคร : กรมป่าไม้, 2523.

ภาษาอังกฤษ

Ahmad, A., Khan, K.A., Ahmad, V.U., and Qazi, S..Antibacterial activity of juliflorine isolated from *Prosopis juliflora* Planta Med. 52 (1986) : 285-288.

Bailey, W.R., and Scott, E.G..Diagnostic Microbiology. 4th ed. Saint Louis : C.V. Mosby Company, 1974.

Barry, L.A..The antimicrobial susceptibility test : Principles and practices. Philadelphia : Lea & Febiger, 1976.

Barton, D.H.R. and Bruun, T.. Some observations on the constitution of usnic acid. J.Chem.Soc. (1953) : 603.

- Bejairi, A.M., Qureshi, S., Chaudhry, M.A., Krishna, Dr., Shah, A.H.. Mitodepressive, clastogenic and biochemical effects of (+)-usnic acid in mice. J.Ethnopharmacol. 33 (1991) : 217-220.
- Benezra, C., Stampf, J.L., Barbier, P., Ducoms, G.. Enantiospecificity in allergic contact dermatitis. A review and new results in *Frullania*-sensitive patients. Contact Dermatitis, 13 (1985) : 110-114.
- Birkbeck, A.A., Sargent, M.V. and Elix, J.A.. The structure of the lichen depsidones fulgidin and isofulgidin. Aust. J. Chem. 43 (1990) : 419-425.
- Bohlmann, F., Gupta, R.K., Jakupovic, J., King, R.M. and Robinson, H.. Naturally occurring terpene derivatives. part 437. four oplopanone and bisabolone and bisabolone derivatives from *Regulia nudicaulis*. Phytochemistry 21 (1982) : 1665-1668.
- Borkowski, B., Wozaniak, W., Gertig, H., and Werakso, B. Bacteriostatic action of some compounds from lichen *Cetraria islandica* and of usnic acid. Dissertations Pharm. 16 (1964) : 189-194.
- Budavar, S., O' Neil, M.J., Smith, A., and Heckelman, P.E. The Merck Index : and encyclopedia of chemicals, drugs, and biologicals. In Budavar, S.ed. 11 eds. The United States : Merck and Co., Inc., 1989.
- Burkill, L.H.. A Dictionary of the Economic Products of the Malay Peninsula vol II University Press Oxford, 1935
- Capriotti, A.. The effect of benzylidimethyl-2-2{2-[p-(1,1,3,3-tetramethylbutyl) phenoxy] ethoxyl} ethylammonium usneate (USNO) on yeasts. Giorn. Microbiol. 7 (1959) : 187-206.
- Caccamese, S., Compagnini, A., Toscano, R.M. and Cascio,O.. Methyl beta- orcinocarboxylate and atranol from the lichen *Stereocaulon vesuvianum*. J. Nat. Prod. 49 (1986) : 1159-1160.

- Caccamese, S., Toscano, R.M., Bellesia, F. and Pinetti, A.. Methyl beta-orcinoiccarboxylate and depsides from *Parmelia furfuracea*. J. Nat. Prod. 481 (1985) : 157-158.
- Conover, M.A. et al..Usnic acid amide, a phytotoxin and antifungal agent from *Cercosporidium henningsii*. Phytochemistry. 31 (1992) : 2999-3001.
- Cooper, A.B. et al..Synthesis of (+)-8-methyl cercosporamide stereochemical correlation of natural (-)-cercosporamide with (+)-usnic acid. Tetrahedron. 48 (1992) : 4757-4766.
- Creaser, E.H. The induced (adaptive) biosynthesis of β -galactosidase in *Staphylococcus aureus*. J. Gen. Microbiol. 12 (1955) : 288.
- Culberson, C.F..Chemical and Botanical Guide to Lichen Products The University of North Carolina Press, 1969.
- _____.The structure of a new depsidone from the lichen *Parmelia livida*. Phytochemistry. 5 (1966) : 815-818.
- _____. The lichen substances of the genus *Evernia*.Phytochemistry. 24 (1963) : 335-340.
- _____. Some constituents of the lichen *Ramalina siliquosa* Phytochemistry. 4 (1965) : 951-961.
- _____.and Culberson, W.L.. Beta-orcinol derivatives in lichens: biogenetic evidence from *Oropogon loxensis*. Exp. Mycol. 2 (1978) : 245-257.
- Curd, H.F. and Robertson, A..Usnic acid part V. J. Chem. Soc.. (1937) : 894-901.
- David, F., Elix, J.A. and Wahid B.. Two new aliphatic acids from the lichen *Parmaotrema prasoredisum*. Aus. J. Chem. 437 (1990) : 1297-1300.
- Denyer, S.P., and Baird, R.M.,ed..Guide to Microbiological Control in Pharmaceuticals. 1st ed. England : Ellis Horwood Limited, 1990.

- Devlin, J.P., Mahandru, M.M., Ollis, W.D. and Smith, C.. Phytochemical examination of the lichen, *Lecanora gangleoides* NYL. J. Chem. Soc. Perkin Trans I. (1986) : 1491-1494.
- Ejiri, H., Sankawa, U. and Shibata, S.. Isolation of methyl 2,4- dihydroxy-3,6-dimethyl benzoate, atranorin, lobaric acid and stictic acid from *Stereocaulon vesuvianum*. Phytochemistry. 15 (1976) : 2025C.
- Elix, J.A.. 2'-O-methylphysodic acid and hydroxyphysodic acid: two new depsidones from the lichen *Hypogymnia billardieri*. Aus. J. Chem. 28 (1975) : 849-858.
- _____ and Bennett, S.A.. 6-O-methylarthothelin and 1,3,6-tri-O-methylarthothelin, two new xanthones from a *Dimelaena* lichen. Aus. J. Chem. 439 (1990) : 1587-1590.
- _____ and Engkaninan, U.. 4,5-di-O-methylhiascic acid, a new tridepside from the lichens *Parmelia pseudofatiscens* and *Parmelia horrescens*. Aust. J. Chem. 29 (1976) : 2701-2705.
- _____ and Engkaninan, U.. The structure of lividic acid, a depsidone from the lichen *Parmelia formosana*. Aus. J. Chem. 29 (1976) : 203-207.
- _____ and Jayanthi, V.K.. The isolation and synthesis of the lichen depside 4-O-demethylmicrophyllinic acid. Aus. J. Chem. 40 (1987) : 1851-1859.
- _____, Bennett, S.A. and Jiang, H.. 2,5-dichloro-6-O-methylnorlichexanthone and 4,5-dichloro-6-O-methylnorlichexanthone, two new xanthones from an australian *Dimelaena* lichen. Aus. J. Chem. 448 (1991) : 1157-1162.
- _____, Gaul, K.L. and James, P.W.. Alpha-acetylhypoconstictic acid, a new depsidone from the lichen *Menegazzia disporea*. Aus. J. Chem. 3811 (1985) : 1735-1737.
- _____, Gaul, K.L., James, P.W., and Purvis O.W. Three new lichen depsidones. Aust. J. Chem. 40 (1987) :417-423.

- Elix, J.A., Jayanthi, VK. and Leznoff, CC.. 2,4-di-O-methylgyrophoric acid and 2,4,5-tri-O-methylhiascic acid. new tridepsides from *Parmelia damaziana*. Aust. J. Chem. 34 (1981) : 1757-1761.
- _____, Whitton, A.A. and Jones, A.J.. Triterpenes from the lichen genus *Physcia*. Aus. J. Chem. 35 (1982) : 641-647.
- Emmerich, R., Giez, I., Lange, O. L. and Proksch, P.. Function of lichen substances : deleterious effect of usnic acid and vulpinic acid towards the generalist herbivore *Spodoptera littoralis*. Planta Med. (1992) : 630-631.
- _____. Toxicity and antifeedant activity of lichen compounds against the polyphagous herbivorous insect *Spodoptera littoralis*. Phytochemistry. 33 (1993) : 1389-1394.
- Ghione, M., Parrello, D., Grasso, L.. Usnic acid revisited, its activity on oral flora. Chemioterapia. 7(1988) : 302-5.
- Goncalo, S., Cabral, F., Goncalo, M.. Contact sensitivity to oak moss. Contact Dermatitis. 19(1988) : 355-7.
- Gonzalez, A.G., Barrera, J.B., Perz, E.M.R. and Padron, C.E.H.. Chemical constituents of the lichen *Ramalina hierrensis*. Planta Med. 582 (1992) : 214-218.
- Guenther, E. Essential oils of the plant family Usneaceae. The essential oils. D.Van Nostrand Company, Princeton, New Jersey, Toronto, New York, London, 1960.
- Gunzinger, J. and Tabacchi, R.. Isolation and identification of furfuric acid a new depsidone from the lichen *Pseudevernia furfuracea* (L) ACH.. Helv. Chim. Acta. 687 (1985) : 1936-1939.
- Hale, E.M.. How to know the lichens. Iowa : WM.C.Brown Company, 1969.
- _____. Lichen handbook. The United states Pennsylvania, Smithsonian Institution Washington D.C., 1961.
- _____. The biology of lichens. 2nd ed. London : William Clowes&sons, 1977.

- Hamada, N. and Ueno, T.. Lecanoric acid from the mycobiont of the lichen *Stereocaulon curtatum*. Phytochemistry. 292 (1990) : 678-679.
- Hamat, A.L.B., Din, L.B., Samsudin, M.W.B. and Felix, J.A.. Two new depsidones from the lichen. *Erioderma phaeorhizum* Vainio Sensu Lato. Aus. J. Chem. 461 (1993) : 153-156.
- Hernandez, D., Asis, V.C.. Plants of the philippines. Quezon City, The University of Philippines Press, 1971.
- Hirayama, T., Fujikawa, F., Yoshioka, I. and Kitagawa,I.. On the constituents of the lichen in the genus *Menegazzia*. menegazziaic acid. Chem. Pharm. Bull. 24 (1976) : 2340-2344.
- _____. Constituents of the lichen in the genus *Hypogymnia*. I. oxphysodic acid, a new depsidone from *Hypogymnia enteromorpha*. Chem. Pharm. Bull. 24 (1976) : 1596-1601.
- _____. Constituents of the lichen in the genus *Hypogymnia*. II vittantolic acid, a new optically active depsidone and 2'-O-methylphysodic acid from *Hypogymnia vittata*. Chem. Pharm. Bull. 24 (1976) : 1602-1608.
- _____. Vittatalic acid, a new depsidone isolated from the lichen *Hypogymnia vittata*. Chem. Pharm. Bull. 23 (1975) : 693-695.
- Holzmann, G. and Leuckert, C.. Application of negative fast atom bombardment and MS/MS to screening of Lichen compounds. Phytochemistry. 29 (1990) : 2277-2283.
- Huneck, S.. Lichen constituents. CIII. secondary constituents of an american lichen. Phytochemistry. 13 (1974) : 2872-2873.
- _____. Lichen constituents. part 111. constituents of *Pyxine coccifera*. Phytochemistry. 15 (1976) : 799-801.
- Huneck, S., Culberson, C.F., Culberson, W.L. and Elix, J.A.. Haematommone, a new

- Huneck, S., Culberson, C.F., Culberson, W.L. and Elix, J.A.. Haematommone, a new pigment from apothecia of *Haematomma puniceum*. Phytochemistry. 302 (1991) : 706-707.
- _____ and Lamb, I.M.. Lichen substances., part 106. 1'-chloropannarin, a new depsidone from *Argopsis friesiana* notes on the structure of pannarin and on the chemistry of the lichen genus *Argopsis*. Phytochemistry. 14 (1975) : 1625-1628.
- _____ and Schreiber, K.. Lichen constituents 102 secondary constituents from some european and indian lichens. Phytochemistry. 13 (1974) : 2315-2316.
- _____, Steglich, W. and Hofle, G.. Canarione, a new naphthoquinone from *Usnea canariensis*. Phytochemistry. 16 (1977) : 121-123.
- _____, Weinberg GDR and Mulholland G.A., "The absolute configurations of (+)-usnic acid and (+)-isousnic acid x-ray analyses of the (-)- α -phenylethylamine derivative of (+)-usnic acid and of (-)-pseudoplacodiolic acid, a new dibenzofuran, from the lichen. Tetrahedron Letters. 22 (1981) : 351-352.
- Hveding-Bergseth, N., Brunn, T. and Kjosen, H.. Isolation of 30-nor-21 alpha-hopan-22-one (isoadiantone) from the lichen *Platismatia glauca*. Phytochemistry. 228 (1983) : 1826-1827
- Hylands, D.J. and Ingolfsdottir, K. The isolation of methyl β -orsellinate from *Stereocaulon alpinum* and comments on the isolation of 4,6-dihydroxy-2-methoxy-3-methylacetophenone from *Stereocaulon* species. Phytochemistry. 24 (1985) : 127-129.
- Johnson, R.B., Feldott, G. and Lardy, H.A. The mode of action of the antibiotic usnic acid Arch. Biochem. 28 (1950) : 317.
- Jones, T.F. and Palmer, J.K.. Optical, crystallographic and x-ray diffraction data for usnic acid. J. Am. Chem. Soc. 72 (1950) : 1820-1822

- Keeton, F.J. and Keogh, F.M.. Caperatic acid from *Usnea alata*. Phytochemistry, 12 (1973) : 721-722.
- Keogh, F.M.. 2'-O-demethylpsoromic acid from *Usnea* sp.. Phytochemistry, 15 (1976) : 1801.
- _____ and Zurita, E.M.. α -(15-Hydroxyhexadecyl) itaconic acid from *Usnea aliphatica*. Phytochemistry, 16 (1977) : 134-135
- Kheir, Y.M. and Patel, M.B.. Isolation of (+)-usnic acid from Sudanese Drug *Usnea molliuscula*. Planta Med. 27(2) (1975) : 171-172.
- Krishna, D.R. and Venkataramana, D. Pharmacokinetics of d(+)-usnic acid in rabbits after intravenous and oral administration. Drug metabolism and disposition, 20 (1992) : 909-911.
- Kupchan, S.M. and Koppelman, H.L.. L-usnic acid : tumor inhibitor isolated from lichens. Experientia, 6 (1975) : 625.
- Lee, P. and Chantrapromma, K.. The major constituents of the lichen *Parmelia tinctorum* NYL. Warasan Songkla Nakkharin, 52 (1983) : 149-150.
- Lennette, E.H.,ed.. Manual of Clinical Microbiology, 4 th ed. Washington D.C. : American Society for Microbiology, 1985.
- Letcher, R.M.. Chemistry of lichen constituents-VI : mass spectra of usnic acid, lichenanthone and their derivatives. Organic mass spectrometry, 1 (1968) : 551-562.
- Lovell, C.R.. Plants and the skin. Oxford : Blackwell Scientific, 1993.
- Maass, W.S.G.. Lichen substances.V.methylated derivatives of orsellinic acid, lecanoric acid, and gyrophoric acid from *Pseudocyphellaria crocata*.Can. J. Bot. 53 (1973) : 1031-1039.

- Marshak, A. and Kuschner, M.. Action of streptomycin and usnic acid on the development of tuberculosis in guinea pigs. U. S. Publ. Health Rep. 65 (1950) : 131-144.
- Mateos, J.L., Conde, E., Miranda, T. and Vicente, C.. Regulation mechanisms of phenolic production in the lichen *Himantormia lugubris*, as reduced from the analysis of metabolite accumulation. Plant Sci. (Limerick Ire) 77 (1991) : 1-9.
- Mitchell, J.C.. Stereoisomeric specificity of usnic acid in delayed hypersensitivity. The Journal of Investigative Dermatology. (1966) : 167-168.
- _____ and Maibach, H.I.. Sensitizing capacity of usnic acid derived from lichenized fungi. Acta Derm. Venereol. 49 (1969) : 498-500.
- Neelakantan, S.. Recent developments in the chemistry of lichen substances Advancing frontiers in the chemistry of natural products. India : Hindustan, 1965.
- Nishikawa, Y., Ohki, K., Takahashi, K., Kurono, G., Fukuoka, F., and Emori, M.. Studies on the water soluble constituents of lichens. II. antitumor polysaccharides of *Lasallia*, *Usnea* and *Cladonia* species. Chem. Pharm. Bull. 22 (1974) : 2692-2702.
- Nishitoba, Y., Nishimura, H., Nishiyama, T. and Mizutani, J.. Lichen acids, plant growth inhibitors from *Usnea longissima*. Phytochemistry. 26 (1987) : 3181-3185.
- O'Donovan, D.G., Roberts, G. and Keogh, M.F.. Structure of the beta-orcinol depsidones, connorstictic acid and consalazinic acids. Phytochemistry. 19 (1980) : 2497-2499.
- Phrukudom, S.. M.Sc.Thesis. Dept. of Pharmacology, Faculty of Pharmaceutical Science, Chulalongkorn University, Thailand, 1993.
- Polonia, M.A.J., Carneiro A.F., Pinto, M. M.. Quantification usnic acid in *Usnea florida* by densitometric high performance thin layer chromatography. J. Liq. Chromatogr. 14 (1991) : 1447-1452.

- Posner, B., Feige, G.B. and Huneck, S.. Studies of the chemistry of the lichen genus *Umbilicaria* Hoffm.. Z. Naturforsch. Ser.C. 471/2 (1992) : 1-9.
- Quilhot, W., Piovano, M., Arancibia, H., Garbarino, JA. and Gambaro, V.. Studies on chilean lichens. XII chemotaxonomy of the genus *Psoroma*. J. Nat. Prod. 52(1989) : 191-192.
- Quilhot, W., Redon, J., Zuniga, E. and Vidal, S.. Depsides from *Lobodirina mahuiiana*. Phytochemistry. 14 (1975) : 1865B-1866.
- Round, F.E.. Introduction to the Lower Plant. London : The Camelot Press, 1969.
- Saklani, A. and Upreti, D.K.. Folk uses of some lichens in sikkim. J. Ethnopharmacol. 373 (1992) : 229-233.
- Sala, T., Sargent, M.V. and Elix, J.A.. Depsidone synthesis. part 15 new metabolites of the lichen *Buellia canescens* (Dicks.) De Not : novel phthalide catabolites of depsidones. J. Chem. Soc. Perkin Trans. I. 1981 (1981) : 849-854.
- Savich, V.P., Litvinov, M.A., and Moiseeva, E.N.. An antibiotic from lichens as a medicinal product. Planta Med. 8 (1960) : 191.
- Sevilla, S., and Patroinio.. Studies on philippine lichens III part II, chemical constituents and antibiotic activity of *Usnea montagnei*. Acta. Manilana, Ser. A. 29 (1980) : 130-149.
- Shaw, P.D.. Usnic acid. Antibiotic. 1 (1957) : 611-612.
- Siintola, S., Heilala, P., Patiala, R.. Usnic acid and its possible use in medicine. Soumen. Kemistilehti. 21A (1948) : 179-183.
- Solberg, Y.. Studies on the chemistry of lichens. XI. chemical investigation of five norwegian *Alectoria* species. Acta. Chem. Scand. Ser. B. 29 (1975) : 145-153.

- Solberg, Y.. Studies on the chemistry of lichens, XVI. chemical investigation of the lichen species *Alectoria ochroleuca*, *Stereocaulon vesuvianum* var. *pulvinatum* and *Icmadophila erectorum*. Z. Naturforsch. Ser. C. 32 (1977) : 182-189.
- _____ and Buzina, GV.. Studies on the chemistry of lichens, IVX. chemical investigation of the lichen species *Anaptychia fusca*, *Peltigera canina* and *Omphalodiscus spodochrous*. Z. Naturforsch. Ser. C. 30 (1975) : 445-450.
- _____ and Remedios, G.. Studies on the chemistry of lichens. XVIII. chemical investigation on the species *Letharia vulpina* and new derivatives of vulpinic acid. Z. Naturforsch. Ser. C. 33 (1978) : 449-451.
- Stark, B.J., Walter, D.E. and Owens, S.H.. Method of isolation of usnic acid from *Ramalina reticulata*. J. Am. Chem. Soc. 72 (1950) : 1819.
- Stoll, A., Brack, A., and Reng, J.. The antibiotic effect of usnic acid on mycobacteria and microorganisms. Experientia. 3 (1947) : 115-117.
- Swinscow T.D.V. and Krog H.. Microlichens of East Africa. London : Henry Ling Ltd., 1988.
- Taguchi, H., Sankawa, U. and Shibata S.. Biosynthesis of natural products. VI. Biosynthesis of usnic acid in lichens.(I).A general scheme of biosynthesis of usnic acid. Chem. Pharm. Bull. 17 (1969) 2054-2060.
- Takahashi, K. and Tanaka, Y.. Usnic acid XII.The oxidation of dihydrousnic acid. Chem. Pharm. Bull. 23(1975) : 623-628.
- Takai, M., Uehara, Y., and Beisler, J.A.. Usnic acid derivaitves as potential antineoplastic agents. J. Med. Chem. 22 (1979) : 1380-1384.
- Thune, P., Solberg, Y., McFadden, N., Staerfelt, F., and Sandberg, M.. Perfume allergy due to oak moss and other lichens. Contact Dermatitis. 8 (1982) : 396-400.

- Tomaselli, R.. New usnic acid derivative used therapeutically. Ed. Pract. Ed. Sci. 12 (1957) : 137.
- Vertia, K.O.. Antibiotics in lichens. Ann. Med. Exp. Biol. Fenn. 27 (1949) : 46-54.
- Virtanen, O.E. and Killpio, O.E.. Fungistatic activity of an usnic acid preparation with the trade name USNO. Suomen Kemistilehti. 30(1957) : 1-5.
- _____. Derivativs of usnic acid with the most important tuberculostatic agents. Suomen Kemistilehti. 28B (1955) : 125.
- Wilkins, A.L., Elix, J.A. , Gaul, K.L. and Moberg, R.. New hopane triterpenoids from lichens in the family physiaceae. Aust. J. Chem. 42 (1989) : 1415-1422.

ศูนย์วิทยบริการ
จุฬาลงกรณ์มหาวิทยาลัย

APPENDIX

ศูนย์วิทยบริการ
จุฬาลงกรณ์มหาวิทยาลัย

1)Chloroform : Benzene (3 : 2)




Figure 12 Thin-layer chromatogram of US-1

2) Petroleum ether : Ethyl acetate (9 : 1)




Figure 13 Thin-layer chromatogram of US-2

3) Chloroform : Methanol (9 : 1)




Figure 14 Thin-layer chromatogram of US-3




Figure 15 UV absorption spectrum of US-1 in MeOH




Figure 16 IR absorption spectrum of US-1 in KBr




Figure 17 Mass spectrum of US-1




Figure 18 $^1\text{H-NMR}$ spectrum of US-1 in CDCl_3





Figure 19 ^{13}C -NMR spectrum of US-1 in CDCl_3




Figure 20 ^{13}C -NMR spectrum of US-1 (DEPT 135) in CDCl_3




Figure 21 UV absorption spectrum of US-2 in MeOH




Figure 22 IR absorption spectrum of US-2 in Nujol




Figure 23 Mass spectrum of US-2




Figure 24 ^1H -NMR spectrum of US-2 in CDCl_3




Figure 24 ^1H -NMR spectrum of US-2 in CDCl_3




Figure 24 $^1\text{H-NMR}$ spectrum of US-2 in CDCl_3




Figure 25 ^{13}C -NMR spectrum of US-2 in CDCl_3




Figure 26 UV absorption spectrum of US-3 in MeOH




Figure 27 IR absorption spectrum of US-3 in Nujol




Figure 28 Mass spectrum of US-3




Figure 29 ^1H -NMR spectrum of US-3 in DMSO-d_6




Figure 30 ^{13}C -NMR spectrum of US-3 in DMSO- d_6



VITA

Miss Narisa Kamkaen was born on September 27, 1966 in Bangkok, Thailand. She received her Bachelor of Science in Pharmacy in 1989 from the Faculty of Pharmaceutical Sciences, Prince of Songkla University, Hat Yai, Thailand. At present, she is a food and drug technical officer in Narcotic Control Division, Food and Drug Administration, Thailand.

ศูนย์วิทยทรัพยากร
จุฬาลงกรณ์มหาวิทยาลัย