



References

- Allen, M.M. and Smith, A.J. 1969. Nitrogen chlorosis in blue-green algae. Arch. Mikrobiol. 69, 114-120.
- Bogorad L. 1975. Phycobiliproteins and complementary chromatic adaptation. Annu. Rev. Plant Physiol. 26 p.369-401.
- Boresch, K. 1922. Die Komplementare chromatische adaptation. Arch. Protistenkd. 44:1-70.
- Boussiba, S. and Richmond A.E. 1980. C-phycocyanin as a storage protein in the blue-green alga Spirulina platensis. Arch. Microbiol. 125. p.143-147.
- _____ and Richmond, A.E. 1979. Isolation and characterization of phycocyanin from the blue - green alga Spirulina platensis. Arch. Microbiol. 120. p.155-159
- _____ and Tel-Or, E. 1980. Production and chemicals from Spirulina platensis. Algae Biomass. (Shelef,G and Soeder, C.J.,eds.). Elsevier / Amsterdam. North-Holland Biomedical Press, p.612.
- Bennet, A. and Bogorad, L. 1971. Properties of subunits and aggregates of blue-freen algal biliproteins. Biochemistry. Vol.10, No.19, p.3625-3634
- Ciferri,O.1983. Spirulina, the edible microorganism. Microbiol.Rev. Dec, p.551-578.

- Conley, P.B., Lemaux, P.G. and Grossman A. 1988. Molecular characterization and evolution of sequences encoding light - harvesting component in the chromatically adaptation cyanobacterium Fremyella diplosiphon. J. Mol. Biol. 199. p.447- 465.
- Dainippon Ink and Chemicals Inc. 1980. Production of highly purified alcoholophilic phycocyanin. Japanese Patent. 80 77,890.
- De. Loura, I.C., Dubacq, J.P. and Thomas, J.C. 1978. The effect of nitrogen deficiency on pigment and lipid of cyanobacteria. Plant.Physiol. 83. p.838-843.
- Drews, G., and Weckesser, J. 1982. The biology of cyanobacteria., Carr, N. G. and Whitton, B.A., Eds. Bot.Monogr. 19. p.333-357.
- Emerson, R. and Lewis, C.M. 1942. The photosynthetic efficiency of phycocyanin in Chroococcus and the problem of caotenoid participation in photosynthesis. J.Gen.Physiol. 25. p. 57-99.
- Engelmann, T.W. 1902. Ueber experimentelle erzengung zweckmassiger aenderungen der farbung pflanzlieher chromophylle dyrch farbiges licht. Arch. Anat. Physiol. Physiol.Abt., p. 333-335.
- Fogg, G.E., 1966. Algal culture and phytoplankton ecology. Milwaukee and London, University of Wisconsin Press, Madison.

- Foulds, I. and Carr, N.G. 1977. A proteolic enzyme degrading phycocyanin in the cyanobacterium Anabaena cylindrica. FEMS microbiol.Lett., 2, 117.
- Francis, F.J. 1987. Lesser-known food colorants. Food Technology. April. p. 62-68.
- Gimmler, H., Wiedemann, C., and Moller, E.M. 1981. The metabolic response of the halotolerant green alga Dunaliella parva to hypertonic shocks. Ber.Dtsch.Bot.Ges.Bd., 94, 613.
- Glazer, A.N. 1984. Phycobilisome : A macromolecular complex optimized for light energy transfer. Biochimica et Biophysica Acta. 768. p.29-51.
- _____. 1988. Phycobiliprotein. Method of Enzymology. Vol 167. p.291-312. 52. p. 125- 157.
- _____. 1976. Phycocyanin : Structure and functions. Photochem.Photobiol.Rew. 1: 71-115.
- Healey, F.P. 1973. Characteristics of phosphorus deficiency in Anabaena. J.Phycol., 9. p. 383.
- Hoshaw, R.W. and Rosowski, J.R. 1973. Method for microscopic algae. Handbook of Phycological Methods and Growth. Measurement. (stein, J.R.,ed.) Cambridge Press.USA.p.54-67.
- Jassby, A. and Platte, T. 1976. Mathematical formulation of the relationship between photosynthesis and light for phytoplankton. Limnol.Oceanogr. 21, 540.

- Kao,O.H.W., Edwards,M.R. and Berns.D.S. 1975. Physical-chemical properties of c-phycocyanin isolated from an acido-thermophilic eukaryote,Cyanidium caldarium.Biochem.J. 147.p.63-70.
- Kaplan,D., Richmond,A., Dubinsky,Z., and Aaronson,S. 1986. Algal nutrition, in Handbook of Microalgal Mass Cultures, Richmond,A., Ed., CRC Press, Boca Raton , Fla., 147.
- Kawasaki,M. and Kaneko,T. 1978. Coloring of food. Jap.patent. 78,047,572.
- Kufer, W. and Scheer, H. 1979. Preparation and characterization of phycobiliproteins with chromophores chemically modified by reduction. Hoppe-Seyler's Z.Physiol.Chem. Bd.360,S.935-956.
- Kuhl, A., 1974. Phosphorus, in algal physiology and biochemistry., Stewart,W.D.P., Ed., Blackwell Scientific, Oxford, p. 636.
- Laemmli,U.K. 1970. Cleavage of structural protein during the assembly of the head of Bacteriophage T4. Nature. (London) 227,680
- Lau,R.H., Mackenzie, M.M., and Doolittle, W.F. 1976. Phycocyanin synthesis and degradation in the blue-green bacterium Anacystis nidulans Lemm, Arch.Microbiol. 107.p.15.
- Lowry,O.H., Rosenbrough,N.J., Farr,A.L. and Randall.R.J. 1951. Protein determination with the phenol reagent. J.Biol.Chem. 193,265-275.

M.Anusuya Devi. 1982. Biochemical and nutritional studies on algal protein, Thesis submit to University of Mysore for the degree of Doctor of Philosophy, Central Food Technology Research Institute Mysore- 570013, India, May 1982.

Mohleji,S.C. and verhoff,F.H.1980. Sodium and potassium ion effect on phosphorus transport in algal cells. J.Water Pollut. Control Fed. 52 (1), 110.

Reungjitchachawaly M., Vetayasuporn S., Chitnumsub P. and Tanticharoen M. 1989 .Abstract from 15 th Conference on Science and Technology of Thailand ,18-20 October 1989, Chiang Mai University.

Rippka,R.,Dernelles,J.,Waterbury,J.B.,Herdman,M. and Stanier,R.Y. 1979. Generic assignment, strain histories and properties of pure culture of cyanobacteria. J.Gen.Microbiol. 111.p.1-61.

Richmond,A.E. 1986. Microalgaculture.CRC Critical Reviews in Biotechnology. Vol.4, p.369-438.

Santillan,C. 1982. Mass production of Spirulina. Experientia. 38,Birkhauser verlag, CH-4010 Basel/Switzerland.

Tandeau De Marsac,N. and Cohen-Bazire,G. 1977. Molecular composition of cyanobacterial phycobilisomes. Proc.Natl.Acad.Sci.USA. 74,No.4,p.1635-1639.

1977. Occurrence and nature of chromatic adaptation in cyanobacteria. J.Bacteriol. Vol.130, No.1 p. 82-91.
- Vaccaro,R.F. 1965. Inorganic nitrogen in sea water. Chemical Oceanography ,Vol.1, Riley,J.P. and Skirrow,G., Eds., Academic Press, London.p.365.
- Venkataraman,L.V. 1983. A monograph on Spirulina platensis. CFIRI Press,Mysore,India.
- and Becker,E.W. 1985. Biotechnology and utilization of algae. The Indian Experience. Sharada Press. Bangalore.
- Verity,P.G. 1981. Effects of temperature, irradiance and daylength on the marine diatom Leptocylindrus danicus cleve.I. Photosynthesis and cellular composition. J.Exp.Mar.Biol.Ecol. 55,79.
- Yamanaka,S., Chiba,M. and Morinaga,Y. 1978. Algal pigment stabilization. Jap.patent. 78,101,532.
- Wood,N.B. and Haselkorn,R. 1976. Protein degradation during heterocyst development in Anabaena , in Proc.2nd Int. Symp.Photosynthesis Prokaryotes, Codd, G. and Stewart, W.D., Eds., Dundee Scotland,125.
- Zafaralla, M.T.,Barril, C.R., Vidal, L.R., Serrana, M.S.,Aguila,L.G. and Tansinsin,L.G. 1985. Protein from Spirulina. NSTA Technology J. July-Sept. p.18- 26.

Zarrouk , C. 1966. Contribution a l'étude d'un cyanophycee. Influence de divers facteurs physiques et chimiques sur la croissance et la photosynthèse de Spirulina maxima. Ph.D thesis, Paris.



ศูนย์วิทยทรัพยากร
จุฬาลงกรณ์มหาวิทยาลัย

APPENDIX

Zarrouk medium (Zarrouk, 1966) contained the follow components in gram per liter.

NaHCO ₃	16.80
NaNO ₃	2.50
K ₂ HPO ₄	0.50
NaCl	1.00
MgSO ₄ .7H ₂ O	0.20
FeSO ₄ .7H ₂ O	0.01
K ₂ SO ₄	1.00
CaCl ₂ .2H ₂ O	0.04
EDTA	0.08
A ₅ Solution	1 ml/l
B ₆ Solution	1 ml/l
pH	8-10

The A₅ Solution contained the following in g/l ; H₃BO₄ : 2.86 ; MnCl₂.4H₂O : 1.81 ; ZnSO₄.7H₂O : 0.22 ; MoO₃ : 0.01 ; and CuSO₄.5H₂O : 0.08.

The B₆ Solution contained the following in mg/l; NH₄VO₃ : 22.9 ; NiSO₄.7H₂O : 47.8 ; Na₂WO₄ : 17.9 ; Ti(SO₄)₂ : 40.0 ; and Co(NO₃)₂.6H₂O : 4.4.

In order to prevent the reactions of some components causing the precipitation of the medium, NaHCO₃ and K₂HPO₄ were autoclaved separately. The medium was sterilized by autoclaving at 15 lb/in² for 15-20 min and the solutions were mixed together.

BIOGRAPHY

Miss Duangrat Inthorn was born on March 8, 1965 in Bangkok, Thailand. She graduated with the Bachelor of Science Degree in Microbiology from Faculty of Science, Chulalongkorn University in 1988.



ศูนย์วิทยทรัพยากร
จุฬาลงกรณ์มหาวิทยาลัย