



## REFERENCES

### ภาษาไทย

เต็ม สมิตินันท์. ชื่อพรรณไม้แห่งประเทศไทย (ชื่อพุกฤษศาสตร์-ชื่อพื้นเมือง). กรุงเทพ : พิพิธภัณฑ์พันลิขซึ่ง, 2523.

เสงี่ยม พงษ์บุญรอด. ไม้เทศ เมืองไทย. กรุงเทพ: เกษมบรรณกิจ, 2522.

### ภาษาอังกฤษ

Akasu, M., Itokawa, H., and Fujita, M. Oxoaporphine alkaloids from callus tissues of *Stephania cepharantha*. *Phytochemistry* 14 (1975) : 1673-1674.

\_\_\_\_\_, Itokawa, H., and Fujita, M. Biscoclaurine alkaloids in callus tissues of *Stephania cepharantha*. *Phytochemistry* 15 (1976) : 471-473.

Amann, M., Wanner, G., and Zenk, M.H. Methyl transfer in formation of aporphine and Protoberberine alkaloids. *Planta*. 167 (1986) : 310-320.

Amarendra, P., et al. The protoberberine alkaloids of *Stephania suberosa*. *Phytochemistry* 26 (1987) : 547-549.

Baarschers, W.H., Arudt, R.R., Pachler, K., Weisbach, J.A., and Douglas, B. Nuclear magnetic resonace study of aporphine alkaloids. *J. Chem. Soc.* (1964) : 4778-4782.

Banerji, J., et al. Kamaline, an unusual aporphine alkaloid from *Stephania venosa*. *Phytochemistry* 36 (1994) : 1053-1054.

Bartley, J.P., Baker, L.T., and Carvalho, C.F. Alkaloids of *Stephania bancroftii*. *Phytochemistry* 36 (1994) : 1327-1331.

Barton, D.H.R., Hesse, R.H., and Kirby, G.W. Phenol oxidation and biosynthesis, part VIII. Investigations on the biosynthesis of berberine and protopine. *J. Chem. Soc.* (1965) : 6379-6389.

- Battersby, A.R., Binks, R., Francis, R.J., McCaldin, D.J., and Ramuz, H. Alkaloid Biosynthesis, part IV. 1-Benzylisoquinolines as precursors of thebaine, codeine and morphine. *J. Chem. Soc.* (1964) : 3600-3610.
- Bhakuni, D.S., and Gupta, S. The alkaloids of *Stephania glabra*. *J. Nat. Prod.* 45 (1982) : 407-411.
- \_\_\_\_\_. Tewari, S., and Kapil, R. S. Biosynthesis of boldine (1,10-dimethoxy-6- $\alpha$ -aporphine-2,9-diol). *J. Chem. Soc.* 1 (1977) : 706-709.
- Bick, I.R.C., Harley-Mason, J., Shrppard, N., and Vernengo, M.J. Structure correlations in the nuclear magnetic resonance spectra of bisbenzylisoquinoline and aporphine alkaloids. *J. Chem. Soc.* (1961) : 1896-1903.
- Brochmann-Hanssen, E., Chen, C.H., Chiang, H.C., Fu, C.C., and Nemoto, H. Opium alkaloids XIV : Biosynthesis of Aporphines-detection of orientaline in opium poppy. *J. Pharm. Sci.* 63 (1971) : 1291-1293.
- Charls, B., et al. Some unusual proaporphine and aporphine alkaloids from *Stephania venosa*. *J. Nat. Prod.* 50 (1987) : 1113-1117.
- Charlwood, V. B., and Rhodes, M. J. C. **Physiology of secondary product formation in plants**. Oxford : Clarendon Press, 1990.
- Cordell, G.A. **Introduction to alkaloids : A biogenic Approach**. New York : John Wiley and Sons : 1981.
- Craib, W.G. Contributions to the Flora of Siam. Additamentum XIII. *Bull. Misc. Inf., R.Bot. Gard.* (1922) : 225-241.
- \_\_\_\_\_. **Flora Siamensis Enumeratio : A List of the Plants Known from Siam with Records of Their Occurrence**. Bangkok : Bangkok Times Press, 1931.

- Deng, J.Z., and Zhao, S.X. Alkaloids from aerial parts of *Stephania longa*. **Phytochemistry** 33 (1993) : 941-942.
- Diels, L. Menispermaceae. In A. Engler (ed.), **Das Pflanzenreich : Regni vegetabilis conspectus**, pp.345. Weinheim : H.R. Engelmann, 1910.
- Forman, L.L. **Studies in the Flora of Thailand**. Odense : Andelsbogtrykkeriet, 1962.
- \_\_\_\_\_. Menispermaceae. **Flora of Thailand** 5 (1991) : 300-365.
- Furuya, T., Yoshikawa, T., and Kiyohara, H. Alkaloid production in cultured cells of *Dioscoreophyllum cumminsii*. **Phytochemistry** 22 (1983) : 1671-1673.
- Galneder, E., Rueffer, M., Wanner, G., Tabata, M., and Zenk, M.H. Alternative final steps in berberine biosynthesis in *Coptis japonica* cell cultures. **Plant Cell Reports** 7 (1988) : 1-4.
- Gamborg, O.L. The effects of amino acids and ammonium on the growth of plant cells in suspension. **Physiol. Plant.** 45 (1970) : 372-375.
- \_\_\_\_\_, Miller, R.A., and Ojima, K. Nutrient requirement of suspension cultures of soy bean root cells. **J. Exp. Cell. Res.** 50 (1968) : 151-158.
- Guinaudeau, H., Leboeuf, M., and Cave, A. Aporphine Alkaloids. **Lloydia** 38 (1975) : 275-338.
- \_\_\_\_\_. Aporphine Alkaloids, II. **J. Nat. Prod.** 42 (1979) : 325-360.
- \_\_\_\_\_. Aporphinoid Alkaloids, III. **J. Nat. Prod.** 46 (1983) : 761-835.
- \_\_\_\_\_. Aporphinoid Alkaloids, IV. **J. Nat. Prod.** 51 (1988) : 389-474.
- \_\_\_\_\_. Aporphinoid Alkaloids, V. **J. Nat. Prod.** 57 (1994) : 1033-1135.

- Gurova, T.F., Popov, Y.G., Fadeeva, I.I., Shain, S.S., and Tokachev, O. Callus formation and accumulation of alkaloids in the tissue culture of *Stephania glabra*. **Rastit. Resur.** 16 (1980) : 412-425.
- Herbert, R.B. **The biosynthesis of secondary metabolites**. New York : Champman and Hall Co., 1981.
- Ikuta, A., and Itokawa, H. Berberine and other protoberberine alkaloids in callus tissue of *Thalictrum minus*.. **Phytochemistry** 21 (1982a) : 1419-1421.
- \_\_\_\_\_. Studies on the alkaloids from tissue culture of *Nandina domestica*. Plant tissue culture 1982, 5th Int. Congr. **Plant Cell Culture** (1982b) : 315-316.
- \_\_\_\_\_, Syona, K., and Furuya, T. Alkaloids of callus tissue and redifferentiated plantlets in the Papaveraceae. **Phytochemistry** 13(1974) : 2175-2179.
- Kunitomo, J., et al. The alkaloids of *Stephania sasakii* : Structure of five new alkaloids. **Phytochemistry** 19 (1980) : 2735-2739.
- \_\_\_\_\_. 4-Hydroxycrebanine a new 4-hydroxyaporphine alkaloids and (R)-roemeroline from *Stephania sasakii*. **Chem. Pharm. Bull.** 29 (1981) : 2251-2253.
- Lane, W. D. Regeneration of pear plants from shoot meristem tips. **Plant. Sci. Lett.** 13 (1979) : 337-342.
- Lao, A., Tang, Z., and Xu, R. Studies on the chemical constituents of *Stephania longa*. **Chem. Abstr.** 96 (1982) : 196523.
- Likhitwitayawuid, K., Angerhofer, C.K., Cordell, G.A., and Pezzuto, J.M. Cytotoxic and antimalarial bisbenzylisoquinoline alkaloids from *Stephania erecta*. **J. Nat. Prod.** 56 (1993a) : 30-38.

- \_\_\_\_\_, Chai, H., Pezzuto, J.M. and Cordell, G.A. Cytotoxic and antimalarial alkaloids from the tubers of *Stephania pierrei*. **J. Nat. Prod.** 56 (1993b) : 1468-1478.
- Linsmaier, E.M., and Skoog, F. Organic growth factor requirements of Tobacco tissue cultures. **Physiol. Plant.** 18 (1965) : 100-127.
- \_\_\_\_\_. Plant Tissue Culture Methods. **Physiol. Plant.** 18 (1965) : 100-127.
- Loeffler, S., Stadler, R., Wagakura, N., and Zenk, M.H. Norcoclaurine as biosynthetic precursor of thebaine and morphine. **J. Chem. Soc., Chem. Commun.** (1987) : 1160-1162.
- Loomis, W.D., and Battaile, J. Plant phenolic compounds and the isolation of plant enzymes. **Phytochemistry** 5 (1966) : 423-438.
- Luckner, M. **Secondary metabolism in microorganisms, plants and animal.** 3rd ed. Berline : INTERDRUCK, 1990.
- Matsui, M., Kabashima, T., Ishida, K., and Takebayashi, T. Alkaloids of the leaves of *Stephania japonica*. **J. Nat. Prod.** 45 (1982) : 497-500.
- Menachery, M.D., et al. Simple isoquinoline alkaloids. **J. Nat. Prod.** 49 (1986) : 745-778.
- Muller, M.J., and Zenk, M.H. The Norcoclaurine pathways in operative in Berberine biosynthesis in *Coptis japonica*. **Planta Med.** 58 (1992) : 524-527.
- Murashige, T., and Skoog, F. A revised medium for rapid growth and bioassay with tobacco tissue cultures. **Physiol. Plant.** 15 (1962) : 473-497.
- Nisch, J.P., and Nitsch, C. Haploid plants from pollen grain. **Science** 163 (1969) : 85-87.
- Patra, A., et al. The bisbenzylisoquinoline alkaloids of *Stephania suberosa*. **J. Nat. Prod.** 49 (1986) : 424-427.

- \_\_\_\_\_, Ghosh, A., and Metra, A.K. Alkaloids of *Stephania glabra*. **Planta Med.** 40 (1980) : 333-336.
- Pharadai, K., et al. (-)-*O*-Acetylsukhodianine and oxostephanosine : Two new aporphinoids from *Stephania venosa*. **J. Nat. Prod.** 49 (1965) : 658-659.
- Prawat, U., Wiriachitra, P., Lojanapiwatra, V., and Nimgirawath, S. Alkaloids of *Stephania erecta* Craib. **J. Sci. Soc. Thailand** 8 (1982) : 65-68.
- Robinson, R. **The Structural Relations of Natural Products**. Oxford : Clarendon Press, 1955.
- Rueffer, M., and Zenk, M.H. Revision of early steps of reticuline biosynthesis. **Tetrahedron Lett.** 26 (1985) : 201-202.
- Sakuta, W. D., and Komamine, A. Cell Growth and Accumulation of Secondary Metabolites. In F. Constabel and I. K. Vasil (eds.), **Cell Culture and Somatic Cell Genetics of Plants**, pp. 97-110. New York : Academic Press, 1987.
- Schneider, B., and Zenk, M.H. Metabolism of secondary products in cell systems. In K. Oono et al. (eds.), **Plant tissue culture and gene manipulation for breeding and formation of phytochemicals**, pp. 323-333. Mito : Sato Press, 1985.
- \_\_\_\_\_. Unusual methyl transfer in the isoquinoline of aporphine and protoberberine alkaloids. **Phytochemistry** 32 (1993) : 897-903.
- Shamma, M. **The isoquinoline alkaloids chemistry and pharmacology**. New York and London : Academic press, 1972.
- \_\_\_\_\_, and Maniot, J.L. **Isoquinoline alkaloid research 1972-1977**. London : Plenum, 1978.

Singh, R.S., Kumar, P., and Bhaduni, D.S.: The alkaloids of *Stephania elegans*. **J. Nat. Prod.** 44 (1981) :182-185.

Spenser, I.O. Biosynthesis of isoquinoline alkaloids. In M. Florkin and E.H. Stotz (eds.), **Comprehensive biochemistry**, Vol XX, pp. 231-232. Amsterdam : Elsevier, 1968.

Stadler, R., et al. Revision of the early steps of reticuline biosynthesis **Tetrahedron Letters** 28 (1987) : 1251-1254.

\_\_\_\_\_, Kutchan, T.M., and Zenk, M.H. (S)-Norcoclaurine is the central intermediats in benzylisoquinoline alkaloids biosynthesis. **Phytochemistry** 28 (1989) : 1083-1086.

Stockigt, J. The biosynthesis of heteroyohimbine-type alkaloids. In J.D. Phillipson and M.H. Zenk (eds.), **Indole and biogenetically related alkaloids**, p. 113-114. London : Academic Press, 1980.

Sugimoto, Y., Sugimura, Y., and Yamada, Y. Effects of culture conditions on Bisbenzylisoquinoline Alkaloids Production in cultured roots of *Stephania cepharantha*. **Agric. Biol. Chem.** 52 (1988a) : 1495-1498.

\_\_\_\_\_. Production of Bisbenzylisoquinoline alkaloids in cultured roots of *Stephania cepharantha*. **Phytochemistry** 27 (1988b) : 1379-1381.

\_\_\_\_\_, and Yamada, Y. Involement of tyramine in bisbenzylisoquinoline biosynthesis in cultured roots of *Stephania cepharantha*. **J. Nat. Prod.** 56 (1993) : 1229-1233.

Tantisewie, B., Amurrio, S., Guinaudeau, H., and Shamma, M. New bisbenzylisoquinolines from *Stephania pierrei*. **J. Nat. Prod.** 52 (1989) : 846-851.

Thornber, C.W. Alkaloids of the Menispermaceae. **Phytochemistry** 9 (1970) : 157-187.

- Webb, W. D., and Street, H. E. Morphogenesis in vitro of Pinus and Picea.  
*Acta Horticulturae* 78 (1977) : 259-269.
- White, P.R. **The cultivation of animal and plant cells.** New York : Ronald Press, 1963.
- Yamamura, Y., and Matsui, M. Alkaloids from the fruits of *Stephania japonica*, structure of oxostephabenine and *N,O*-dimethyloxostephine. *J. Nat. Prod.* 48 (1980) : 746-750.
- Yeoman, M. M. **Plant Cell Culture Technology.** vol 23. Guildford : Biddles Ltd., 1986.
- Zenk, M.H. The impact of plant cell culture on industry. **Publ. Int. Assoc. Plant Tissue Culture**, p. 1. Calgary : 1978.
- \_\_\_\_\_. Enzymology of Benzylisoquinoline Alkaloids Formation. In J.D. Phillipson, M.F. Roberts, and M.H. Zenk (eds.), **The Chemistry and Biology of Isoquinoline Alkaloids**, pp. 240-256. Berlin : Springer-Verlg, 1985.

ศูนย์วิทยทรัพยากร  
 จุฬาลงกรณ์มหาวิทยาลัย



## VITA

Mr. Charlee Thongruang was born on June, 12 1966 in Singburi, Thailand. He received his Bachelor of Pharmaceutical Science in 1990 from Chulalongkorn University , Bangkok, Thailand. At present, he has been appointed as a student in University Development Commission (UDC) Program of the Department of Pharmaceutical Chemistry and Pharmacognosy, Faculty of Pharmaceutical Science, Naresuan University, Pisanulok, Thailand.

ศูนย์วิทยทรัพยากร  
จุฬาลงกรณ์มหาวิทยาลัย