

REFERENCES

- Abdel-Kawi, M., Slatkin, D.J., Schiff, P.L. Additional alkaloids of *Pachygone ovata*. J. Nat. Prod. (1984): 459-464.
- Amarendra, P., et al. The protoberberine alkaloids of *Stephania suberosa*. Phytochemistry. 26 (1987): 547-549.
- Backer, C.A. and Evan Den-Brind, R. Flora of Java. 1 (1953):156.
- Beecher, C.W.W. , and Kelleher, W.J.. Enzymatic study of the late stages of protoberberine alkaloid biosynthesis. Tetrahedron letter 25 (1984):538-542.
- Beek, T., Verpoorte, R., and Baerheim Svendsen, A.. Colorflammine and berbacolorflammine, two new orange-caloured bis [benzylisoquinoline] alkaloids from *Pycnarrhena longifolia*. J. Org. Chem. 47 (1982) : 898.
- Bembenek, M.E., et al. Inhibition of monoamine oxidases A and B by simple isoquinoline alkaloids. J.Med. Chem. 33 (1990) ; 147-152.
- Bhakuni, D.S.. Some aspects of protoberberine alkaloids. J. Indian Chem. Soc. 61 (1984) : 1016-1022.
- Bhakuni, D.S., and Jain, S.. Protoberberine alkaloids. J. Indian. Chem. Soc. 61 (1984) : 95-181.
- Bhakuni, D.S. and Gupta, S.. The alkaloids of *Stephania glabra*. J. Nat. Prod. 45 (1982): 407-411.

Blasko, G. et al.. ^{13}C -NMR assignments of berberines.

Heterocycles. 27 (1988) : 911-916.

Cave, A., Leboeuf, M., Hocquemiller, R., Bouquet, A.

Alkaloids du *Limaciopsis loangensis*. Planta Med.

35 (1978) : 31-41.

Chang, H.M., El-Fishawy, A.M., Slatkin, D.J., and Schiff, P.L.. Quaternary alkaloids of *Tinospora capillipes*.

Planta Med. 1984 : 35-38.

Charls, B., et al. Some unusual proaporphine and aporphine alkaloid from *Stephania venosa*. J. Nat. Prod. 50 (1987) : 1113-1117.

Clark, E.P.. Preparation of picrotoxin. J. Am. Chem. Soc. 57 (1935) : 1111.

Cordell, G.A.. Introduction to alkaloids : A biogenetic approach. New York : A wiley-interscience, 1981.

Dwuma-Badu, D., Ayim, J.S.K., and Fiagbe, N.Y.. Constituents of West African Medicinal Plants XV. Dinlkacorine, a new biphenyl dibenxodioxin alkaloid from *Tiliacora clinlkagei*. Lloydia. 39 (1973) : 213-217.

Forman, L.L.. A revision of the tribe coscinieae. Hook.f. & Thoms. (Menispermaceae). Kew. Bull. 32 (1978) : 323-338.

Forman, L.L.. The correct names for the tribes of Menispermaceae. Kew. Bull. 37 (1982) : 368.

Forman, L.L.. Menispermaceae in Flora Malesiana. 10² (1986) : 157-253.

Forman, L.L.. Menispermaceae in Flora of Thailand. 5 (1992) : 300-365.

Garcia, L.M., et al. The alkaloids of *Archangelisia laureirii* and *Coscinium wallichianum*. Phytochemistry: 9 (1970) : 663-664.

Guinaudeau, H., Shamma, M., Tantisewie. B., Pharadai, K. Aporphine alkaloids oxygenated at C-7. J. Nat. Prod. 45 (1983) : 335-337.

Guinaudeau, H. Aporphinoid alkaloids III. J. Nat. Prod. 46 (1983) : 761-835.

Guinaudeau, H., et al. Bisbenzylisoquinoline alkaloid from *Cocculus pendulus*. Phytochemistry 26 (1986):829-832.

Hegnauer, R.. Chemotaxonomie der pflanzen. 5 (1969):428-431.

Hussain, R.A., Kim, J.W., Beechu, C.W.W., and Kinghorn, D.. Unambiguous carbon-13 NMR assignments of some biologically active protoberberine alkaloids. Heterocycles. 29 (1989) : 2257-2260.

Kametani, T., et al. Conformational analysis of tetrahydroprotoberberines by carbon-13 NMR spectroscopy. Heterocycles 3 (1975) : 371-377.

Kunitoma, J., et al. The alkaloids of *Stephania sasakii* : Structure of five new alkaloids. Phytochemistry 19 (1980) : 2735-2739.

Kunitoma. 4-Hydroxycrebanine a new 4-hydroxyaporphine alkaloid and (R)-roemeroline from *Stephania sasakii*. Chem. Pharm. Bull. 29 (1981) : 2251-3.

Kupchan, M.S., Liepa, J.A.; Baxter, L.R.; and Hintz, P.J.. New alkaloids and related artifacts from *Cyclea peltata*. J. Org. Chem. 38 (1973) : 321-322.

Lao, A.; Tang, Z, and Xu, R.. Studies on the chemical constituents of *Stephania longa*. Chem. Abstr. 96 (1982) : 196523.

Leboeuf, M., et al. The phytochemistry of the annonaceae. Phytochemistry 21 (1982) : 2783-2813.

Leboeuf, M., Abouchara, M.L., Sevenet, T., Cave, A.. Alkaloids of *Albertisia papuana* Becc. Plant. Med. 16 (1982) : 280-291.

Leboeuf, M., Abouchara, M.L., Cave, A., Debray, M.. Alkaloids of *Pachygone loyaltiensis* Diels. Plant. Med Phytothr. 21 (1987) : 106-155.

Malhotra, S., Taneja, S.C., and Dhar, K.L.. Minor alkaloids from *Coscinium fenestratum*. Phytochemistry 28 (1989) : 1998-1999.

Matsui, M., Kabashima, T., Ishida, K., and Takebayashi, T. Alkaloids of the leaves of *Stephania japonica*. J. Nat. Prod. 45 (1982) : 497-500.

Menachery, M.D., et al. Simple isoquinoline alkaloids. J. Nat Prod. 49 (1986) : 745-778.

Monkovic, I., and Spenser, T.D.. Biosynthesis of berberine incorporation of catecholamines. Can. J. chem. 43 (1965) : 2017-2026.

Ohiri, F.C., Verpoorte, R., Baerheim Svendsen, A.. Alkaloids from *Chasmanthera dependens*. Planta Med. 46 (2983): 228-230.

Ohiri, F.C.; Verpoorte, R.; and Baerheim Svendsen, A.. Cycleanine N-oxide, a new alkaloid from *Synclisia scabrida*. Planta Med. 47 (1983) : 87-89.

- Ohiri, F.C.; Verpoorte, R.; and Baerheim Svendsen, A.. Tertiary phenolic alkaloids from *Chasmanthera dependens*. Planta Med. 49 (1983) : 17-19.
- Ohiri, F.C.; Verpoorte, R., Baerheim Svendsen, A.. Proton NMR chemical values for aromatic proton in 2,3,9,10- and 2,3,10,11-tetrasubstituted tetrahydroprotoberberine alkaloid. Planta Med. 49 (1983) : 162-164.
- Owusu, P.D., Slatkin, D.J.; Inapp, J.E; and Schiff, P.L.. Constituents of West African Medicinal Plant XXVIII. Additional alkaloids of *Triclisia giliett*. J. Nat. Prod. 44 (1981) : 63-65.
- Pachaly, Alkaloid from *Tiliacora triandra* Diels. VII. Tilianagine, a new bisbenzylisoquinoline. Arch. Pharm. 319 (1986): 872-877.
- Pachaly, P., and Tjie, J.. Alkaloids from *Tiliacora triandra* Diels. II. Yanangine, a new bisbenzylisoquinoline alkaloids. Arch. Pharm 319 (1986) : 841-849.
- Patra, A., Ghosh, A., and Metra, A.K.. Alkaloids of *Stephania glabra*. Planta Med. 40 (1980) : 333-336.
- Patra, A., et al. The bisbenzylisoquinoline alkaloids of *Stephania suberosa*. J.Nat.Prod. 49 (1986) : 424-427.
- Pavelka, S., and Kova, R.J.. Interaction of liver alcohol dehydrogenase with protoberberine alkaloids. Collect. Czech Chem Commun. 40 (1975) : 753-768.
- Pharadai, K., et al. The absolute configuration of (-) crebanine. Heterocycles 15 (1981) : 1067-1068.

- Pharadai, K., et al. (-)-O-Acetylsukhodianine and oxostephanosine : Two new aporphinoids from *Stephania venosa*. J. Nat. Prod. 49 (1965) 658-659.
- Ruangrungsi, N., Lange, G.L. and Lee, M.. Constituents of *Parabaena sagittata*. Two new tetrahydroprotoberberine alkaloids. J. Nat. Prod. 49(1986):253-258.
- Sethi, M.L.. Enzyme inhibition VI : inhibition of reverse transcriptase activity by protoberberine alkaloid and structure activity relationships. J. Pharm. Sc. 72 (1983) : 538-541.
- Sethi, M.L.. Comparision of inhibition of reverse transcriptase and antileukemic activities exhibited by protoberberine and benzophenanthridine alkaloids and structure activity relationship. Phytochemistry 24 (1985) : 447-454
- Shamma, M.. The isoquinoline alkaloids chemistry and pharmacology. New York and London : academic press 1972.
- Shamma, M., and Maniot, J.L.. The systematic classification of bisbenzylisoquinoline. Heterocycles 4 (1976) : 1817-1824.
- Simieon, S., Rios, J.L., Villar, A.. Pharmacological activities of protoberberine alkaloids. Plant. Med. Phytothr. 23 (1989) : 202-250.
- Singh, R.S., Kumar, P., and Bhakuni, D.S.. The alkaloids of *Stephania elegans*. J. Nat. Prod. 44 (1981) 182-185.

- Siwon, J., Verpoorte, R., Van Essen, G.F.A. and Baerheim Svendsen, A.. Studies on Indonesian medicinal plants. Planta Med. 38 (1980) 24-32.
- Siwon, J., Verpoorte, R., and Baerheim Svendsen, A. Dihydrodiscretine, an quaternary protoberberine alkaloids from *Heptacyclum zenkeri*. Planta Med. 41 (1981) : 65-68.
- Siwon, J., Verpoorte, R. and Baerheim Svendsen, A.. Studies on Indonesian medicinal plants VI. Further alkaloid from *Fibraurea chloroleuca*. Planta Med. 41 (1981) : 65-68.
- Siwon, J., et al. The studies of Indonesian medicinal plants VII : Further alkaloids from *Arcangelisia flava*. Planta Med. 43 (1982): 72-74.
- Southon, I.W. and Buckingham, J. (eds). Dictionary of alkaloids. London: Chapman and Hall 1989.
- Spiff, A.I.. Constituents of West African medicinal plants. Tridictyophlline, a new morphinan alkaloids from *Trilisia dictyophylia*. J. Nat. Prod. 44 (1981) : 161-165.
- Suffness, M., and Cordell, G.A.. The alkaloids. Academic press, New York. 25 (1985).
- Takani, M., Takasu, Y., and Takahashi, K.. Studies on constituents of the vines of *Menispermum dauricum* DC. Chem. Pharm. Bull. 11 (1983) : 3091-3093.

Tantisewie, B., et al.. (+) - N - Formylnornantenine, a new aporphine alkaloid from *Cyclea atjehensis*. J. Nat. Prod. 52 (1989) : 653-654.

Tantisewie, B., et al. A new subgroup of bisbenzylisoquinoline alkaloid (+) cycleatjehenine and cycleatjehine. Tetrahedron 46 (1990) ; 325-330.

Thornber, C.W.. Alkaloids of the Menispermaceae. Phytochemistry 9 (1970) : 157 - 187.

Verpoorte, R., Rijzen, A.H.M., Siwon, J., and Baerheim Svensen, A.. Tertiary alkaloids of *Pycnarrhena novoquinensis*. Planta Med. 34 (1978) : 274-279.

Verpoorte, R., et al. Studies on Indonesian medicinal plants V. The alkaloids of *Anamirta coccus*. J. Nat. Prod. 44 (1981) : 221-224.

Verpoorte, R., et al. Studies on Indonesian medicinal plants VII. Alkaloids of *Arcangelisia flava*. J. Nat. Prod. 45 (1982) : 582-584.

Yamamura, Y., and Matsui, M.. Alkaloids from the fruits of *Stephania japonica*, structrue of oxostephabenine and N,O-dimethyloxostephine. J. Nat. Prod. 48(1985): 746-750.

APPENDIX




Figure 2 *Coscinium fenestratum* Colebr.

1) Chloroform : Methanol (9:1)




Figure 3 Thin layer chromatogram of the isolated alkaloids,
Cf-1 to Cf-4

2) Chloroform : Methanol : Ethyl acetate (7:2:1)




Figure 4 Thin layer chromatogram of the isolated alkaloids,
Cf-1 to Cf-4

3) Chloroform : Acetone (6:4)




Figure 5 Thin layer chromatogram of the isolated alkaloids,
Cf-1 to Cf-4

4) n-Hexane : Methanol (7:3)




Figure 6 Thin layer chromatogram of the isolated alkaloids,
Cf-1 to Cf-4

5) Chloroform : Methanol : Acetone (5:3:2)




Figure 7 Thin layer chromatogram of the isolated alkaloids,

Cf-1 to Cf-4




Figure 8 IR absorption spectrum of Cf-1 in KBr




Figure 9 ^1H NMR spectrum of Cf-1 in CDCl₃





Figure 10 ^{13}C NMR spectrum of Cf-1 in CDCl_3




Figure 11 Mass spectrum of Cf-1





Figure 12 ^1H -NMR spectrum of Cf-2 in DMSO-d₆




Figure 13 Mass spectrum of Cf-2




Figure 14 IR absorption spectrum of Cf-2 in KBr




Figure 15 ^{13}C -NMR spectrum of Cf-2 in DMSO-d_6




Figure 16 ^1H -NMR spectrum of Cf-3 in DMSO-d_6




Figure 17 Two dimension ^1H -NMR spectrum (COSY) of Cf-3

in DMSO-d_6




Figure 18 ^{13}C -NMR spectrum of Cf-3 in DMSO-d_6




Figure 19 ^{13}C -NMR spectrum of Cf-3 (DEPT. 90) in DMSO-d_6




Figure 20 ^{13}C -NMR spectrum of Cef-3 (DEPT. 135) in $\text{DMSO}-\text{d}_6$




Figure 21 IR absorption spectrum of Cf-3 in KBr






Figure 22 Mass spectrum of Cf-3

Figure 23 ^1H -NMR spectrum of Cf-4 in DMSO-d_6






Figure 24 ^{13}C -NMR spectrum of Cf-4 in DMSO- d_6




Figure 25 IR absorption spectrum of Cf-4 in KBr




Figure 26 Mass spectrum of Cf-4




Figure 27 ^1H -NMR spectrum of reduced Cf-2 in $\text{DMSO}-\text{d}_6$




Figure 28 ^1H -NMR spectrum of reduced Cf-4 in $\text{DMSO}-\text{d}_6$




Figure 29 Mass spectrum of reduced Cf-2



VITA

Name Mrs. Supranee Keawpradub.

Education B.Sc. in pharm., Faculty of Pharmaceutical Science, Prince of Songkla University, Haadyai, Songkla, Thailand, 1987.

Position and site of the Employer's office.

Medical representative in Pharmaceutical Division,
Ciba-Geigy (Thailand) Limited Bangkok, Thailand.