

รายการเอกสารอ้างอิง

ภาษาไทย

- เต็ม สมิตินันท์. 2523. ข้อพิรภวไม้แห่งประเทศไทย (ข้อพุกเศษศาสตร์-ชื่อพืชเมือง). กรุงเทพมหานคร : พนีพับลิชชิ่ง 70
- ประทีป พุตระกูล. 2521. การหาปริมาณเซนโนไซด์ ในใบและผักมะขามแขกในประเทศไทย. กรุงเทพมหานคร : มหาวิทยาลัยศรีนครินทร์ วิโรฒประสาณมิตร
- ลัดดาวัลย์ บุญรัตนกรกิจ. 2536. สมุนไพรนำไปใช้. เล่ม 1 กรุงเทพมหานคร : แท่นทองบรรณาธิค์เชอร์วิส
- วัลย์ลด้า ทรงส์ทอง และ นฤมล รื่นราษฎร์. 2533. คู่มือสมุนไพรมะขามแขก. กรุงเทพมหานคร : สถาบันวิจัยวิทยาศาสตร์และเทคโนโลยีแห่งประเทศไทย.
- ศูนย์ข้อมูลสมุนไพร คณะเภสัชศาสตร์ มหาวิทยาลัยมหิดล. 2530. ก้าวไปกับสมุนไพร. เล่ม 1 กรุงเทพมหานคร : ชาร์มกมลการพิมพ์.
- . 2532. สมุนไพรนำไปใช้ยานมือ. กรุงเทพมหานคร
เศรษฐกิจการพาณิชย์ (กรม). 2533. รายงานผลการศึกษาโครงการวิจัย
ตลาดฟืชสมุนไพรและเครื่องเทศ. กรุงเทพมหานคร. 503-509
- ข้อมูล ล้วนรัตน์. 2534. ยาและผลิตภัณฑ์ที่แสดงฤทธิ์ต่อระบบทางเดินอาหาร.
ยาและผลิตภัณฑ์จากธรรมชาติ. กรุงเทพมหานคร : โรงพิมพ์
วุฒิลังกรณ์มหาวิทยาลัย. 295-296
- อาคารฯ ชาติเสถียร. 2534. การวิเคราะห์สารประกอบจากพืชสมุนไพร
มะขามแขก ดองดึง และชีลีบัน โดยใช้ HPLC. การประชุมวิชาการ
วิทยาศาสตร์และประเทศไทย ครั้งที่ 17. กรุงเทพมหานคร :
สมาคมวิทยาศาสตร์แห่งประเทศไทยในพระบรมราชูปถัมภ์. 356-357

ການຫຼັງກຸມ

- Aimon Somanabandhu et al. 1986. Specification of Thai Medicinal Plants. Bangkok : Aksornsampan Press.
- Alarm, N. and Gupta, P.C. 1986. Structure of a Water Soluble Polysaccharide from the Seed of *Cassia angustifolia*. Plant Biochem. 106:308-310
- Alfermann, A.W. and Reinhard, E. 1978. Production of Natural Compounds by Plant Cell Culture Method. p. 27. Munchen:GSF-BPT Report.
- Andrews, F.W. 1952 The Flowering Plants of the Anglo-Egyptian Sudan. 2. Scotland : T. Buncle and Co. Ltd.
- Asahina, Y. and Fujikawa, F. 1935 "Lichen Substance XLIV. Salazinic Acid and LV Endocrocin, a New Hydroxy-Derivatives" Ber. 68 : 1500-1503 (Though CA29 : 7308⁵)
- Atzorn, R., Weiler, E.W., and Zenk, M.H. 1981. Formation and Distribution of Sennosides in *Cassia angustifolia*, as Determined by a Sensitive and Specific Radioimmunoassay. Planta Medica. 41:1-14
- Baars, A.J., Vermeulen, R.J. and Breimer D.D. 1976. Gas Chromatographic determination of the laxative 1,8-dihydroxyanthraquinone in urine and faeces. J. Chromatogr. 120:217-220

- _____. 1974b. Quantitative determination in folia and fructus sennae. Planta Medica. 26 : 124-127
- Baier, S., and Friedrich, H. 1970. Anthracene derivatives in callus cultures of senna. Naturwissenschaften. 57(11):548-549
- Barz, W., Reinhard, E., and Zenk, M.H. 1977. Plant Tissue Culture and its Biotechnological Application. New York. Springer-Verlag.
- Bentham, G. 1871. Revision of the genus *Cassia*. Trans. Linn. Soc. 27:503-591
- Berlin, J. 1986. Secondary Products from plant cell cultures. Biotechnology. 4:630-658
- Bhatia, R.K., Lohar, D.R., Chawan, D.D., and Garg, S.P. 1978a. Chemical control of seedling growth and sennosides content in senna. Broteria, Ser. Cienc. Nat. 47(3-4):93-98 (Through CA 90. 1979 : 133959Z)
- _____. 1978b. Chemical control of yield of sennosides in senna leaves. Planta Medica. 34:437-441
- _____. 1978c. Effect of growth retardants on early seedling growth and sennoside contents in *Cassia angustifolia* Vahl. 1975. Biol. Conterm. 2(3):129-131
- Brenan, J.P.M. 1958. New and noteworthy *Cassias* from tropical Africa. Kew Bulletin (2)231-252

- Brendel, W.D. and Berger, L. 1974. Quantitative Determination in Senna Leaves and Fruits and their Preparations III Statistical Evaluation of the result. Planta Medica. 26(2):124-127
- British Pharmacopoeia. 1988. Cambridge, England: Her Majesty's Stationary Office, University Printing House-502-504
- Butcher, D.N., Ingram, D.S., 1976. Plant Tissue Culture 1th Edition. Great Britain : The Gametol Press Ltd.
- Caleffi, C; Ventura, M.T; Ciasca, M.A. 1970. New chemical method for the quantitative determination of anthracene derivatives in drugs. Farmaco. Ed. Prat. 25(1):47-58 (Through CA 72. 1970 : 93355C)
- Chaleff, R.S. 1983. Isolation of agonomically useful mutants from plant cell cultures. Science 219 : 676-682.
- Charlwood, V.B. and Rhodes, M.J.C. 1990. Physiology of secondary product formation in plants. Oxford : Clarendon Press.
- Chopra, R.N., Nayer, S.L., and Chopra, I.C. 1956. Glossary of Indian Medicinal Plants. New Delhi : Council of Scientific and Industrial Research.
- _____. 1969. Supplement to Glossary of Indian Medicinal Plants. New Delhi : Council of Scientific and Industrial Research.

- Clarke. E.G.C. Isolation and Identification of Drugs p.180
- Claus, E.P. 1949. Pharmacognosy. 2 nd ed. USA.
- Claus, P.S. 1992. Anthranoid Laxatives and Colorectal Cancer. TiPS. 13:229-231
- Daiya, K.S.; Sharma, H.K.; Chawan, D.D.; Sen, D.N. Effect of some growth retardants and their interactions with humic acid on the growth of *Cassia* species. Cienc. Nat. 48(1979) : 19-23 (Through CA 92. 1980 : 210116t)
- Davey, M.R. 1938. Recent development in the culture and regeneration of plant protoplast. Experientia Suppl. 46 : 19-20.
- Denee, R. and Huizing, H. 1981. Purification and Separation of Anthracene Derivatives on a Polystyrenedivinybenzene Copolymer. J. Nat. Prod. 44:257-240
- Dressen, M., Stragier, J. and Eyssen, H. 1982. The Metabolism of Sennosides by Intestinal Bacteria, Bio. Act. Nat. Prod. 111/40:246
- Duez, P., Vanhaelen, M., Hanacq, M., and Molle, L. 1984. Comparison between high-performance thin-layer chromatography-fluorometry and high-performance liquid chromatography for the determination of sennoside A and B in senna (*Cassia* spp.) pods and leaves. J. Chromatogr. 270:390-394

- Eilert, U., Kurz, W.G.W., and Constable, F. 1985. Stimulation of sanguinarine accumulation in *Papaver somniferum* cell cultures by fungal elicitors. J. Plant Physiol. 199 : 65-67.
- Erni, F. and Frei, R.W. 1976. A low-east gradient system for high-performance liquid chromatography, quantitation of complete pharmaceutical raw materials. J. Chromatogr. 125 : 265-274
- _____. 1978. Two dimensional column liquid chromatographic Technique for resolution of complex mixtures. J. Chromatogr. 149 : 561-569
- Fairbrain, J.W., and Shrestha, A.B. 1967. The taxonomic validity of *Cassia acutifolia* and *Cassia angustifolia*. Lloydia. 30(1):67-72
- Farnsworth, N.R. and Bunyapraphatsara, N. 1992. Thai Medicinal Plants. Bangkok : Prachachon Co., Ltd.
- Fernandes, Pruscilla; Kokate, C.K.; and Tipnis, H.P. 1982. Anthraquinones in tissue cultures of *Cassia angustifolia* Vahl. Indian Drugs. 19(11):442-445
(Through CA 98 : 1983, 14396U)
- Fowler, M.W. 1983. Commercial applications and economic aspects of mass plant cell culture. Plant Biotechnology. Cambridge : Cambridge University Press.
- Friedrich, H; and Baier, S. 1973a. Anthracene derivatives in callus cultures from *Cassia angustifolia*. Phytochem. 12(6):1459-1462

- Frischknecht, P.M., Baumann, T.W., and Tanner, H. 1977. Tissue culture of *Coffea arabica* : Growth and caffeine formation. Planta medica 31 : 344-350.
- Fujita, Y. and Tabata, M. 1986. Secondary metabolites from plant cells : Pharmaceutical application and progress in commercial product. Plant tissue and cell culture. Minnesota : University of Minneapolis Press.
- Fukui, H. et al. 1982. Production of isoquinoline alkaloids by cell suspension cultures of *Coptis japonica*. Plant tissue and cell culture. Tokyo : Maruzen Co.
- Furuya, F. et al. 1983. Saponin production in cell suspension cultures of *Panax ginseng*. Planta Medica. 48 : 83-87.
- Gaitonde, R.V., Bhat, U. 1987. TLC-Spectrophotometric determination of total sennoside from market galenical preparation. Phytochem. Lab. 3(1-4) : 137-5
- Gennaro, A.R. 1990. Reminton's Pharmaceutical Sciences. 18th Edition. Easton, Pennsylvania : The Mack Publishing Company.
- George, E. 1943. Senna leaflets-their palisade ratio values and ranges. Quart. J. Pharm. Pharmacological. 16:226-232
- Gorler, K., Mutter, S., and Westphal, C. 1979. HPLC of Senna Preparation. Planta Medica. 37:308-315

- Gritsanapan, W., 1983. Anthraquinone compounds of *Cassia* species growing in Thailand. Varasarn Presachasarathara. 10(3):90-96
- Gupta, G.K., Rao, P.R. and Vishwapaul. 1975. Production of sennoside concentrate from senna leaves and pods. Res. Ind. 20(3):124-128
- Gupta, R., Srivastava, V.K. and Maheshwari W.L. 1977. Sennoside contents in senna leaves and pods as influenced by growth and nutrition. Indian J. Pharm. 39(5):109-111
- Habit, A.M., and El-Sebakhy, N.A. 1980. Spectrophotometric estimation of sennosides and rhein glycosides in senna and its preparation. J. Nat. Prod. 43(4):452-458
- Harborne, J.B. 1982. Introduction of ecological biochemistry. 2nd Edition New York : Academic Press.
- Harry Swartz. M.d. 1981. Prescriber's Handbook of Therapeutic drugs. USA. 146-149
- Hiroshi, N., and Ushio, S. 1982. Formation of germichrynone by tissue cultures of *Cassia torosa* : induction of secondary metabolism in the lag phase. Phytochemistry. 21:319-323
- Hooker, J.D. 1973. The Flora of British India. Vol. VI. New Delhi : Bishen Singh Mahendra Pal Singh.

- Huang, K.C. 1993. The Pharmacology of Chinese Herb, Florida : CRC Press, Inc.
- International Trade Centre. UNCTAD/GATT. 1982. Market for Selected Medicinal Plants and Their Derivatives.
- James E.F. Reynolds, Martindale, 19 edition, London, 1989, p.1105-6
- Jech, J. 1974. Problem in the sennoside determination in leave and fruit of senna comparison of the DAB7 and Ph. Eur. I method Dtsch. Apoth-Ztg., 114(50): 1978-1979 (Through CA 82. 1975 : 116126t)
- Kalashnikova, G.K., Romanova, A.S., and Shevlinisk, A.N. 1983. Anthracene derivatives from the roots of *Cassia acutifolia*. Khim.-Farm. Zh. 19(5): 569-573 (Through CA 103. 1985 : 51212y)
- Kalyanasundaram, N., Patel, A.I., and Dalal, K.C. 1981. Effect of nitrogen and phosphorus on yield and quality of senna leaflet. Indian J. Pharm. Sci. 43(3):110-112
- Kaul, B. and Staba, E.J. 1968. *Dioscorea* tissue cultures. I. Biosynthesis and isolation of diosgenin from *Dioscorea deltoidea* callus and suspension cells. Lloydia 31 : 171-179
- Khafagy, Saad M.; Girgis, A.N.; Khayyal, S.E.; Helmi, M.A. Estimation of sennoside A, B, C and D in senna leaves, pods, and formulations. Planta Medica. 21 (1972):304-309

Kirtikar, K.B., and Basu, B.D. 1935. Indian Medicinal Plants. Vol. IV. 2nd Edition. Allahabad, India : Latit Mohan Basu.

Kisa, K., Sasak, K., Yamauchi, K., and Kuwano, S. 1981. Potentiating Effect of Sennoside C on Purgative Activity of Sennoside A in Mice. Planta Medica. 42:302-312

Kitanaka, S. et al. 1985. Component of laxative crude drug and Quantitative analysis of sennoside A and B in senna cultivar in Japan by thin-layer Chromatography-densitometry method. Coll. Sci. Technol. 39(2):106-110

Koblitz, H. et al. 1975. Gewebekulturen aus alkaloidpflanzen : IV. *Macleaya microcarpa* (Maxim.) Fedde. Experientia 31 : 768-769.

Krieger, Th. M., Verpoorte, R., Kreek, M., and Baerheim, A. 1984. Identification of Alkaloids and Anthraquinones in *Cinchona pubescens* Callus Cultures; the Effect of Plant Growth Regulators and Light on The Alkaloid Content. Planta Medica. 50:17-20

Krieger, Th.M., Verpoorte, P., Water, A., Gessel, M.V., Oeveren, B.C.J.A and Baerheim, A. 1982. Identification of the Alkaloids and Anthraquinones in *Cinchona ledgeriana* Callus Cultures. Planta Medica. 46:19-24

- Kumano, M., Handa, S., and Hirayama, F. 1988. Analysis of senna leaf components in tea with weight-reducing effect. Nagasaki-ken Eisei Kogai Kenkyushoho. 30: 129-132 (Through CA 111. 1989 : 95688C)
- Kyung, K., et al. 1988. Production of 20 metabolites by hairy roots and regenerated paltn transformed with Ri Plasmids. Chem. Pharm. Bull. 36(10):4217-4220
- Lemli, J., and Cuveele, J. 1972. Anthraquinone drug XXII detection of anthrones in purgative drug. Plant. Med. Phytochem. 6(4):271-274
- _____. 1974a. Anthraquinone drugs XXIII. Chromatography of anthrone glycoside purgative durg. Planta Medica. 26/2:193-5
- _____. 1974b. Chromatography of anthronglycosides in purgative drug. Planta Medica 26 : 192-193
- Lemli, J. Toppet, S., Cuveele, J., and Janssen, G. 1981. Napthalene Glycoside *Cassia senna* and *Cassia angustifolia*. Stuides in the field of drugs containing anthracene derivatives. Planta Medica. 43(1):11-17
- Lemli, J. et al. 1983. Chemical Identification of Alexandrian and Tinnevelly Senna. Planta Medica. 49:36-37
- Lemmens. 1977. Determination of Dihydroxydianthrones by Densitometry after Thin-Layer Chromatographic Separation. J. Chromatogr. 132(2):363-365

- Leistner, E. "Mode of Incoporation of Precursors into Alizarin (1,2-dihydroxy-9,10-anthraquinone) Phytochemistry. 12 (1973) : 337-345
- Leistner, E and Zenk, M.H. "Chrysophanol (1,8-dihydroxy-3-methyl-anthraquinone, Biosynthesis in Higher Plants" Chem. Commun. (1969):210-211
- Levin, D.A. 1976. The Chemical defences of plants to pathogen and microorganism Annual Review of Ecological System. 7:121-159
- Linneus, C. 1799. Species Plantarum Genus Cassia. 4 th Edition. Facsmile Ray Society.
- Lohar, D.R., Bhatia, R.K., and Garg, S.G. 1979. Seasonal variation in the content of sennoside senna leaves. Weekbl. Sci. Ed. 1(1):206-208 (Through CA 90. 1979 :18537j)
- Luckner, M. and Nover. L. 1977. Expression of secondary metabolism. An aspect of cell specialization of microorganisms, higher plants, and animals. Secondary metabolism and cell differentiation. Berlin : Springer-Verlag.
- Macek, K. 1972. Pharmaceutical Applications of Thin-Layer and Paper Chromatography. Amsterdam : Elsevier Publishing Co.
- Matsumoto, T. et al. 1981. Selection of cultured tobacco cell strains producing high levels of ubiquinone 10 by a cell cloning technique. Agric. Biol. Chem. 45 : 1627-1633.

- Mietheing, H., Boventer, W., and Haensel, R. 1986a. Comparison of the photometric sennoside content determination in the European Pharmacopoeia with a new specific densitometric method. Dtsch. Apoth. Ztg. 126:2158-2159 (Through CA 106. 1987:32531u)
- . 1986b. Densitometric content determination of aloë-emodin glycosides in senna pods and leaves and their aqueous preparations. Inst. Pharmakognosie Phytochem. 127(49):2587-2588
- . 1987. Sennosides in aqueous formulations, determination of release kinetics. Inst. Pharmakognosie Phytochem. 127(49):2587-2588
- Momonoki, T., and Momonoki, Y.S., 1987. Estimation of germinability of gramineous and leguminous seeds in long-term storage by mean of peroxidase activity and TTC reduction. JARQ. 20(4):296-301
- Morton, J.F. 1977. Major Medicinal Plant : Botany Culture and Uses. Springfield, Illinois : Charles C Thomas Publisher.
- Mueller, B.M., Kraus, J., and Franz, G. 1989. Chemical Structure and Bioogical activity of H_2O -solubles polysacc. from *Cassia angustifolia* leave Planta Medica. 55(6):536-539
- Muffat, F., Bernard, P., and Sabot, J.F. 1986. Determination of sennoside A and sennoside B in senna extract by HPLC J. Chromatogr. 369(1): 261-264

- Nakanishi, K., Goto, T., Ito, S., Natori, S. and Nozoe, S. 1974. Nat. Prod. Chem. Vol 2. Tokyo : Kodansha Scientific Ltd.
- Nandi, R.P., and Chatterjee, S.K. 1978. Physiological studies in some medicinal plants with special reference to synthesis of active principles in relation to photoperiodic and GA₃ treatment. Environ. Physiol. Ecol. Plants. 373-384 (Through CA 94. 1981:61801Z)
- Narong Chomchalow and Oradee Sahavacharin. 1981. The role of tissue culture in the development of medicinal plants and spices. Proceedings of the International Symposium Held at National University of Singapore, April 28-30 : 162-166.
- Nathalie, C., Denis, R., and Jean, A. 1990. Solid-phase extraction and fluorimetric detection of benzophenanthridine alkaloid from *Sanguinaria canadensis* cell culture. J. Chromatogr. 519:99-107
- Nishimura, H. and Hanba, K. 1977. Sennosides Calcium Salt from Senna. Japan Kokai. 77(12):916
- Noguchi, H., and Sangawa, U. 1982. Formation of germichrysone by tissue cultures of *Cassia torasa* : induction of secondary metabolism in the lag phase. Phytochemistry. 21(2):319-323
- Ohshima, Y., and Takahashi, K. 1983. Separation Method for Sennosides. J. Chromatogr. 258 : 292-296

- Pierre, D., Vanhaelen, M., Vanhaelen-Fastre, R., Hanoco, M. and Molle, L. 1984. Comparison between High-Performance Thin-Layer Chromatography Fluorometry and High Performance Liquid Chromatography for the Determination of Sennosides A and B in Senna (*Cassia* spp.) Pods and Leaves. J. Chromatogr. 303:391-395
- Razzaque, A. and Ellis, B. E. 1977. Rosmarinic acid production in *Coleus blumei*. Planta 137 : 287-292.
- Reinert, J. and Bajaj, Y.P.S. 1977. Applied and Fundamental Aspects of Plant Cell, Tissue and Organ Culture. p. 649. New York : Springer-Verlag
- Reynolds, J.E.F. 1989. Martindales, The Extra Pharmacopoeia 29. 29 th Edition. London : The Pharmaceutical Press.
- Rhodes, M.J.C. et al. 1987. Secondary product formation in plant cell cultures. Journal of Applied Bacteriology Symposium Supplement : 105S-114S.
- Robinson, T. 1967. In The Organic Constituents of Higher Plants 2nd ed., pp. 106-112 Mass, U.S.A.: Burgess Publishing Company, Menaplis.
- Romanova, A.S. et al. 1982. Method for the quantitative determination of anthracene derivative in senna leaves and fruits. Khim. Farm. Zh. 16(3):330-334 (Through CA 96. 1982:205476S)

- Rosental, G.A. and Janzen, D.H. 1979. *Herbivores : Their interaction with secondary metabolites.* London : Academic Press.
- Ross, M.S.F. and Brain, K.R. in An Introduction to Pharmacy, pp 166-175, Pitnam Press, Bath, Great Britain, 1977.
- Rowson, J.M. 1943. The significance of stomatal index as a differential character. Quart. J. Pharm. Pharmacol. 16:24-31
- Saad, M., Khafagy, Gergis, A.N., Khayyal, S.E. and Helmi, M.A. 1972. Estimation of Sennosides A, B, C and D in Senna Leaves, Pods and Formulations. Planta Medica. 21:304-309
- Samia, M.E., and Samir, A.R. 1982. A phytochemical study of some *Cassia* species cultivated in Egypt. J. Nat. Prod. 46(3):431-432
- Sasaki, K., Yamauchi, K., and Kuwano, S. 1979. Metabolic activation of Sennoside A in mice 37:370-378
- Sastri, B.N. 1950. The Wealth of India Raw Material. Vol III. Delhi. India Council of Scientific and Industrial Research.
- Sato, F. and Yamada, Y. 1984. High berberine producing cultures of *Coptis japonica* cells. Phytochemistry 23 :281-285.
- Shargool, P.D. 1982. Appl. Biochem. Biotech. 7:239-257

- Sharma, H.K., Daiya, K.S., and Chawan, D.D. 1982. Sennoside and carbohydrate in developing pod of *Cassia* species. Pharm. Weekbl. 4(3):83-84
- Sharma, O.P. and Khanna, P. 1980. Studies on steroid sapogenins from tissue cultures of *Agave wightii*. Lloydia 43 : 459-462.
- Shuler, M.L. 1981. Production of Secondary metabolite from plant tissue culture. Annul. N.Y. Acad. Sci. 369:65-79
- Smith, J.T. et al. 1987. Stimulation of indole alkaloid production in *Catharanthus roseus* (L.) G. Don by vanadyl sulfate. Plant Cell Report 6: 142-145.
- Srivastava, V.K., Maheshwari, M.L., and Mandal, S. 1983a. A Rapid HPLC Method for Analysis of Sennosides in Senna (*Cassia angustifolia* Vahl). Ind. J. Pharm. Sci. 73:230-232
- _____. 1983b. Investigation of chemical method of assay of sennosides in senna (*C. angustifolia* Vahl). Int. J. Trop. Agric. 1(3):231-238
- _____. 1981. Dynamics and localization of anthracene compounds during growth of seena plants, *Cassia angustifolia* Vahl. Natl. Bur. Plant Genet. Resour. 20(1-2), 75-81
- Staba, E.J. 1980. Plant tissue culture as a source of biochemicals Florida : CRC-Press.

- Stephen, C.Y.S., and Ferguson, N.M. 1973. Extraction and Separation of Anthraquinone Glycosides. J. Pharm. Sci. Vol 62. No 6 : 899-901
- Su, Stepphen C.Y., Ferguson, N.M. 1973. Extraction and separation of anthraquinone glucosides. J. Pharm. Sci. 62(6):899-901
- Swain, T. 1977. Secondary compounds as protective agents. Ann. Rev. Plant Phys. 28:479-501
- Tabata, M. and Noboru, H. 1976. Variation of alkaloid in *Nicotina rustica* callus cultures. Physiol. Plant. 38 : 19-23.
- Tabata, M. et al. 1975. The production of anthraquinones in callus cultures of *Cassia tora*. Lloydia 38 : 131-134.
- _____. 1978. Selection of cell lines with higher yield of secondary products. Frontiers of plant tissue culture. Canada : University of Calgary.
- Takahashi, Kunio, Kaizuka, Hitomi, Oshima and Yukio. 1983. Application of Three-Dimensional UV Aborbance-High -Performance Liquid Chromatographic Patterns for the Analysis of Plant Extracts. J. Chromatogr. 268(3):522-527
- Thai Pharmacopoeia. 1987. Vol 1. Bangkok: The Drug committee and the Food and Drug Administration of Thailand.

- Thomson, R.H. "Distribution and Biogenesis" in Naturally Occuring Quinones 2nd ed., pp 1-38, London : William Clowers and Sars, 1971
- Trease, C.E. and Evan, W.C. 1966. Textbook of Pharmacognosy. 9th Edition Baillier Tindall and Cassell.
- _____. 1978. Pharmacognosy. 11th Edition. London Baillier Tindall.
- _____. 1989. Trease and Evan's Pharmacognosy. 13 Edition. Great Britain : Alden Press.
- Trease, G.E. 1931. Notes on some Leguminous seeds. Pharm. J. 126:211-212
- Upadhyaya, S.K., and Singh, V. 1989. Phytochemical Identify of *C. angustifolia* and *C. acutifolia*. Natl. Acad. Sci. Lett. 12(6):181-182
- Vahl, M. 1790. Symbolae Botanicae. 1, 29
- Van Horne, K.C., et al., 1990. Sorbent Extraction Technology, 1st Edition.
- Vasil K.I. 1987. Cell culture in phytochemistry. Cell culture and somatic cell genetics of plants 4. San Diego : Academic Press, Inc.
- Wallis, T.E. 1948. Practical Pharmacognosy. London : J.A. Churchill. 136-137
- Walter, H.E., and Memory, P.F. 1976. Medical Botany, America, 1th Edition, 271-282

- Wanchai De-Eknamkul and Ellis, B.E. 1984. Rosmarinic acid production and growth characteristics of *Anchusa officinalis* cell suspension cultures. Planta Medica 51 : 346-350.
- Wijnsma, R., Vliet, T.B., Harkes, P.A.A., Groningen, H.J., Heijden, R., Verpootte, R., and Baerheim A. 1987. A Method for the Quantitative Determination of Anthraquinones and Alkaloids in Cell and Tissue Cultures of *Cinchona* Sp.. Planta Medica. 56:80-84
- Windholz, M., et al., 1983. The Merck Index an Encyclopedia of Chemicals, Drugs, and Biologicals. 10th Edition. USA : Merck & Co., Inc
- Wink, M. 1988. Plant breeding : Importance of plant secondary metabolites for production against pathogens and herbivores. Theoretical and Applied Genetics. 75:225-233
- Yasmin, A., Kapadia, Z., and Bader, Y. 1986. A comparative study of sennoside contents of *Cassia* species. Pak. J. Sci. Ind. Res. 20(5) : 348-349
(Through CA 106. 1987:192738y)
- Youngken, H.W. 1950. Textbook of Pharmacognosy. 6th Edition. New York : McGraw-Hill Book Company.
- Yukio, O., and Kunio, T. 1982. Separation method for sennosides. J. Chromatogr. 301:292-296

Zaka, S., Akhtar, M.W. and Khan, S.A. 1989. Lipd metabolism in germinating seeds of *Cassia*. Pak. J. Sci. Ind. Res. 32(5):323-332 (Through CA 108. 1988 : 147263v)

Zenk, M.H. et al. 1975. Anthraquinone production by cell suspension cultures of *Morinda citrifolia*. Planta Medica. Supplement. 75:79-81

_____. 1977. Formation of the indole alkaloids serpentine and ajmalicine in cell suspension cultures of *Catharanthus roseus*. Plant Tissue Culture and Its Biotechnological Application. Berlin : Springer-Verlag.

Zwaving, J.H., and Elema, E.T. 1976. A comparative investigation of two methods for the determination of 1,8-dihydroxyanthracene derivatives in vegetable drugs. Pharm. Weekbl. 1(52-53):1315-1320

VITA

Miss Supreeya Pomprasirt was born on September 23, 1965 in Phichit, Thailand. She received her Bachelor degree of Science of Pharmacy in 1988 from Chiang Mai University, Chiang Mai, Thailand. At present, she is the head of Herbal I section, Pharmaceutical IV Division of Production Department of The Government Pharmaceutical Organization of Thailand.

