



References

1. Fickett, J.W., and Hatzigeorgiou, A.G. Eukaryotic promoter recognition. Genome Res 7(9)(1997): 861-878.
2. Mahadevan, I., and Ghosh, I. Analysis of E.coli promoter structures using neural networks. Nucleic Acids Research 22(1994): 2158-2165.
3. Pedersen, A.G., Baldi, P., Brunak, S., and Chauvin, Y. Characterization of prokaryotic and eukaryotic promoters using hidden markov models. Proceedings of the Third International Conference on Intelligent Systems for Molecular Biology 1998.
4. Pedersen, A.G., and Engelbrecht, J. Investigations of Escherichia coli promoter sequences with artificial neural networks: new signals discovered upstream of the transcriptional startpoint. Proceedings of the Third International Conference on Intelligent Systems for Molecular Biology vol. 3, pp. 292-299, 1995.
5. Ma, Q., Wang, T.L., Shasha, D., and Wu, C.H. DNA Sequence Classification via an Expectation Maximization Algorithm and Neural Networks: A Case Study. IEEE Trans. Systems, Man and Cybernetics, Part-C: Applications and Reviews vol. 31, pp. 468-475, 2001.
6. Matsuda, T., Motoda, H., and Washio, T. Graph based induction and its applications. Advanced Engineering Informatics 16(2002): 135-143.
7. Matsuyama, Y., and Kawamura, R. Promoter Recognition for E.coli DNA Segments by Independent Component Analysis. Proceedings of the 2004 IEEE Computational Systems Bioinformatics Conference 2004.
8. Huang, Y.F., and Wang, C.M. Integration of Knowledge-Discovery and Artificial-Intelligence Approaches for Promoter Recognition in DNA Sequences. In Proceedings of the third conference on Information Technology and Applications 2005.
9. Bucher, P. Weight matrix descriptions of four eukaryotic RNA polymerase II promoter elements derived from 502 unrelated promoter sequences. Journal of Molecular Biology 212(4)(1990): 563-578.
10. Prestridge, D.S. Predicting Pol II promoter sequences using transcription factor binding sites. Journal of Molecular Biology 249(5)(1995): 923-932.
11. Hutchinson, G.B. The prediction of vertebrate promoter regions using differential hexamer frequency analysis. Comp Appl Biosc 12(5)(1996): 391-398.
12. Audio, S., and Claverie, J.-M. Detection of eukaryotic promoters using Markov transition matrices. Comput Chem 28(1997): 1997.
13. Chen, Q.K., Hertz, G.Z., and Stormo, G.D. A computer programme that predicts eukaryotic pol II promoters using strings and IMD matrices. Bioinformatics 28(1997): 39-35.

14. Solovyev, V., and Salamov, A. The Gene-Finder computer tools for analysis of human and model organisms genome sequences. Proceedings of International Conference Intelligent Systems for Molecular Biology vol. 8, pp. 294-302, 1997.
15. Reese, et al. Genome Annotation Assessment in *Drosophila melanogaster*. Genome Res 10(4)(2000): 483-501.
16. Scherf, M., Kingenhoff, A., and Werner, T. Highly specific localization of promoter regions in large genomic sequences by PromoterInspector: a novel context analysis approach. Journal of Molecular Biology 297(2000): 599-606.
17. Ohler, U., Niemann, H., Liao, G., and Rubin, G.M. Joint modeling of DNA sequence and physical properties to improve eukaryotic promoter recognition. Bioinformatics 17(2001): 199-206.
18. Knudsen, S. For the recognition of pol II promoter sequences. Bioinformatics 15(1997): 356-361.
19. Bajic, V.B., et al. Computer model for recognition of functional transcription start sites in RNA polymerase II promoters of vertebrates. Journal of Molecular Graphics and Modelling 21(2002): 323-332.
20. Howard, D., and Benson, K. Promoter Prediction with a GA-Automaton. LNCS 2611(2003): 44-53.
21. Chun-bo, and Toa, L. A hybrid neural network system for prediction and recognition of promoter regions in human genome. Journal of Zhejiang University SCIENCE 6B(5)(2005): 401-407.
22. Gangal, R., and sharma, P. Human pol II promoter prediction: time series descriptors and machine learning. Nucleic Acids Research 33(2005): 1332-1336.
23. Jeffrey, H.J. Chaos game representation of gene structure. Nucleic Acids Research 18(8)(1990): 2163-2170.
24. Goldman, N. Nucleotide, dinucleotide and trinucleotide frequencies explain patterns observed in chaos game representations of DNA sequences. Nucleic Acids Research, 21(10)(1993): 2487-2491.
25. Almeida, J.S., et al. Analysis of genomic sequences by Chaos Game Representation. Bioinformatics 17(5)(2001): 429-437
26. Deschavanne, P.J., et al. Genomic Signature: Characterization and Classification of Species Assessed by Chaos Game Representation of Sequences. Mol. Biol. Evol 16(10)(1999): 1391-1399.
27. Quinlan, J.R. Induction of decision trees. Machine Learning PP. 81-106, 1986.
28. Quinlan, J.R. C4.5: programs for machine learning. Morgan Kaufmann Los Altos, CA, 1993.
29. O'Neill, M.C. *Escherichia coli* promoters: neural networks develop distinct descriptions in learning to search for promoters of different spacing classes. Nucleic Acids Research 20(1992): 3471-3477.

30. Huang, -W.J., Yang, -B.C., and Tseng, -T.K. Promoter Prediction in DNA Sequences. Proc. of National Computer Symposium, Workshop on Algorithm and Computation Theory Taichung, Taiwan, 2003.
31. Kim, B.H., Park, -B.S., and Zhang, -T.B. PromSearch: A Hybrid Approach to Human Core-Promoter Prediction. LNCS 3177(2004): 125-131.

Biography

Name: Ms.Orawan Tinnungwattana.

Date of Birth: 11th August, 1974.

Educations:

- Ph.D., Program in Computer Science, Department of Mathematics, Chulalongkorn University, Thailand, (June 2002 - March 2006)
- M.Sc. Program in Information Technology, King Mongkut's Institute of Technology Ladkrabang, Thailand. (June 1996 - March 1998).
- B.Sc. Program in Applied Mathematics, King Mongkut's Institute of Technology Ladkrabang, Thailand. (June 1992 - March 1996).

Publication papers:

- O.Tinnungwattana and C.Lursinsap. Statistical Feature Selection From Chaos Game Representation for Promoter Recognition, *Proceedings of the 2nd International Conference on Bioinformatics*.