

REFERENCES

- Arai, S. 1990. What kind of magma could be equilibrated with ophiolitic peridotites? In J. Malpas, E. M. Moores, A. Pnayiotou, and C. Xenophontos (eds.), Ophiolites. Oceanic Crustal Analogues Proceedings of Symposium "Troodos 1987", pp. 555-565. The Geological Survey Department, Ministry of Agriculture and Natural Resources, Nicosia, Cyprus.
- Arai, S. 1992. Chemistry of chromian spinel in volcanic rocks as a potential guide to magma chemistry. Mineralogical Magazine 56: 173-184.
- Arai, S. 1994. Characterization of spinel peridotites by olivine-spinel compositional relationships: Review and Interpretation. Chemical Geology 113: 191-204.
- Arai, S., and Okada, H. 1991. Petrology of serpentinite sandstone as a key to tectonic development of serpentinite belts. Tectonophysics 195: 65-81.
- Asama, K. 1967. Permian plants from Maiya, Japan. Cathaysiopteris and Psygmorphylum. Bull. Nat. Sci. Museum. Tokyo 10: 139-153.
- Asama, K., and other 1985. Early Carboniferous plants from the Hikoroichi Formation, southern Kitakami massif, Northeast Japan. J. Geol. Soc. Japan 91: 421-426. (in Japanese)
- Bando, Y. 1970. Lower Triassic ammonoids from the Kitakami massif. Trans Proc. Palaeont. Soc. Japan, N. S. 79: 337-354.
- Bando, Y., and Shimoyama, S. 1974. Late Scythian ammonoids from the Kitakami massif. Trans. Proc. Palaeont. Soc. Japan, N. S. 94: 293-312.
- Batia, M. R. 1983. Plate tectonics and geochemical composition of sandstones. Geology Journal 91: 611-627.
- Bowie, S. H. U. 1967. Microscopy: Reflected light. In J. Zussman (ed.), Physical Methods in Determinative Mineralogy, pp. 103-159. New York: Academic Press.

- Cookenboo, H. O., Bustin, R. M., and Wilks, K. R. 1997. Detrital chromian spinel compositions used to reconstruct the tectonic setting of provenance: implications for orogeny in the Canadian Cordillera. Journal of Sedimentary Research 67: 116-123.
- Copper, P., Tazawa, J., and Mori, K. 1982. *Atrypa* (Devonian brachiopod) from Japan. Trans. Proc. Palaeont. Soc. Japan. N.S. 127: 368-374.
- Dick, H. J. B., and Bullen, T. 1984. Chromian spinel as a petrogenetic indicator in abyssal and alpine-type peridotites and spatially associated lavas. Contributions to Mineralogy and Petrology 86: 54-76.
- Dickinson, W. R., and others 1983. Provenance of North American Phanerozoic sandstones in relation to tectonic setting. Geological Society of American Bulletin 94: 222-235.
- Dozen, K., and Ishigi, H. 1997. Geochemical indication of provenance of Permian and Mesozoic shales of Southern Kitakami Terrane, Northeast Japan. Earth Science 51: 232-335.
- Ehiro, M., and Kanisawa, S. 1996. Formation and evolution of the South Kitakami Paleoland during the Paleozoic, with special reference to geodynamic of Eastern Asia (IGCP-321). Japan Contribution to the IGCP 43-50.
- Ehiro, M., and others 1986. Discovery of *Trimerella* (Silurian brachiopoda) from the Odagoe Formation south of Mt. Hayachine in the Kitakami Massif, Northeast Japan and its significance. J. Geol. Soc. Japan 92: 753-756. (in Japanese)
- Folk, R. L. 1974. Petrology of sedimentary rocks. Texas: Hemphills.
- Fujimaki, H., and Yomogida, K. 1986. Petrology of Hayachine ultramafic complex in contact aureole, NE Japan. (II) metamorphism and origin of the complex. J. Japan Assoc. Min. Petr. Econ. Geol. 81: 59-66.
- Haggerty, E. 1991. Oxide mineralogy of the Upper Mantle. In D. H. Linsley (ed.), Oxide Minerals: Petrology and Magnetic Significance, pp. 355-416. Michigan: BookCrafters, Inc.

- Harada, T. 1890. Die japanischen Inseln, eine topographisch-geologische Uebersicht. Berlin: Verlag von Paul Parcy. (in German)
- Hekinian, R. 1985. Petrology of the ocean floor. Elsevire Oceanography Series, no. 33. Amsterdam: Elsvire.
- Hisada, K., and Arai, S. 1993. Detrital chrom spinels in the Cretaceous Sanchu sandstone, central Japan: impicator of serpentinite protrusion into a fore-arc region. Palaeogeography, Palaeoclimatology, Palaeoecology 105: 95-109.
- Hisada, K., and Arai, S. 1994. Occurrence of detrital chromian spinels from the end-Permian sedimentary complex, Shikoku, and its implications for the tectonic of the Kurosegawa landmass. Ann. Rep., Inst. Geosci., Univ. Tsukuba 20: 23-27.
- Hisada, K., Arai, S., and Negoro, A. 1994. Devonian serpentinite protrusion onfirmed by detrital chromian spinels in outer zone of SW Japan. In Proceedings of the International Symposium on: Stratigraphic Conrrelation of Southeast Asia, pp. 76-80. 15-20 November, Bangkok, Thailand.
- Hisada, K., Arai, S., and Ishii, T. 1995. Occurrence of detrital chromian spinels and its implication: tectonic setting of Matsugadaira-Motai belt, NE Japan. Ann. Rep., Inst. Geosci., Univ. Tsukuba 21: 15-19.
- Hisada, K., Arai, S., Ishii, T., and Mogi, N. 1995. Gondwana Marginal Subduction Zone Deduced from Chemistry of Detrital Chromian Spinels from Hayachine Tectonic Belt, NE Japan. Journal of Geology, Series B, 5-6: 204-211.
- Hisada, K., and Arai, S. 1996. Occurrence and its tectonic significance of chromian spinels from the Kanmon Group, SW Japan. Ann. Rep., Inst. Geosci., Univ. Tsukuba 22: 31-36.
- Hisada, K., Arai, S., and Miyamoto, T. 1996. Detrital chromian spinels and Gondwana marginal subduction zone. Japan Contribution to the IGCP 35-41.
- Hisada, K., Mogi, N., and Arai, S. 1997. Detrital chromian spinels from the Tobigamori and Karaumedate Formations in Mt. Tobigamori area, South Kitakami Mountains. NE Japan. Ann. Rep., Inst. Geosci., Univ. Tsukuba 23.

- Igo, H. 1990. Paleozoic strata in the Hida "Gien" belt. In K. Ichikawa, S. Mizutani, I. Hara, S. Hada, and A. Yao (eds.), Pre-Cretaceous Terranes of Japan, pp. 41-48. Osaka: Publication of IGCP #224.
- Irvine, T. N. 1965. Chromian spinel as a petrogenetic indicator. Part 1. Theory. Canadian Journal of Earth Sciences 2: 648-672.
- Isozaki, Y. 1996. Anatomy and genesis of a subduction-related orogen: A new view of geotectonic subdivision and evolution of the Japanese Islands. The Island Arc 5: 289-320.
- Isozaki, Y., and Maruyama, S. 1991. Studies on orogeny based on plate tectonics in Japan and new geotectonic subdivision of the Japanese Islands. Journal of Geography 100: 679-761. (in Japanese with English abstract)
- Kadoshima, K., and Arai, S. 1999. Primary petrological characteristics of peridotites in the Sangun zone of northern Kyushu: a preliminary note from detrital chromian spinels. J. Min. Petr. Econ. Geol. 94: 97-108.
- Kato, M. 1979. Silurian Paleontology. In M. Minato, M. Hanahashi, J. Watanabe, and M. Kato (eds.), Variscan Geohistory of Northern Japan: the Abean Orogeny, pp. 60-61. Tokyo: Tokai University Press
- Kato, M. 1990. Paleozoic corals. In K. Ichikawa, S. Mizutani, I. Hara, S. Hada, and A. Yao (eds.), Pre-Cretaceous Terranes of Japan, pp. 307-312. Publication of IGCP #224, Osaka.
- Kato, M., and Minato, M. 1979. Silurian Introduction. In M. Minato, M. Hanahashi, J. Watanabe, and M. Kato (eds.), Variscan Geohistory of Northern Japan: the Abean Orogeny, pp. 54-56. Tokyo: Tokai University Press.
- Kawamura, M. 1985. Lithostratigraphy of the Carboniferous Formations in the Setamai region, Southern Kitakami Belt, Northeast Japan. (Part I). Shimoarisu district of the Setamai Subbelt. Jour. Geol. Soc. Japan 91, 3: 165-178 (in Japanese with English abstract)

- Kawamura, M. 1985. Lithostratigraphy of the Carboniferous Formations in the Setamai region, Southern Kitakami Belt, Northeast Japan. (Part II). Yokota district of the Setamai Subbelt. Jour. Geol. Soc. Japan 94, 4: 245-258. (in Japanese with English abstract)
- Kawamura, M. 1985. Lithostratigraphy of the Carboniferous Formations in the Setamai region, Southern Kitakami Belt, Northeast Japan. (Part III). Karosawa and Oide districts of the Omata Subbelt. Jour. Geol. Soc. Japan 91, 5: 341-352. (in Japanese with English abstract)
- Kawamura, M., Kato, M., and Kitakami Paleozoic Research Group 1990. Southern Kitakami Terrane. In K. Ichikawa, S. Mizutani, I. Hara, S. Hada, and A. Yao (eds.), Pre-Cretaceous Terranes of Japan, pp. 249-266. Osaka: Publication of IGCP #224.
- Kawamura, T. 1983. The Lower Carboniferous formations in the Hikoroichi region, southern Kitakami Mountains, northeast Japan. (Part I). Stratigraphy of the Hikoroichi Formation. Jour. Geol. Soc. Japan 89, 12: 707-722. (in Japanese with English abstract)
- Kawamura, T. 1984. The Lower Carboniferous formations in the Hikoroichi region, southern Kitakami Mountains, northeast Japan. (Part II). Sedimentological study of sandstone and limestone. Jour. Geol. Soc. Japan 90, 11: 831-847. (in Japanese with English abstract)
- Kawamura, T., and Kawamura, M. 1989. The Carboniferous system of the South Kitakami Terrane, Northeast Japan. (Part I). Summary of stratigraphy. Earth Science 43, 2: 84-97. (in Japanese with English abstract)
- Kawamura, T., and Kawamura, M. 1989. The Carboniferous System of the South Kitakami Terrane, Northeast Japan. (Part II). Sedimentary and tectonic environment. Earth Science 43, 3: 157-167. (in Japanese with English abstract)

- Kawamura, T., and others 1984. A new occurrence of Silurian fossils in the northern marginal part of the Southern Kitakami Belt. J. Geol. Soc. Japan 90: 61-64. (in Japanese)
- Kawamura, T., and others 1985. The Lower Carboniferous Odaira and Onimaru Formations in the Setamai-Yukisawa districts, southern Kitakami Mountains, Northeast Japan. J. Geol. Soc. Japan 91: 851-866. (in Japanese with English abstract)
- Kimura, T., Hayami, I., and Yoshida, S. 1991. Geology of Japan. University of Tokyo Press.
- Klein, C., and Hurlbut, C.S., Jr. 1993. Manual of Mineralogy. 21st ed. New York: John Wiley & Sons, Inc.
- Kobayashi, T., and Hamada, T. 1977. Devonian trilobites of Japan in comparison with Asian, Pacific and other faunas. Palaeont. Soc. Japan, Spec. Pap. 20.
- Kobayashi, T., and Hamada, T. 1987. A new Carboniferous trilobites from the Hida plateau, West Japan. Proc. Japan Acad. 63B: 115-118.
- Kobayashi, T., and Mori, K. 1954. Studies on the Jurassic trigonians in Japan. Part 2. *Prosogyrotrigonia* and the trigonea. Japan J. Geol. Geogra. 25: 155-175.
- Komatsu, M. 1990. Hida "Gaien" belt and Joetsu belt. In K. Ichikawa, S. Mizutani, I. Hara, S. Hada, and A. Yao (eds.), Pre-Cretaceous Terranes of Japan, pp. 25-40. Publication of IGCP #224, Osaka.
- Maruyama, S., Isozaki, Y., Kimura, G., and Terabayashi, M. 1997. Paleogeographic maps of the Japanese Islands: Plate tectonic synthesis from 750 Ma to the present. The Island Arc 6: 121-142.
- Maruyama, S., Liou, J. G., and Zhang, R. 1994. Tectonic evolution of the ultrahigh-pressure (UHP) and high-pressure (HP) metamorphic belts from central China. The Island Arc 3: 112-121.

- Minato, M. 1979. Introduction to the stratigraphy of the non-metamorphosed Paleozoic deposits of the Aben terrain. In M. Minato, M. Hanahashi, J. Watanabe, and M. Kato (eds.), Variscan Geohistory of Northern Japan: the Aben Orogeny, pp. 52-54. Tokyo: Tokai University Press.
- Minato, M., and Kato, M. 1977. Two spiriferida from the Hikoroichi formation (Tournaisian) in the Kitakami mountains, Japan. J. Fac. Sci. Hokkaido Univ. 4, 17: 613-617..
- Minato, M., and others 1953. Biostratigraphie des karbons in Kitakami-Gebirge, Nordostliches Honshu, Japan. J. Geol. Soc. Japan 59: 385-399.
- Mori, K., Okami, K., and Ehiro, M. 1992. Field Trip Guide Book 1. In M. Adatochi, and K. Suenki (eds.), 29th IGC: pp. 81-114. Nagoya Univ.
- Mori, K., and Tazawa, J. 1980. Discovery and significance of Visean rugose coral and brachiopods from the type locality of the Lower Carboniferous Hikoroichi Formation. J. Geol. Soc. Japan 86: 143-146. (in Japanese)
- Nanayama, F., Nakagawa, M., and Katch, T. 1993. Chemistry of detrital chromian spinel in latest Cretaceous to Paleocene sediments from eastern Hokkaido. Jour. Geol. Soc. Japan 99: 629-642.
- Nanayama, F., Nakagawa, M., and Okada, H. 1994. Chemistry of detrital chromian spinels from the Eocene Urahoro Group, in the eastern Hokkaido and its origin. Jour. Geol. Soc. Japan 100: 383-398.
- Nelles Verlag GmbH, D-80935 München. Explore the world. (Nelles Maps 1 : 5,000,000 of Japan with special maps: Tokyo environs, Osaka environs, Nagoya environ). Germany: Nelles.
- Noda, M., and Tachibana, K. 1959. Some Upper Devonian Cyrtospiriferids from the Nagasaki district, Kitakami mountainland. Sci. Bull. Fac. Lib. Art. Educ.. Nagasaki Univ. 10: 15-22.

- Ogasawara, K. 1988. Early Cretaceous fresh-water molluses from the Monomiyama Formation, Southern Kitakami mountains, Japan. Saito Ho-on Kai Pub. (Prof. T. Kotaka Comm. Vol.), 301-315.
- Qgawa, Y., Tamaki, K., Isozaki, Y., and Shikazono, N. 1997. Japan. In E. M. Moores, and R. W. Fairbridge (eds.), Encyclopedia of European and Asian Regional Geology, pp. 436-449. Lndon: Chapman & Hall.
- Okami, K., and others 1987. *Leptophloeum* bearing formation in the "Hayachine Tectonic Belt", Kitakami Massif, Northeast Japan. J. Geol. Soc. Japan 93: 321-327. (in Japanese with English abstract)
- Onuki, Y. 1937. New discovery of Gotlandian deposits from the Kesengun, Iwate Prefecture, Kitakami mountains, and Paleozoic stratigraphy (Preliminary report). J. Geol. Soc. Japan 44: 600-604.
- Onuki, Y. 1969. Geology of the Kitakami Massif, Northeast Japan. Contr. Inst. Geol. Pal. Tohoku Univ. 69: 1-239. (in Japanese)
- Ozawa, K. 1988. Ultramafic tectonic of the Miyamori ophiolite complex in the Kitakami mountains, Northeast Japan. Hydrous upper mantle in an island arc. Contribution of Mineralogy and petrology 99: 159-175.
- Pober, E., and Faupl, P. 1988. The chemistry of detrital chromian spinels and its implications for the geodynamic evolution of the Eastern Alps. Geologische Rundschau 77: 641-670.
- Press, S. 1986. Detrital spinels from Alpinotype source rocks in Middle Devonian sediments of the Rhenish Massif. Geologische Rundschau 75: 333-340.
- Roser, B. P., and Korsch, R. J. 1988. Provenance signatures of sandstone-mudstone suites determined using discriminat function analysis of major-element data. Chemical Geology 67: 119-139.
- Sack, R. O., and Ghiorso, M. S. 1991. Chromite as a petrogenetic indicator. In D. H. Linsley (ed.), Oxide Minerals: Petrologi and Magnetic Significance, pp. 355-416. Michigan: BookCrafters, Inc.

- Saito, Y., and Hashimoto, M. 1982. South Kitakami Region : An Allochthonous Terrane in Japan. Journal of Geophysical Research 87: 3691-3696.
- Sakagami, S. 1962. Lower Carboniferous Bryozoa from the Hikoroichi Series, Japan. Trans. Proc. Palaeont. Soc. Japan, N.S. 46: 227-242.
- Shikama, T., Kamei, T., and Murata, M. 1978. Early Triassic Ichthyosaurus, *Utatsusaurus hataii gen. et sp. nov.*, from the Kitakami massif, Northeast Japan. Sci. Rep. Tohoku Univ. 2, 48: 77-97.
- Sugiyama, T. 1944. On Japanese trilobites. Studies Geol. Min. Inst. Tokyo Bunrika Daigaku 1: 21-31. (in Japanese)
- Taira, A., Tokuyama, H., and Soh, W. 1989. Accretion Tectonics and Evolution of Japan. In Z. Ben-Avraham (ed.), The Evolution of Pacific Ocean Margins, pp. 100-123. Oxford: Oxford University Press.
- Takeuchi, M. 1994. Origin of detrital garnet, chromian spinel and chloritoid from the Lower Jurassic Shizugawa Group in the South Kitakami Belt. Journal of the Geological Society of Japan 100: 234-248.
- Tazawa, J. 1988. *Zdimir* (Devonian Brachiopoda) from the Kitakami Mountains, Northeast Japan and its paleobiogeographical significance. J. Geol. Soc. Japan 94: 1013-1016. (in Japanese)
- Tazawa, J. 1989. Comparison between the Paleozoic formations of the Southern Kitakami Mountains, NE Japan and the Hida Marginal Belt, SW Japan: a preliminary note. Earth Science 43: 224-230.
- Tazawa, J., 1993. Pre-Neogene tectonics of the Japanese Islands from the viewpoint of paleobiogeography. Jour. Geol. Soc. Japan 99: 525-543. (in Japanese with English abstract)
- Tazawa, J., and Kurita, H. 1986. Brachiopods and their correlation with the Lower Carboniferous Shittakazawa Formation, Southern Kitakami Mountains, Northeast Japan. J. Geol. Soc. Japan 92: 167-170. (in Japanese)

- Tazawa, J., and Osawa, M. 1979. Occurrence of *Martinia* sp. (Brachiopoda) from the Lower Carboniferous Karaumedate Formation, Southern Kitakami Mountains and its significance. J. Geol. Soc. Japan 85: 775-777. (in Japanese)
- Tazawa, J., and others 1979. Discovery of Early Cretaceous bivalves from the "Ubaishi Formation", Southern Kitakami mountains and their significance. J. Geol. Soc. Japan 85: 261-263. (in Japanese)
- Tucker, M. E. 1991.. Sedimentary Petrology: An Introduction to the Origin of Sedimentary Rocks. London: Blackwell Scientific Publications.
- Yoshida, K., Kawamura, M., and Kitakami Paleozoic Research Group. 1995. Detrital chromian spinels from the Silurian in Ohasama district, South Kitakami Terrane, Northeast Japan. Journal of the Geological Society of Japan 101: 817-820.
- Zhang, Z. M. 1985. Tectonostratigraphic terranes of Japan that bear on the tectonics of mainland Asia. In D. G. Howell (ed.), Tectonostratigraphic Terranes of the Circum-Pacific Region, pp. 409-420. Houston: The Circum-Pacific Council for Energy and Mineral Resources.
- Zimmerle, W. 1984. The Geotectonic Significance of Detrital Brown Spinel in Sediments. Mitt. Geol. Palaont. Inst. Univ. Hamburg 56: 337-360.

APPENDICES

APPENDIX I

List of rock samples from the Southern Kitakami area together with their investigation-methods.

Sample No.	Age	Roack name	Point counting	XRF	EPMA (spinel)
93120205	Carboniferous	Volcanic arenite	*		
93120206	Carboniferous	Volcanic arenite	*		*
93120207	Carboniferous	Siltstone			*
93120208	Carboniferous	Sandy siltstone			*
93120209	Carboniferous	Feldspathic litharenite	*		
97031501	Triassic	Plagioclase arenite			
97031502	Cretaceous	Plagioclase arenite			
97031503	Triassic	Plagioclase arenite		*	
97031601	Cretaceous	Plagioclase arenite	*		
97031603	Devonian	Feldspathic litharenite	*		
97031604	Devonian	Lithic arkose			*
97031605-1	Carboniferous	Feldspathic litharenite			
97031605-2	Carboniferous	Volcanic arenite			
97031605-3	Carboniferous	Feldspathic litharenite			*
97031605-4	Carboniferous	Sandy siltstone			
97031605-5	Carboniferous	Volcanic arenite	*		
97031606	Carboniferous	Plagioclase arenite	*		
97031607	Carboniferous	Siltstone			
97031611	Devonian	Feldspathic litharenite		*	
97031612	Devonian	Lithic arkose		*	
97031613	Devonian	Feldspathic litharenite		*	
97031614	Devonian	Feldspathic litharenite		*	
97031701	Permian	Plagioclase arenite	*		

Sample No.	Age	Roack name	Point counting	XRF	EPMA (spinel)
97031702	Permian	Plagioclase arenite	*		
97031703	Triassic	Plagioclase arenite	*		* (without EPMA)
97031704	Jurassic	Plagioclase arenite	*		
97031705	Jurassic	Plagioclase arenite	*		
97031706	Jurassic	Plagioclase arenite	*		
97031707	Jurassic	Plagioclase arenite			
97031708	Triassic	Plagioclase arenite	*		
97031709	Jurassic	Plagioclase arenite	*		
97031710	Jurassic	Plagioclase arenite	*		
97031711	Jurassic	Plagioclase arenite		*	
97031712	Jurassic	Plagioclase arenite		*	
97031713	Jurassic	Plagioclase arenite		*	
97031714	Jurassic	Plagioclase arenite		*	
97031715	Jurassic	Plagioclase arenite		*	
97031801	Permian	Calclithite			
97031803	Permian	Plagioclase arenite		*	
97031804	Permian	Plagioclase arenite		*	
97112101	Carboniferous	Volcanic arenite	*		
97112102	Carboniferous	Feldspathic litharenite			
97112103	Carboniferous	Volcanic arenite	*		
97112104	Carboniferous	Basalt			*
97112105	Carboniferous	Volcanic arenite	*		
97112106	Carboniferous	Volcanic arenite	*		
97112201	Carboniferous	Volcanic arenite			
97112202	Carboniferous	Basalt			
97112203	Carboniferous	Feldspathic litharenite	*		
97112204	Carboniferous	Lithic arkose	*		
97112205	Carboniferous	Feldspathic litharenite	*		

Sample No.	Age	Roack name	Point counting	XRF	EPMA (spinel)
97112206	Carboniferous	Feldspathic litharenite	*		
97112207	Carboniferous	Siltstone			
97112208	Carboniferous	Siltstone			
97112209	Carboniferous	Metamorphosed sandy siltstone			
97112210	Carboniferous	Feldspathic litharenite			
97112211	Carboniferous	Volcanic arenite	*		
97112212	Carboniferous	Volcanic arenite	*		
97112213	Carboniferous	Feldspathic litharenite			
97112214	Carboniferous	Feldspathic litharenite			
97112215	Carboniferous	Lithic arkose			
97112216	Carboniferous	Feldspathic litharenite			
97112217	Carboniferous	Calclithite			*
97112218	Carboniferous	Siltstone			
97112219	Carboniferous	Volcanic arenite	*		
97112301	Carboniferous	Feldspathic litharenite	*		
97112302	Carboniferous	Feldspathic litharenite	*		*
97112303	Carboniferous	Calclithite			
97112304	Carboniferous	Lithic arkose	*		
97112305	Carboniferous	Plagioclase arenite	*		
97112306	Carboniferous	Feldspathic litharenite			
97112307	Carboniferous	Calclithite			
97112308	Carboniferous	Calclithite			
97112309	Carboniferous	Lithic arkose			
97112310	Carboniferous	Lithic arkose			
97112311	Carboniferous	Lithic arkose			
97112312	Carboniferous	Andesite			
97112313	Carboniferous	Siltstone			

Sample No.	Age	Roack name	Point counting	XRF	EPMA (spinel)
97112314	Carboniferous	Andesite			
97112315	Carboniferous	Feldspathic litharenite	*		
97112316	Carboniferous	Sandy siltstone			*
97112317	Carboniferous	Feldspathic litharenite	*		
97112318	Carboniferous	Lithic arkose	*		

APPENDIX II

Major oxides of sandstones in the Southern Kitakami area analysed by XRF

Sample No.	Age	SiO ₂	TiO ₂	Al ₂ O ₃	Fe ₂ O ₃	CaO	MnO	MgO	K ₂ O	Na ₂ O	P ₂ O ₅	LOI	SUM
97031711	Jurassic	79.7	0.29	10.4	1.14	0.39	0.008	0.32	3.28	2.73	0.029	1.68	99.97
97031712	Jurassic	79.9	0.32	10.2	1.29	0.43	0.01	0.33	3.19	2.61	0.033	1.75	100.06
97031803	Permian	57.0	0.97	16.2	9.02	3.91	0.159	2.70	3.39	3.24	0.215	3.04	99.84
97031804	Permian	53.8	1.00	17.0	9.49	4.62	0.163	3.02	2.84	4.00	0.217	3.43	99.58
97031503	Triassic	60.2	0.90	18.3	7.34	1.19	0.052	2.53	1.89	3.22	0.188	5.07	100.88
97031713	Jurassic	71.2	0.20	13.0	1.32	3.86	0.018	0.57	2.83	3.45	0.036	4.26	100.74
97031714	Jurassic	70.6	0.21	13.3	1.32	3.66	0.019	0.58	2.82	3.52	0.038	4.18	100.25
97031715	Jurassic	69.5	0.38	12.9	1.81	4.44	0.027	0.62	2.36	3.70	0.052	4.57	100.36
97031611	Devonian	83.3	0.26	7.7	2.96	0.22	0.046	1.12	1.83	1.33	0.041	1.44	100.25
97031612	Devonian	70.1	0.40	12.9	6.06	0.82	0.113	2.33	2.18	2.75	0.076	2.49	100.22
97031613	Devonian	74.7	0.45	13.1	2.11	0.81	0.044	0.84	3.88	3.48	0.083	1.31	100.81
97031614	Devonian	64.6	0.54	15.4	7.42	0.94	0.135	2.85	2.82	2.71	0.096	3.04	100.55

APPENDIX III

Selected fossils from the Southern Kitakami area relating to the well-known area.

Age	Fossil	Relative Area
Silurian	Coral fauna: - <i>Nipponophyllum</i> - <i>Schedohalysites</i> Trilobite fauna: - <i>Enocrinurus kitakamiensis</i>	Eastern Australia New South Wales
Late Devonian	Plant: - <i>Leptophloeum rhombicum</i> - <i>Leptophloeum australe</i> - <i>Lepidodendron</i> - <i>Cyclostigma</i>	Southern China Australia and China
Early Carboniferous	Characteristic Corals: - <i>Pseudouralinia tangpakouensis</i> - <i>Kueichouphyllum heishihkuanense</i> - <i>Yuanophyllum kansuense</i> - <i>Spongophyllum</i> - <i>Australophyllum</i> Brachiopod fauna: <i>Kitakamithyris</i>	Southern China Bohemia and Australia Eastern Australia
	<i>Archaeocalamites</i> flora	Southern China
Late Carboniferous	Coral fauna (contain Boreal elements)	Tethys Character
Early Permian	Maiya flora: - <i>Cathaysiopteris</i> - <i>Noeggerathiopsis</i>	Cathysian (tropical area) Gondwana

BIOGRAPHY

The author was born in Chaiyaphum on May 1973. He got the diploma in Analytical Chemistry at the Institute of Analytical Chemistry Training, Department of Science Service, Ministry of Science, Technology, and Environment on March 1994. Then, he started the geology program at the Department of Geology, Faculty of Science, Chulalongkorn University, and got a Bachelor of Science there in March 1997. After that, he continued his Master program in geology at the Department of Geology, Faculty of Science, Chulalongkorn University.

