

รายการอ้างอิง

Cook, C. E. , Twine, M. E. , Myer, M. , Amenson, E. , Kepler, J. A., and Taylor, G. F. 1976. Theophylline radioimmunoassay: Synthesis of antigen and characterization of antiserum. Res. Commun. Chem. Pathol. Pharmacol. 13: 497-505.

Crowther, J. R. 1995. ELISA theory and practice. Institute for animal health, woking, UK: 177-205.

Daly, J. W. , Padgett, W. , Shamin, M. T. , Butts-Lamb , P. , and Water, J. 1985. 1,3-Dialkyl-8-(p-sulfonyl)xanthines: Potent water-soluble antagonists for A₁ and A₂-adenosine receptors. J. Med. Chem. 28: 87-492.

Ferencik Miroslav. 1993. Handbook of immunochemistry. 115: 142-143.

Habeeb, A. F. 1966. Determination of free amino groups in proteins by trinitrobenzene sulfuric acid. Anal. Biochem. 14: 328-336.

Hosada, H. 1979. The preparation of steroid N-hydroxysuccinimide esters and their reactivities with bovine serum albumin. Chem. Pharm. Bull. : 742-746.

Hosada, H. , Kawamura, N., and Nambara, T. 1981. Effect of bridge heterologous combination on sensitivity in enzyme immunoassay for cortisol. Chem. Pharm. Bull. 29(7): 1969-1974.

Hosada, H. , Kobayashi, N. , Ishii, N. , and Nambara, T. 1986. Bridging phenomena in steroid immunoassays. The effect of bridge length on sensitivity in enzyme immunoassay. Chem. Pharm. Bull. 34(5): 2105-2111.

Hosoda, H. , Kobayashi, N. , Ishii, N. , and Nambara, T. 1985. The bridge length effect on sensitivity in steroid enzyme immunoassay. Chem. Pharm. Bull. 33: 902-904.

Hosoda, H. , Kobayashi, N. , and Nambara, T. 1983. Effect of bridge heterologous combination on sensitivity in enzyme immunoassay for 11-deoxycortisol. Chem. Pharm. Bull. 31: 953-958.

Hosoda, H. , Sakai, Y. , Yoshida, H. , Miyairi, S. , Ishii, K. , and Nambara, T. 1979. The preparation of steroid *N*-hydroxysuccinimide esters and their reactivities with bovine serum albumin. Chem. Pharm. Bull. 27: 742-746.

Hosada, H. , Yashida, H. , Sakai, Y. , et al. 1980. Sensitivity and specificity in enzyme immunoassay of testosterone. Chem. Pharm. Bull. 28(10): 3035- 3040.

Hu, C. E. , Singh, P., and Ullman, E. F. 1980. Use of (Pivaloxy)methyl as a protecting group in the synthesis of antigenic theophylline (1,3-dimethyl xanthine) derivative. J. Org. Chem. 45: 1711-1713.

Ishikawa, E. , Kawai, T., and Miyai, K. 1981. Enzyme immunoassay. 133-134.

Kerr, M. A., and Thorpe, R. 1994. Immunochemistry. Lab Fax. Oxford. Blackwell scientific publication: 63-70.

Kim, H. O. , Ji, X. , Melman, N. , Olah , M. E. , Stiles, G. L., and Jacobson, K. A. 1994. Structure-activity relationships of 1, 4-dialkylxanthine derivatives at rat A₃ adenosine receptors. J. Med. Chem. 37: 3373-3382.

Kirton, K. T. et al. 1972. Characterization of antibody to prostaglandin F_{2α}. Biochem. Biophys. Res. Com. 47(4): 903-909.

Litman, et al. 1983. An internally referenced test strip immunoassay for morphine. Clin. chem. 29: 1589-1603.

McDonald, J. M., and Ladenson, J. H. 1978. Theophylline toxicity. Clin. Chem. 24(9): 1603-1608.

Nakane, P. K. , and Kawaoi, A. 1974. Peroxidase-Labeled antibody a new method of conjugation. J. Histo. Cytochem. 22: 1084-1091.

Nishikawa, T. , Saito, M. , and Kubo, H. 1984. Effect of heterologous combination on competitive nephelometric immunoassay I. site-heterologous combination for theophylline immunoassay. Chem. Pharm. Bull. 32: 4951-4957.

Okuyama, T., and Satake, K. 1960. On the preparation and properties of 2,4,6-trinitrophenyl-amino acids and peptides. J. Biochem. 47: 454-466.

Patrono, C., and Peskar, B. A. 1987. Producing of antisera by conventional techniques. Radioimmunoassay in basic and clinical pharmacology : 23-29.

Sadee, W., and Beelen, G. C. M. 1980. Drug level monitoring:Analytical technique metabolism and pharmacokinetics. USA: John Wiley & Son: 45.

Satake, K. , Kuyama, T. , Ohashi, M. , and Shinoda, T. 1960. The spectrophotometric determination of amine, amino acid and peptide with 2,4,6-trinitrobenzene 1-sulfonic acid. J. Biochem. 47: 645-660.

Sikora, K., and Smedley, H. M. 1984. Monoclonal antibodies : 6-8.

Standefer, J. C., and Saunders, G. C. 1978. Enzyme immunoassay for gentamicin Clin. chem. 24: 1903-1907.

Tijssen, P. 1985. Practical and theory of enzyme immunoassay. : Elsevier-North Holland: 279-283.

Van weemen, B. K., and Schuurs, A. H. 1972. Immunoassay using hapten enzyme conjugates. FEBS letters. 24(1) : 77-81.

Van weemen, B. K., and Schuurs, A. H. 1975. The influence of heterologous combinations of antiserum and enzyme labeled estrogen on the characteristics of estrogen enzyme-immunoassay. Immunochemistry. 12: 667-670.

Wilson, M. B., and Kakane, P. K. 1978. Recent developments in the periodate method of conjugating HRP to antibodies. Immunofluorescence and related staining techniques : 215-224.

Yoshida, H. , Sakai. , Y. , Miyairi, S. , and Nambara, T. 1980. Sensitivity and specificity in enzyme immunoassay of testosterone. Chem. Pharm. Bull. 28(10): 3035-3040.

Zuk, R. F. , et al. 1985. Enzyme immunochromatography-A quantitative immunoassay requiring no instrument. Clin. chem. 29: 1144-1150.

ประวัติผู้เขียน

นางสาวสุชาสินี พิชญ์วนิชิน เกิดเมื่อวันที่ 16 เมษายน พ.ศ. 2514 จบการศึกษาชั้นมัธยมศึกษาตอนปลายจากโรงเรียนนารีรัตน์จังหวัดแพร่ อำเภอเมือง จังหวัดแพร่ และสำเร็จการศึกษาชั้นปริญญาตรีเกียรตินิยมบัณฑิต จากคณะเภสัชศาสตร์ มหาวิทยาลัยเชียงใหม่ จังหวัดเชียงใหม่ เมื่อปีการศึกษา 2537

