REFERENCES

- [1] Tobin DJ: The biogenesis and growth of human hair, 2005.
- [2] Stough D, Stenn K, Haber R, Parsley WM, Vogel JE, Whiting DA, et al.: Psychological effect, pathophysiology, and management of androgenetic alopecia in men. Mayo Clinical Proceedings. Mayo Foundation, 2005. pp. 1316.
- [3] Ellis JA, Sinclair R, Harrap SB. Androgenetic alopecia: pathogenesis and potential for therapy. Expert Reviews in Molecular Medicine 2002; 2: 1-11.
- [4] Price VH: Androgenetic alopecia in women. Journal of Investigate Dermatology Symposium Proceedings. Nature Publishing Group, 2003. pp. 24-27.
- [5] Trüeb RM. Molecular mechanisms of androgenetic alopecia. Experimental Gerontology 2002; 37: 981-990.
- [6] Azzouni F, Godoy A, Li Y, Mohler J. The 5 alpha-reductase isozyme family: a review of basic biology and their role in human diseases. Advances in Urology 2011; 2012.
- [7] Harris WP, Mostaghel EA, Nelson PS, Montgomery B. Androgen deprivation therapy: progress in understanding mechanisms of resistance and optimizing androgen depletion. Nature Clinical Practice Urology 2009; 6: 76-85.
- [8] Kwack MH, Sung YK, Chung EJ, Im SU, Ahn JS, Kim MK, et al. Dihydrotestosteroneinducible dickkopf 1 from balding dermal papilla cells causes apoptosis in follicular keratinocytes. Journal of Investigative Dermatology 2007; 128: 262-269.
- [9] Meidan VM, Touitou E. Treatments for androgenetic alopecia and alopecia areata: current options and future prospects. Drugs 2001; 61: 53-69.
- [10] Takahashi T, Kamimura A, Hamazono-Matsuoka T, Honda S. Phosphatidic acid has a potential to promote hair growth in vitro and in vivo, and activates mitogenactivated protein kinase/extracellular signal-regulated kinase kinase in hair epithelial cells. Journal of Investigative Dermatology 2003; 121: 448-456.
- [11] Iino M, Ehama R, Nakazawa Y, Iwabuchi T, Ogo M, Tajima M, et al. Adenosine stimulates fibroblast growth factor-7 gene expression via adenosine A2b

receptor signaling in dermal papilla cells. Journal of Investigative Dermatology 2007; 127: 1318-1325.

- [12] Suraja R, Rejitha G, Anbu Jeba SB, Anandarajagopala K, Promwichita P. In vivo hair growth activity of *Prunus dulcis* seeds in rats. Biology & Medicine 2009; 1: 34-38.
- [13] Dinh QQ, Sinclair R. Female pattern hair loss: current treatment concepts.Journal of Clinical Intervention in Aging 2007; 2: 189.
- [14] Kumar T, Chaiyasut C, Rungseevijitprapa W, Suttajit M. Screening of steroid 5αreductase inhibitory activity and total phenolic content of Thai plants. Journal of Medicinal Plant Research 2011; 5: 1265-1271.
- [15] Pandit S, Chauhan NS, Dixit V. Effect of *Cuscuta reflexa* Roxb on androgeninduced alopecia. Journal of Cosmetic Dermatology 2008; 7: 199-204.
- [16] Raynaud J-P, Cousse H, Martin P-M. Inhibition of type 1 and type 2 5α-reductase activity by free fatty acids, active ingredients of Permixon[®]. Journal of Steroid Biochemistry & Molecular Biology 2002; 82: 233-239.
- [17] Liu J, Kurashiki K, Shimizu K, Kondo R. Structure–activity relationship for inhibition of 5α-reductase by triterpenoids isolated from *Ganoderma lucidum*. Bioorganic & Medicinal Chemistry 2006; 14: 8654-8660.
- [18] Murata K, Noguchi K, Kondo M, Onishi M, Watanabe N, Okamura K, et al. Inhibitory activities of *Puerariae Flos* against testosterone 5a-reductase and its hair growth promotion activities. Journal of Natural Medicine 2012; 66: 158-165.
- [19] Hiipakka RA, Zhang H-Z, Dai W, Dai Q, Liao S. Structure–activity relationships for inhibition of human 5α-reductases by polyphenols. Biochemical Pharmacology 2002; 63: 1165-1176.
- [20] Botchkarev VA, Paus R. Molecular biology of hair morphogenesis: development and cycling. Journal of Experimental Zoology Part: B Molecular Developmental Evolution 2003; 298: 164-180.
- [21] Tsuboi R. Growth factors and hair growth. Journal of Investigative Dermatology 1997; 4: 103-108.
- [22] Randall VA. Androgens and human hair growth. Clinical Endocrinology (Oxf) 1994;40: 439-457.

- [23] Messenger AG. The control of hair growth: an overview. Journal of Investigative Dermatology 1993; 101: 4S-9S.
- [24] Rho S-S, Park S-J, Hwang S-L, Lee M-H, Kim CD, Lee I-H, et al. The hair growth promoting effect of *Asiasari radix* extract and its molecular regulation. Journal of Dermatological Science 2005; 38: 89-97.
- [25] Alonso L, Fuchs E. The hair cycle. Journal of Cell Science 2006; 119: 391-393.
- [26] Huh S, Lee J, Jung E, Kim SC, Kang JI, Kim YW, et al. A cell-based system for screening hair growth-promoting agents. Archives of Dermatological Research 2009; 301: 381-385.
- [27] Randall VA. Androgens and hair growth. Dermatologic therapy 2008; 21: 314-328.
- [28] Cotsarelis G, Millar SE. Towards a molecular understanding of hair loss and its treatment. Trends in molecular medicine 2001; 7: 293-301.
- [29] Ahluwalia G: Cosmetics Applications of Laser & Light-Based Systems. William Andrew, 2008.
- [30] Paus R, Cotsarelis G. The biology of hair follicles. The New England Journal of Medicine 1999; 341: 491-497.
- [31] Kamimura A, Takahashi T. Procyanidin B-3, isolated from barley and identified as a hair-growth stimulant, has the potential to counteract inhibitory regulation by TGF- β 1. Experimental Dermatology 2002; 11: 532-541.
- [32] Itami S, Inui S: Role of androgen in mesenchymal epithelial interactions in human hair follicle. Journal of Investigative Dermatology Symposium Proceedings. 2005. pp. 209.
- [33] DeVillez RL, Jacobs JP, Szpunar CA, Warner ML. Androgenetic alopecia in the female: treatment with 2% topical minoxidil solution. Archives of dermatology 1994; 130: 303.
- [34] Matsuda H, Yamazaki M, Naruto S, Asanuma Y, Kubo M. Anti-androgenic and Hair Growth Promoting Activities of Lygodii Spora (Spore of Lygodium japonicum) Ι. Active Constituents Inhibiting Testosterone 5α-Reductase. Biological & Pharmaceutical Bulletin. 2002; 25: 622-626.

- [35] Han JH, Kwon OS, Chung JH, Cho KH, Eun HC, Kim KH. Effect of minoxidil on proliferation and apoptosis in dermal papilla cells of human hair follicle. Journal of Dermatological Science 2004; 34: 91-98.
- [36] Hamada K, Thornton MJ, Laing I, Messenger AG, Randall VA. The Metabolism of Testosterone by Dermal Papilla Cells Cultured From Human Pubic and Axillary Hair Follicles Concurs with Hair Growth in 5α-Reductase Deficiency. Journal of Investigative Dermatology 1996; 106: 1017-1022.
- [37] Battmann T, Bonfils A, Branche C, Humbert J, Goubet F, Teutsch G, et al. RU 58841, a new specific topical antiandrogen: a candidate of choice for the treatment of acne, androgenetic alopecia and hirsutism. The Journal of Steroid Biochemistry and Molecular Biology 1994; 48: 55-60.
- [38] Chen W, Orfanos C. The 5α-Reductase System and Its Inhibitors. Dermatology 1996; 193: 177-184.
- [39] Bayne E, Flanagan J, Einstein M, Ayala J, Chang B, Azzolina B, et al.
 Immunohistochemical localization of types 1 and 2 5*a*-reductase in human scalp. British Journal of Dermatology 1999; 141: 481-491.
- [40] Li Y-H, Yang Y-F, Li K, Jin L-L, Yang N-Y, Kong D-Y. 5 Alpha-Reductase and Aromatase Inhibitory Constituents from *Brassica rapa* L. Pollen. Chemical and Pharmaceutical Bulletin 2009; 57: 401-404.
- [41] Park WS, Son ED, Nam GW, Kim SH, Noh MS, Lee BG, et al. Torilin from *Torilis joponica*, as a new inhibitor of testosterone 5α-reductase. Planta medica 2003; 69: 459-461.
- [42] Seo E-K, Kim KH, Kim MK, Cho M-H, Choi EW, Kim KN, et al. Inhibitors of 5α reductase type I in LNCaP cells from the roots of *Angelica koreana*. Planta medica 2002; 68: 162-163.
- [43] Park W-s, Lee C-h, Lee B-g, Chang I-s. The extract of *Thujoe occidentalis* semen inhibited 5α -reductase and androchronogenetic alopecia of B6CBAF1/j hybrid mouse. Journal of Dermatological Science 2003; 31: 91-98.
- [44] Roh S-S, Kim CD, Lee M-H, Hwang S-L, Rang M-J, Yoon Y-K. The hair growth promoting effect of *Sophora flavescens* extract and its molecular regulation. Journal of Dermatological Science 2002; 30: 43-49.

- [45] Park SY, Kwack MH, Chung EJ, Im SU, Han IS, Kim MK, et al. Establishment of SV40T-transformed human dermal papilla cells and identification of dihydrotestosterone-regulated genes by cDNA microarray. Journal of Dermatological Science 2007; 47: 201-208.
- [46] Jones CD, Audia JE, Lawhorn DE, McQuaid LA, Neubauer BL, Pike AJ, et al. Nonsteroidal inhibitors of human type I steroid 5alpha-reductase. Journal of Medicinal Chemistry 1993; 36: 421-423.
- [47] Cha T-L, Qiu L, Chen C-T, Wen Y, Hung M-C. Emodin down-regulates androgen receptor and inhibits prostate cancer cell growth. Cancer Research 2005; 65: 2287-2295.
- [48] Pais P. Potency of a novel saw palmetto ethanol extract, SPET-085, for inhibition of 5α -reductase II. Advances in therapy 2010; 27: 555-563.
- [49] Cho C-H, Bae J-S, Kim Y-U. 5α-Reductase Inhibitory Components as Antiandrogens From Herbal Medicine. Journal of Acupuncture and Meridian Studies 2010; 3: 116-118.
- [50] Kumar N, Rungseevijitprapa W, Narkkhong N-A, Suttajit M, Chaiyasut C. 5αreductase inhibition and hair growth promotion of some Thai plants traditionally used for hair treatment. Journal of Ethnopharmacology 2012; 139: 765-771.
- [51] Hirata N, Tokunaga M, Naruto S, Iinuma M, Matsuda H. Testosterone 5α-reductase inhibitory active constituents of *Piper nigrum* leaf. Biological and Pharmaceutical Bulletin 2007; 30: 2402-2405.
- [52] Roh S-S, Park M-K, Kim Y-u. Abietic Acid from Resina Pini of Pinus Species as a Testosterone 5ALPHA-Reductase Inhibitor. Journal of Health Science 2010; 56: 451-455.
- [53] Lesuisse D, Berjonneau J, Ciot C, Devaux P, Doucet B, Gourvest J, et al. Determination of oenothein B as the active 5α-reductase-inhibiting principle of the folk medicine *Epilobium parviflorum*. Journal of Natural Products 1996; 59: 490-492.

- [54] Shimizu K, Fukuda M, Kondo R, Sakai K. The 5α-reductase inhibitory components from heartwood of Artocarpus incisus: structure-activity investigations. Planta medica 2000; 66: 16-19.
- [55] Shimizu K, Kondo R, Sakai K, Shoyama Y, Sato H, Ueno T. Steroid 5α-reductase inhibitory activity and hair regrowth effects of an extract from *Boehmeria nipononivea*. Bioscience, biotechnology, and biochemistry 2000; 64: 875-877.
- [56] Kim Y-u, Son HK, Song HK, Ahn M-J, Lee SS, Lee SK. Inhibition of 5*a*-reductase activity by diarylheptanoids from *Alpinia officinarum*. Planta medica 2003; 69: 72-74.
- [57] Ishiguro K, Oku H, Kato T. Testosterone 5*a*-reductase inhibitor bisnaphthoquinone derivative from *Impatiens balsamina*. Phytotherapy Research 2000; 14: 54-56.
- [58] Sawaya ME, Price VH. Different Levels of 5α -Reductase Type I and II, Aromatase, and Androgen Receptor in Hair Follicles of Women and Men with Androgenetic Alopecia. Journal of Investigative Dermatology 1997; 109: 296-300.
- [59] Zhu F, Chen X, Yuan Y, Huang M, Sun H, Xiang W. The chemical investigations of the mangrove plant Avicennia marina and its endophytes. Open Natural Products Journal 2009; 2: 24-32.
- [60] Ando Y, Yamaguchi Y, Hamada K, Yoshikawa K, Itami S. Expression of mRNA for androgen receptor, 5α -reductase and 17β -hydroxysteroid dehydrogenase in human dermal papilla cells. British Journal of Dermatology 1999; 141: 840-845.
- [61] Ito C, Katsuno S, Kondo Y, Tan H, Furukawa H. Chemical Consituents of Avicennia alba. Isolation and Structural Elucidation of New Naphthoquinones and Their Analogues. CHEMICAL AND PHARMACEUTICAL BULLETIN-TOKYO- 2000; 48: 339-343.
- [62] Rui J, Yuewei G, Huixin H. Studies on the chemical constituents from leaves of *Avicennia marina*. Chinese Journal of Natural Medicine 2004; 2: 16-19

VITA

Miss Ruchy Jain, 5377104033, was born on 12th October, 1989 in Bangkok, Thailand. She has graduated with the Bachelor degree in Engineering (Bionanoengineering) with 1st class honors from the International School of Engineering, Chulalongkorn University in 2010. She is currently pursuing her Doctor of Philosophy at the Department of Pharmaceutics and Industrial Pharmacy, majoring in International Graduate Program in Pharmaceutical Technology, Faculty of Pharmaceutical Sciences, Chulalongkorn University.