

## REFERENCES

- Abebe, W. and Macleod, K.M. 1990. Protein kinase C-mediated contractile responses of arteries from diabetic rats. Br.J.Pharmacol. 101:465-471.
- Ali, M. and Mohammad,S.V. 1986. Selective suppression of platelet thromboxane formation with sparing of vascular prostacyclin synthesis by aqueous extract of garlic in rabbit. Prostaglandins Leukot. Med. 25(2-3) : 139-146.
- Anchalee Jatapai. 1994, Effects of garlic extract on changes of serum lipid profiles and cardiac functions in induced diabetic rats. Master's Thesis, Physiology, Graduate School, Chulalongkorn University.
- Apitz-Castro, R., Ledezma, E., Escalante, J., and Jain, M.K. 1986. The molecular basis of the antiplatelet action of ajoene: direct interaction with the fibrinogen receptor. Biochem. Biophys.Res.Commun. 141(1):145-150.
- Aqel,M.B., Gharaibah, M.N., and Salhab, A.S. 1991. Direct relaxation effects of garlic juice on smooth and cardiac muscle. J.Ethnopharmacol. 33(1-2):13-19.
- Arbogast, B.W., Berry, D.L., and Newell,C.L. 1984. Injury of arterial endothelial cells in diabetic sucrose fed and aged rats. Atherosclerosis. 51:31-45.
- Barnett, A.H.1991. Pathogenesis of diabetic microangiopathy. Am. J.Med. 90 (suppl 6A) :675-735.
- Becker,K.L.1990. Principles and Practice of endocrinology and metabolism. Philadelphia: - J.B.Lippincott Co.
- Block, E. 1985. The chemistry of garlic and onions. Science Am. 252(3):94-99.
- Cameron, N.E.,and Cotter, M.A. 1992. Impaired contraction and relaxation in aorta from streptozotocin-diabetic rats: role of polyol pathway. Diabetologia. 35:1011-1019.
- Chang, M.W.,and Jonhson, M.A. 1980. Effect of garlic on carbohydrate metabolism and lipid synthesis in rats. J.Nutr.110:931-936.

- Das,I.,Khan,N.S., and Sooranna, S.R.1995. Nitric oxide synthase activation is a unique mechanism of garlic action. Biochem. Soc.Trans. 23(1):1365.
- Drury, P.L.1983. Diabetes and arterial hypertension. Diabetologia.24:1-9.
- Durante, W.,Sen,A.K.,and Sunahare, F.A. 1988. Impairment of endothelium dependent relaxation in aortae from spontaneously diabetic rats. Br.J.Pharmacol.94:463-468.
- Ernst, E. 1987. Cardiovascular effects of garlic (*Allium Sativum*): a review. Pharmatherapeutica. 5(2):83-89.
- Fenwick,G.R., and Hanley, A.B. 1985. The genus Allium.Part 3. (Reviews). Crit.Res.Food Sci.Nutr. 23(1):1-73 (abstract).
- Furchtgott, R.F., and Zawadzki, J.V.1980. The obligatory role of endothelial cells in the relaxation of arterial smooth muscle by acetylcholine. Nature Lond. 329:373-376.
- Gadkari, J.V.,and Joshi, V.D. 1991. Effect of ingestion of raw garlic on serum cholesterol level, clotting time and fibrinolytic activity in normal subject. J.Postgrad.Med. 37 (3):128-131.
- Greenspan, F.S.,and Baxter, J.D. 1994. Basic and clinical endocrinology. 4th ed. New York: Printice-Hall International Inc.
- Harrison, D.G.,and Ohara, Y.1995. Physiologic consequences of increased vascular oxidant stresses in hypercholesterolemia and atherosclerosis: implications for impaired vasomotion. Am.J.Cardiol. 75:758-818.
- Hsueh, W.A. and Anderson, P.W. 1992. Hypertension, the endothelial cell, and the vascular complications of diabetes mellitus. Hypertension. 20:253-263.
- Iberl, B.,Winkler, G., and Knoblock, K. 1990 Products of allicin transformation: ajoenes and dithiins characterization and their determination by HPLC. Planta Med. 56:202-211.
- Jackson, C.V., McGrath, G.M., Tahiliani, A.G., Vadlamudi, R.V.,and McNeill, J.H. 1985. A functional and ultrastructural analysis of experimental diabetic rat myocardium:manifestation of a cardiomyopathy. Diabetes. 34:876-883.

- Jain,R.C., and Vyas,C.R.1975. Garlic in alloxan-induced diabetes rabbits. Am.J.Clin.Nutr. 28:684-685.
- Jansen,H.,Muller,B.,and Knoblich, K.1987.Allicin characterization and its determination by HPLC. Planta Med. 6:559-562.
- Kahn, C.R.,and Weir, G.C.,1994. Joslin's diabetes mellitus. 13th ed. New York; Lea and Febiger a Warerly Co.
- Kaley,G., Rodenburg, J.M., Messina, E.J.,and Wolin, M.S. 1989. Endothelium-associated vasodilators in rat skeletal muscle microcirculation. Am.J.Physiol. 256 (Heat Circ. Physiol. 25) :H720-H725.
- Kaley, G., Koller, A., Rodenburg, J.M., Messian, E.J.,and Wolin, M.S.1992. Regulation of arteriolar tone and responses via L-arginine pathway in skeletal muscle. Am.J. Physiol. 262 (Heat Circ. Physiol.31):H987-H992.
- Kamata,K., Miyata, N.,and Kausya, Y.1989.Impairment of endothelium-dependent relaxation and changes in levels of cyclic GMP in aorta from streptozotocin-induced diabetic rats. Br. J.Pharmacol. 97:614-618.
- Kapen,C.W., Pritchard, K.A.,Arnold,J.H.,Cornwell,D.G.,and Panganamala, R.V. 1982. Restoration of prostacyclin/thromboxan A<sub>2</sub> balance in the diabetic rat. Diabetes. 31:947-951.
- Kiesewetter, H.,Jung,F.,Pinder,G.,Jung,E.M.,Mrowietz,C.,and Wenzel, E. 1991. Effect of garlic on thrombocyte aggregation, microcirculation, and other risk factors. Int.J.Clin.Pharmacol.Ther.Toxicol.29:151-155.
- Kimball, T.R.,Daniels, R.S.,Khoury, P.R., Magnotti, R.A., Turner, A.M.,and Dolan, L.M. 1994. Cardiovascular status in yong patients with insulin-dependent diabetes mellitus. Circulation. 90(1):367-361.

- Koller, A., Messina, E.J., Wolin, M.S., and Kaley, G. 1989a. Effects of endothelial impairment on arteriolar dilator responses in vivo. Am.J.Physiol. 257 (Heart Circ. Physiol 26): H1485-H1489.
- Koller, A., Messina, E.J., Wolin, M.S., and Kaley, G. 1989 b. Endothelial impairment inhibits prostaglandin and EDRF-mediated arteriolar dilation in vivo. Am.J.Physiol. 257 (Heart Cir. Physiol.26):H1966-H1970.
- Kunisaki, M., Fumio, U., Nawata, H., and King, G.L. 1996. Vitamin E normalizes diacylglycerol-protein kinase C activation induced by hyperglycemia in rat vascular tissues. Diabetes. 45 (suppl.3):S117-S119.
- Lee, T., Saltman, K.A., Ohashi, H., and King, G.L. 1989. Activation of protein kinase C by elevation of glucose concentration: proposal for a mechanism in the development of diabetic vascular complication. Proc.Natl.Acad.Sci USA. 56:5141-5145.
- Leech, C.J., and Faver, J.E. 1996. Difference alpha-adrenoceptor subtypes mediate constriction of arterioles and venules. Am.J.Physiol. 270(Heart Circ.Physiol.39) : H710-H722.
- Ledoux, S.P., Hall, C.R., Forbes, P.M., Patton, N.J., and Wilson, G.L. 1988. Mechanisms of nicotinamide and thymidine protection from alloxan and streptozotocin toxicity. Diabetes. 37:1015-1019.
- Martinez, R. 1992. Diabetic cardiomyopathy:possible pathogenetic role of coronary microcirculation (review). Minerve Cardioangiol. 40(1-2):1-5.
- Mathew, P.T., and Gugusti, K.T., 1973. Studies on the effect of allicin (diallyl disulphide-oxide) on alloxan diabetes:part I-hypoglycemic action and enhancement of serum insulin effect and glycogen synthesis. Indian J. Biochem.Biophys. 10:209-212.
- Mayhan, W.G., Faraci, F.M., and Heistad, D.D. 1987. Impairment of endothelium dependent responses of cerebral arterioles in chronic hypertension. Am.J.Physiol. 253 (Heart Circ. Physiol.22) :H1435-H1440.

- Mayhan, W.G., Simmons,L.K.,and Sharpe, G.M.1991. Mechanism of impaired responses of cerebral arterioles during diabetes mellitus. Am.J.Physiol. 260 (Heart Circ. Physiol.29):H319-H326.
- McDonagh, P.F.1983.Both protein and blood cells reduce coronary vascular permeability to macromolecular. Am.J.Physiol. (Heart Cir. Physiol.14):H698-H706.
- McDonagh, P.F.,Niven, A.T., and Roberts,D. 1984. Direct visualization of the coronary microcirculation for pharmacologic and physiologic studies. Microvasc. Res. 28:180-196.
- Meraji,S.,Jayakody,L.,Senaratne, M.P.J.,Thomson, A.B.R., and Kappagoda, T.1987. Endothelium-dependent relaxation in aorta of BB rat. Diabetes.36:978-981.
- Miller, T.B.1979.Cardiac performance of isolated perfused hearts from alloxan diabetic rats. Am.J.Physiol. 5(Heart Circ.Physiol.6) :H808-H812.
- Mirhadi,SA., Singh, S., and Gupta, P.P.1991. Effect of garlic supplementation to cholesterol-rich diet on development of atherosclerosis in rabbits. Indian J. Exp.Biol.,29(2):162-168.
- Nitenberg,A.,Valensi,P.,Sacha,R.,Dali,M.,Aptecar,E.,and Attale,J.1993. Impairment of coronary vascular reserve and Ach-induced coronary vasodilation in diabetic patients with angiographically normal coronary arteries and normal left ventricular systolic function: Diabetes. 42:1017-1025.
- Oyama, Y.,Kawasaki, H.,Hattor,Y.,and Kanno,M.1986. Attenuation of endothelium-dependent relaxation in aorta from diabetic rats. Eur. J. Pharmacol. 1986 ; 131 : 75-78.
- Ozturk,Y.,Aydin,S., Kosar, M.,and Baser,K.H. 1994. Endothelium-dependent and independent effects of garlic on rat aorta. J.Ethnopharmacol. 44:109-116.
- Patumraj,S.,Ritter, A.B.,and Duran, W.N.1992. A study of coronary macromolecular transport in diabetic rats .Chula Med.J.36(3):175-181.

- Penpargkul,S.,Schaible, T.,Yipintsoi, T., and Scheuer, J. 1980. The effect of diabetes on performance and metabolism of rat hearts. Circ. Res. 47: 911-921.
- Pentoja, C.V.,Chiang, L.C.H., Norris,B.C.,and Concha, J.B.1991. Diuretic, natriuretic and hypotensive effects produced by *Allium Sativum* (garlic) in anesthetized dogs. J. Ethnopharmacol. 31 : 325-331.
- Phelops.S.,and Harris, W.S. 1993. Garlic supplementation and lipoprotein oxidation susceptibility. Lipids.28(5) : 475-477.
- Poolsanong, N.1984. Pharmaceutical production of sugar-coated tablets and capsules of garlic, Production Division, Armed-Force Pharmaceutical Factory (unpublished).
- Rapoport, R.M.,and Murad, F.1983. Agonist -induced endothelium-dependent relaxation in rat thoracic aorta may be mediated through cGMP. Circ. Res. 52:352-357.
- Rashid,A.,and Khan, H.H. 1985. The mechanism of hypotensive effect of garlic extract. J.P.M.A.35:357-362.
- Rifkin, H.1991. Diabetes mellitus theory and practice . 4th ed. New York: Elsevier Co.
- Ross, R. 1986. The pathogenesis of atherosclerosis. New Eng.J.Med.20:448-500.
- Rossini, A.A.,Greiner,D.L.,Friedman,H.P., and Mordes,J.P.1993. Immunopathogenesis of diabetes mellitus. Diabetes. 1 (1):43-48.
- Saifon Saritdikul.1993. Effect of allicin on contraction of pregnant uterine segment in rats. Master's Thesis, Physiology, Graduate School, Chulalongkorn University.
- Sail, A.Y.,and Ahman, S.I. 1982. Study of the hypotensive action of garlic extract experimental animals. J.P.M.A. 32:237-239.
- Sheela, C.G.,and Augusti,K.T.1992. Antidiabetic effects of S-allyl cysteine sulphoxide isolated from garlic *Allium sativum* Linn. J.Exp.Biol. 30:523-526.
- Shimizu, K.,Muramatsu, M.,kakegawa, Y.,Asano, H., Toki,Y.,Miyazaki, Y.,Okumura, K., Hashimoto, H., and Ito, T. 1993. Role of prostaglandin H<sub>2</sub> as an endothelium-derived contracting factor in diabetic state. Deabetes. 42:1246-1252.

- Srivastava, K.C. 1986. Evidence for the mechanism by which garlic inhibits platelet aggregation. Prostaglandins Leukot.Mec. 22(3):313-321.
- Taylor, P.D. McCarthy, A.L., Thomas, C.R., and Poston, L. 1992. Endothelium-dependent relaxation and noradrenaline sensitivity in mesenteric resistance arteries of streptozotocin-induced diabetic rats. Br.J.Pharmacol. 107:393-399.
- Taylor, P.D. Wickenden, A.D., Mirrlees, D.J., and Poston, L. 1994. Endothelial function in the isolated perfused mesentery aortae of rats with streptozotocin-induced diabetes: effect of treatment with the aldose reductase inhibitor, ponalrestat. Br.J.Pharmacol. 111:42-48.
- Taylor, P.D., and Poston, L. 1994. The effect of hyperglycemia on function of rat isolated mesenteric resistance artery. Br.J.Pharmacol. 113:801-808.
- Tejada,S., Goldstein, I., Azadzoi, K., Krane, R. and Cohen, R.A. 1989. Impaired neurogenic and endothelium-dependent relaxation of human penile smooth muscle : the pathophysiological basis of impotence in diabetes mellitus. N Engl J Med. 320:1025-1030.
- Tesfamariam, B., Jakubowski, J.A., and Cohen, R.A. 1989. Contraction of diabetic rabbit aorta caused by endothelium-derived PGH<sub>2</sub>-Tx<sub>A</sub><sub>2</sub>. Am. J. Physiol. 257 (Heart Circ. Physiol. 26) H1327-H1333.
- Tesfamariam, B., Brown, M.L., Deykin, D., and Cohen, R.A. 1990. Elevated glucose promotes generation of endothelium-derived vasoconstrictor prostanoids in rabbit aorta. J. Clin. Invest. 85:929-932.
- Tesfamariam, B., Brown, M.L., and Cohen, R.A. 1991. Elevated glucose impairs endothelium-dependent relaxation by activating protein kinase C. J.Clin.Invest. 87:1643-1648.

- Tesfamariam, B., and Cohen, R.A. 1992 a. Superoxide anion and endothelium in vasoconstriction action of prostaglandin endoperoxide. Am.J.Physiol. 262 (Heart Circ. Physiol.31) :H1915-H1919.
- Tesfamariam, B., and Cohen,R.A. 1992b. Free radicals mediate endothelial cell dysfunction caused by elevated glucose. Am.J.Physiol. 263 (Heart Circ. Physiol.32) :H321-H326.
- Tesfamariam, B. 1994. Selective impairment of endothelium-dependent relaxations by prostaglandin endoperoxide. J.Hypertens. 12:41-47.
- Van, J.R., and Botting, R.M.1995. Pharmacodynamic profile of prostacyclin. Am.J.Cardiol. 75: 3A-10A.
- White, R.E., and Carrier, G.O.1988. Enhanced vascular alpha-adrenergic neuroeffector system in diabetes: importance of calcium. Am.J.Physiol. 255 (Heart Cir. Physiol.24):H1036-H1042.
- Williamson, J.R.,Ostrow,E.,Eabes,D.,Chang,K.,Alison, W., and Sherman, W.R. 1990. Glucose induced microvascular functional changes in nondiabetic rats are stereospecific and are prevented by and aldose reductase inhibitor. J.Clin.Invest. 80:1167-1172.
- Wilson, G.L.,Patton,N.J.,McDord, J.M.,Mullins,D.W.,and Mossman, B.T.1984. Mechanism of streptozotocin-and alloxan-induced damage in rat B cells. Diabetologia. 27 :587-591.
- Wilson,J.D., and Foster,D.W.1992. William textbook of endocrinology. 8th ed. New York: W.B.Saunders Co.
- Yeh,Y.Y., and Yeh, S.M. 1994. Garlic reduces plasma lipids by inhibiting hepatic cholesterol and triacylglycerol synthesis. Lipids. 23(3) :189-193.



APPENDIX

สถาบันวิทยบริการ  
จุฬาลงกรณ์มหาวิทยาลัย

## APPENDIX

### GAS CHROMATOGRAPH / MASS SPECTOMETER

Gas chromatography analysis is a method using for detection of evaporated sulfur compound in garlic oil. Garlic oil contained in closed vial is incubated in distilled water at  $60^{\circ}\text{C}$  for 30 min. Then, evaporation of garlic oil is injected into column of Gas Chromatograph / Mass Spectrometer (GC/MS) instrument. GC/MS is used to detect structural formula of organic compound that could evaporate. The data from this detection demonstrate as mass spectrum of organic compound. And, this spectrum could be shown as structural formula of detected organic compound.

The conditions used in GC/MS were :

Column

Carbovax 100/120 Mersh, 2 metres

FFAP 10 % on CW

Headspace sample control

sample temperature  $60^{\circ}\text{C}$

incubation time 30 min

Gas chromatograph

range x100

column temperature  $80^{\circ}\text{C}$

injection temperature  $150^{\circ}\text{C}$

Gas pressure

$\text{N}_2$  flow rate 30 ml/min

Sample injection

injection volume 0.5 ml




Fig.19 The result of mass spectrum indicated that garlic oil contained cis-trans propenyl methyl disulfide.

สถาบันวิทยบริการ  
 จุฬาลงกรณ์มหาวิทยาลัย




Fig.20 Structural formula of disulfide,methyl 1-propenyl that indicated by mass spectrum of garlic oil was shown.

สถาบันวิทยบริการ  
จุฬาลงกรณ์มหาวิทยาลัย

From this analysis by using GC/MS demonstrated that garlic oil consists of propenyl methyl disulfide as showed in Figure 19 and 20. Its structure is most likely ajoene. The structural formula of ajoene as showed in Figure 21 (Block, 1985). GC/MS is a method using for detection of evaporated sulfur compound in organic compound. Since the garlic oil is very sticky and could not inject into GC/MS. Moreover, the suitable solvent could not obtain for diluting this garlic oil, therefore, the analysis is performed using evaporation of oil. However, other compounds that have higher boiling temperature will not be able to detect by this technique.




Fig.21 Structural formula of ajoene ( Block , 1985 ).

## BIOGRAPHY

Miss Patcharin Sermpukdeekul was born on June 17, 1969, in Bangkok. She graduate Bachelor degree of Nursing from Mahidol University in 1991. After that, she was a nurse at gynecological unit, Siriraj hospital until 1994.



สถาบันวิทยบริการ  
จุฬาลงกรณ์มหาวิทยาลัย