

References

- Alford, R. A., Cohen, M. P., Crossland, M. R., Hearnden, M. N., James, D., and Schwarzkopf, L. 1995. Population biology of *Bufo masinus* in northern Australia. In Finlayson, M. (ed.). Wetland Research in the Wet-Dry Tropics of Australia. Supervising scientist report. pp. 173-181. Canberra: Office of the supervising scientist.
- Augert, D., and Joly, P. 1993. Plasticity of age at maturity between two neighbouring populations of the common frog (*Rana temporaria* L.). Canadian Journal of Zoology 71: 26-33.
- Barr, T. C. (Jr.). 1953. Notes on the occurrence of ranid frog in caves. Copeia 1953:60-61.
- Barreto, L., and Moreira, G. 1996. Seasonal variation in age structure and spatial distribution of a savanna larval anuran assemblage in Central Brazil. Journal of Herpetology 30(1): 87-92.
- Bee, M. A. 2002. Territorial male bullfrogs (*Rana catesbeiana*) do not assess fighting ability based on size-related variation in acoustic signals. Behavioral Ecology 13 (1): 109-124.
- Bee, M. A., and Gerhardt, H. C. 2001. Neighbour-stranger discrimination by territorial male bullfrogs (*Rana catesbeiana*): II. perspective basis. Animal behavior 62: 1141-1150.
- Bee, M. A., Korzich, C. E., Blackwell, K., and Gerhardt, H. C. 2001. Individual variation in advertisement calls of territorial male green frogs, *Rana clamitans*: Implications for Individual Discrimination. Ethology 107: 65-84.
- Beklemishev, V.N. 1960. "Spatial and Functional Structure of Populations". Bulletin MOIP section Biology 65(2): 41-50.
- Berry, P. Y. 1970. "The food of the Giant Toad *Bufo asper*". Zoological Journal of the Linnean Society. 49: 61-68.
- Berven, K. A. 1990. Factors affecting population fluctuations in larval and adult stages of the wood frog (*Rana sylvatica*). Ecology 71: 1599-1608.
- Bille, T. 2000. Microhabitat Utilization of the Mexican Salamander, *Pseudoeurycea leprosa*. Journal of Herpetology 34(4): 588-590.
- Blair, W. F. (ed.). 1972. Evolution in the genus Bufo. Austin: University of Texas Press. 460 pp.

- Blaustein, A. R., and Wake, D. B. 1990. Declining amphibian populations: global phenomenon? Trends Ecology Evolution 5: 203-204.
- Brattstrom, B. H. 1979. Amphibian temperature regulation studies in the field and in the laboratory. American Zoology 19: 345-356.
- Clarke, R. D. 1974. Postmetamorphic growth rates in a natural population of Fowler's toad, *Bufo woodhousei fowleri*. Canadian Journal of Zoology 52: 1489-1498.
- Cohen, M. P., and Alford, R. A. 1993. Growth, survival and activity patterns of recently metamorphosed *Bufo marinus*. Wildlife Research 20: 1-13.
- Crump, M. L. 2002. Amphibian Monitoring in Latin America: a protocol manual. Herpetological Review 33(3): 233-235.
- C-Tech International Co., Ltd. 2002. Chaloem Rattana Kosin National Park Management Project. Basic Information Report. Bangkok. (Mimeoographed)
- Duellman, W. E. 1999. Pattern of distribution of amphibians. Baltimore: The Johns Hopkins University Press. 633 pp.
- Duellman, W. E. and Trueb, L. 1999. Biology of amphibians. Baltimore: The Johns Hopkins University Press. 670 pp.
- Ehrlich, P. R., Roughgarden, J. 1987. The science of Ecology. New York: Macmillan Publishing Company. 710 pp.
- Finlayson, M. (ed.). 1995. Wetland Research in the Wet-Dry Tropics of Australia. Supervising scientist report. Canberra: Office of the supervising scientist.
- Frazer, J. F. D., 1967. A breeding colony of toads (*Bufo bufo* (L.)) in Kent. British Journal of Herpetology 3(2): 236-252.
- Freeland, W. J., and Kerin, S. H. 1988. Within-habitat relationships between invading *Bufo marinus* and Australian species of frog during the tropical dry season. Australian Wildlife Research 15: 293-305.
- Freeland, W. J., and Kerin, S. H. 1991. Ontogenetic alteration of activity and habitat selection by *Bufo marinus*. Wildlife Research 18: 431-443.
- Friedl, T. W. P., and Klump, G. M. 1997. Some aspects of population biology in the European treefrog, *Hyla arborea*. Herpetologica 53(3): 321-330.
- Green, D. M. 1992. Fowler's Toads (*Bufo woodhousei fowleri*) at Long Point, Ontario: changing abundance and implications for conservation. Canadian Wildlife Service Occasional Paper 76: 37-43.
- Green, D. M. 1997. Temporal variation in abundance and age structure in Fowler's toad, *Bufo fowleri*, at Long Point, Ontario. Herpetological Conservation 1: 45-56.

- Griffin P. C., and Case, T. J. 2001. Terrestrial habitat preferences of adult arroyo southwestern toads. Journal o f Wildlife Management 65(4): 633-644.
- Heyer, W. R., Donnelly, M. A., McDiarmid, R. W., Hayek, L. A. C., and Foster, M. S. (ed.). 1994. Measuring and monitoring biological diversity: standard methods for amphibians. Washington: Smithsonian Institution Press. 364 pp.
- Heying, H. E. 2001. Social and reproductive behaviour in the Madagascan poison frog, Mantella laevigata, with comparison to the dendrobids. Animal behaviour 61: 567-577.
- Houck, L. D., Arnold, S. J., and Thisted, R. A. 1985. A statistical study of mate choice: Sexual selection in a plethodontid salamander (Desmognathus ochripheaeus). Evolution 39: 370-386. Cited in Howard, R. D., Whiteman, H. H., and Schueller, T. I. 1994. Sexual selection in American toads: a test of a good-genes hypothesis. Evolution 48(4): 1286-1300.
- Houlahan, J. E., Findlay, C. S., Meyer, A. H., Kuzmins, S. L., and Schmidt, B. R. 2001. Brief communication. Nature 412: 500.
- Howard, R. D., Whiteman, H. H., and Schueller, T. I. 1994. Sexual selection in American toads: a test of a good-genes hypothesis. Evolution 48(4): 1286-1300.
- Inger, R. F. 1966. "The Systematics and Zoogeograaphy of the amphibia of Borneo". Fieldiana: Zoology. 52: 402 pp.
- IUCN. 2002. Global Amphibian Assessment: Species Asia Summarize. Southeast Asia workshop. XX pp. Sep. 30 to Oct. 4, 2002. Bangkok. (Mimeographed)
- Joglar, R. L. 1998. Los Coquies de Puerto Rico: Su historia natural y conservation. Universidad de Puerto Rico, San Juan. 232 pp. Cited in Prather, J. W., and Briggler, J. T. 2001. Use of small caves by anurans during a drough period in the Arkansa Ozarks. Journal of Herpetology 35(4): 675-678.
- Kellner A., and Green, D. M. 1995. Age structure and age at maturity in Fowler's toads, Bufo woodhousii fowleri, at their northern range limit. Journal of Herpetologica 29 (3): 485-489.
- Krebs, C. J. 1989. Ecological Methodology. New York: Harper & Row. 654 pp.
- Kusano, T., Maruyama, K., and Kaneko, S. 1995. Post-breeding dispersal of the Japanese Toad, Bufo japonicus formosus. Journal of Herpetology. 29(4): 633-638.

- Langhelle, A., Lindell, M. J., and Nystrom, P. 1999. Effects of ultraviolet radiation on amphibian embryonic and larval development. Journal of Herpetology 33(3): 449-456.
- Lehtinen, R. M. 2002. A historical study of the distribution of Blanchard's Cricket Frog (*Acris crepitans blanchardi*) in Southeastern Michigan. Herpetological Review 33 (3): 194-197.
- Licht, L. E. 1974. Survival of embryos, tadpoles, and adults of the frog *Rana aurota aurota* and *Rana pretiosa pretiosa* sympatric in southwestern British Columbia. Canadian Journal of Zoology 52: 613-627.
- Lowe, J. 2001. Population structure and habitat relationships of sympatric populations of the terrestrial salamanders *Plethodon elongatus* and *Ensatina escholtzii* a prescribed fire. Master's Thesis, Department of Biology, Faculty of Humboldt State University. 77 pp.
- Magnusson, W. E., Lima, A. P., Hero, J. M., and Araujo, M. C. 1999. The rise and fall of a population of *Hyla boans*: reproduction in a neotropical gladiator frog. Journal of Herpetology 33(4): 647-656.
- Marsh, D. M, Rand, A. S., and Ryan, M. J. 2000. Effects of inter-pond distance on the breeding ecology of tungara frogs. Oecologia 122: 505-513.
- Matsui, M. 1984. Morphometric variation analyses and revision of the Japanese Toads (Genus *Bufo*, Bufonidae). Contributions from the Biological Laboratory, Kyoto University 26(3): 209-428.
- Matthews, K. R., and Pope, K. L. 1999. A telemetric study of the movement patterns and habitat use of *Rana muscosa*, the mountain yellow-legged frog, in a high-elevation basin in Kings Canyon National Park, California. Journal of Herpetology 33(4): 615-624.
- Mazerolle, M. J. 2001. Amphibian activity, movement patterns, and body size in fragmented peat bogs. Journal of Herpetology 35(1): 13-20.
- Miaud, C., Sanuy, D., and Avrillier, J. N. 2000. Terrestrial movements of natterjack toad *Bufo calamita* (Amphibia, Anura) in a semi-arid, agricultural landscape. Amphibia-Reptilia 21: 357-369.
- Monello, R. J., and Wright, R. G. 1999. Amphibian habitat preferences among artificial ponds in the Palouse region of northern Idaho. Journal of Herpetology 33(2): 1999.

- Moreira, G., and Lima, A. P. 1991. Seasonal patterns of juvenile recruitment and reproduction in four species of leaf litter frogs in central Amazonia. Herpetologica 47: 295-300.
- Navas, C. A. 1996. The effect of temperature on the vocal activity of tropical anurans: a comparison of high and low-elevation species. Journal of Herpetology 30(4): 488-497.
- Navas, C. A., and Bevier, C. R. 2001. Thermal dependency of calling performance in the eurythermic frog *Colostethus subpunctatus*. Herpetologica 57(3): 384-395.
- Patto, C. E. G., and Pie, M. R. 2001. Notes on the population dynamics of *Hyloides asper* in Southeastern Brazil (Anura: Leptodactylidae). Journal of Herpetology. 35(2): 684-686.
- Pechmann, J. H. K., and Wilbur, H. M., 1994. Putting declining amphibian population in perspective: Natural fluctuations and human impacts. Herpetologica 50: 65-84.
- Pough, F. H., and Kamel, S. 1984. Post-metamorphic change in activity metabolism of anurans in relation to life history. Oecologia 65: 138-144. Cited in Cohen, M. P., and Alford, R. A. 1993. Growth, survival and activity patterns of recently metamorphosed *Bufo marinus*. Wildlife Research 20: 1-13.
- Prather, J. W., and Briggler, J. T. 2001. Use of small caves by anurans during a drough period in the Arkansa Ozarks. Journal of Herpetology 35(4): 675-678.
- Reading, C. J., Loman, J., and Madsen, T. 1991. Breeding pond fidelity in the common toad, *Bufo bufo*. The Zoological Society of London 225: 201-211.
- Resetarits, W. J. (Jr), and Aldridge, R. D. 1988. Reproductive biology of a cave-associated population of the frog *Rana palustris*. Canadian Journal of Zoology 66:329-333. Cited in Prather, J. W., and Briggler, J. T. 2001. Use of small caves by anurans during a drough period in the Arkansa Ozarks. Journal of Herpetology 35(4): 675-678.
- Schlupp I., and Podloucky, R. 1994. Changes in breeding site fidelity: a combined study of conservation and behavior in the common toad *Bufo bufo*. Biological Conservation 69: 285-291.
- Seebacher, F. and Alford, R. A. 1999. Movement and microhabitat use of a terrestrial amphibian (*Bufo marinus*) on a tropical island: seasonal variation and environmental correlates. Journal of Herpetology. 33 (2): 208-214.
- Semlitsch, R. D. 1981. Terrestrial activity and summer home range of the mole salamander (*Ambystoma talpoideum*). Canadian Journal of Zoology 59: 315-322.

- Shepard, D. B. 2002. Spatial relationships of male green frogs (*Rana clamitans*) throughout the activity season. American Midland Naturalist. 148: 394-400.
- Spieler, M., and Linsenmair K. E. 1998. Migration patterns and diurnal use of shelter in a ranid frog of a West African savannah: A telemetric study. Amphibia-reptilia 19 (1): 43-64.
- Squire, T., and Newman, R. A. 2002. Fine-scale population structure in the wood frog (*Rana sylvatica*) in a northern woodland. Herpetologica 58(1): 119-130.
- Taylor, E. H. 1962. "The amphibian fauna of Thailand". The University of Kansas Science Bulletin. 43(8): 265-599.
- Thornton, W. A., 1961. Population dynamics in *Bufo woodhousei* and *Bufo valliceps*. The Texas Journal of Science 12 (3/4): 176-200.
- Wake, D. B., 1994. Foreword. In Heyer, W. R., Donnelly, M. A., McDiarmid, R. W., Hayek, L. A. C., and Foster, M. S. (ed.). Measuring and monitoring biological diversity: standard methods for amphibians. pp xv-xvi. Washington: Smithsonian Institution Press.

ศูนย์วิทยทรัพยากร
จุฬาลงกรณ์มหาวิทยาลัย



Appendices

ศูนย์วิทยทรัพยากร
จุฬาลงกรณ์มหาวิทยาลัย

Appendix A

Area Characteristics inside Tarn Lord Noi Cave

ศูนย์วิทยทรัพยากร
จุฬาลงกรณ์มหาวิทยาลัย




Figure A-1 The nine sections inside Tarn Lord Noi Cave (Modified from C-Tech International Co, Ltd., 2002).

Table A-1 The area characteristic inside Tarn Lord Noi Cave, Chaloem Rattana Kosin National Park, Kanchanaburi Province during July 2001 to June 2002.

Section	Distance from downstream exit (m)	Stream appearance	Dominant Sediment
1	0-30	+	Rock and gravel
2	30-60	+	Rock and gravel
3	60-90	+	Rock and gravel
4	90-120	+	Sand and gravel
5	120-150	+	Sand
6	150-180	+	Rock
7	180-210	--	Rockfall
8	210-240	+	Rockfall
9	240-270	+	Rockfall



Figure A-2 Section 1 (0-30 m from the down stream exit).




Figure A-3 Section 2 (30-60 m from the down stream exit).



Figure A-4 Section 3 (60-90 m from the down stream exit).



Figure A-5 Section 4 (90-120 m from the down stream exit).




Figure A-6 Section 6 (150-180 m from the down stream exit).



Figure A-7 Section 7 (180-210 m from the down stream exit).



Figure A-8 Section 8 (210-240 m from the down stream exit).





Figure A-9 Section 9 (240-270 m from the down stream exit).





Appendix B

Mark and Recapture Data

ศูนย์วิทยบรังษักษ์
จุฬาลงกรณ์มหาวิทยาลัย

Table B-1 Mark-recapture data for a series of 11 samples of a total population of *Bufo asper* inhabiting Tarn Lord Noi Cave.

Time of last capture	Time of capture										
	Jul-01	Aug-01	Sep-01	Nov-01	Dec-01	Jan-02	Feb-02	Mar-02	Apr-02	May-02	Jun-02
Jul-01	59	1	0	0	0	1	0	0	0	0	0
Aug-01	49	4	8	0	1	0	0	0	1	1	0
Sep-01		18	12	2	1	1	2	0	0	0	0
Nov-01		15	2	2	2	1	1	0	0	0	0
Dec-01		23	24	3	3	1	2	0	0	0	0
Jan-02			19	5	1	0	1	0	1	1	1
Feb-02				42	4	4	1	1	0	0	0
Mar-02					34	7	3	3	22	6	6
Apr-02											18
May-02											
Total marked	0	59	50	22	35	27	48	53	43	33	28
Total unmarked	76	19	9	6	26	2	6	3	3	11	6
Total caught	76	78	59	28	61	29	54	56	46	44	34
Total released	76	78	58	28	61	29	54	56	46	44	33

Remark Fifteen surveys were carried out each month except in April 2002, 13 surveys were done.

Table B-2 Mark-recapture data for a series of 11 samples of a male population of *Bufo asper* inhabiting Tarn Lord Noi Cave.

Time of last capture	Time of capture										
	Jul-01	Aug-01	Sep-01	Nov-01	Dec-01	Jan-02	Feb-02	Mar-02	Apr-02	May-02	Jun-02
Jul-01	33	0	0	0	0	0	0	0	0	0	0
Aug-01		24	2	5	0	1	0	0	0	0	0
Sep-01			13	7	2	0	1	0	0	0	0
Nov-01				12	1	0	0	1	0	0	0
Dec-01					21	17	3	0	1	0	0
Jan-02						17	5	0	0	0	0
Feb-02							29	3	1	0	0
Mar-02								24	4	2	2
Apr-02									15	2	2
May-02										13	
Total marked	0	33	24	15	24	24	35	38	28	21	17
Total unmarked	38	4	1	3	21	0	2	1	0	0	0
Total caught	38	37	25	18	45	24	37	39	28	21	17
Total released	38	37	25	18	45	24	37	39	28	21	17

Remark Fifteen surveys were carried out each month except in April 2002, 13 surveys were done.

Table B-3 Mark-recapture data for a series of 11 samples of a female population of *Bufo asper* inhabiting Tarn Lord Noi Cave.

Time of last capture	Time of capture										
	Jul-01	Aug-01	Sep-01	Nov-01	Dec-01	Jan-02	Feb-02	Mar-02	Apr-02	May-02	Jun-02
Jul-01	11	0	0	0	0	0	0	0	0	0	0
Aug-01		12	1	1	0	0	0	0	0	1	0
Sep-01			3	3	0	0	0	1	0	0	0
Nov-01				2	1	2	1	0	0	0	0
Dec-01					0	3	0	1	1	1	0
Jan-02						1	0	0	0	0	0
Feb-02							8	0	0	0	0
Mar-02								6	0	1	
Apr-02									5	1	4
May-02											
Total marked	0	11	12	4	6	1	6	9	8	7	6
Total unmarked	14	8	5	2	1	0	2	1	2	11	4
Total caught	14	19	17	6	7	1	8	10	10	18	10
Total released	14	19	16	6	7	1	8	10	10	18	9

Remark Fifteen surveys were carried out each month except in April 2002, 13 surveys were done.

Table B-4 Mark-recapture data for a series of 11 samples of a young population of *Bufo asper* inhabiting Tarn Lord Noi Cave.

Time of last capture	Time of capture										
	Jul-01	Aug-01	Sep-01	Nov-01	Dec-01	Jan-02	Feb-02	Mar-02	Apr-02	May-02	Jun-02
Jul-01	15	1	0	0	1	0	0	0	0	0	0
Aug-01		13	1	1	0	0	0	0	0	0	0
Sep-01			1	1	0	1	0	1	0	0	0
Nov-01				1	0	0	1	0	0	0	0
Dec-01					2	4	0	0	0	0	0
Jan-02						1	0	1	0	1	1
Feb-02							4	1	0	0	0
Mar-02								4	3	0	0
Apr-02									2	3	3
May-02										1	1
Total marked	0	15	14	2	3	2	7	5	7	5	5
Total unmarked	24	7	3	2	6	2	2	1	0	0	2
Total caught	24	22	17	4	9	4	9	7	8	5	7
Total released	24	22	17	4	9	4	9	7	8	5	7

Remark Fifteen surveys were carried out each month except in April 2002, 13 surveys were done.

Biography

Miss Tassanee Eamkamon was born on the 18th of September 1976 in Chaiyaphume Province. She graduated her bachelor's degree of Science in Biology in 1998 from the Department of Biology, Faculty of Science, Chulalongkorn University. She continued her graduated study for a master's degree of science in zoology at the same University in 1999.



ศูนย์วิทยบริพยากรณ์
จุฬาลงกรณ์มหาวิทยาลัย