

รายการอ้างอิง

ภาษาไทย

- นิรุชา มงคลแสงสุรีย์, ศิริมาศ สุขประเสริฐ, อัจฉราภรณ์ เปี่ยมสมบุรณ์, ณิชฐารัตน์ ปภาวสิทธิ์ และวราพร ธารางกูร. “การเปลี่ยนแปลงองค์ประกอบของแพลงก์ตอนพืชในป่าชายเลนบ้านคลองโคน จังหวัดสมุทรสงคราม” บรรยาย ณ อาคารสารนิเทศ 50 ปี มหาวิทยาลัยเกษตรศาสตร์, 20 มีนาคม 2546. (เอกสารไม่ตีพิมพ์เผยแพร่)
- บัณฑิต สัจฉนทกสมิต. 2545. การแปรผันในรอบปีของประชากร COPEPOD, CLADOCERA และ ROTIFER ในป่าชายเลนบ้านคลองโคน จังหวัดสมุทรสงคราม. วิทยานิพนธ์ปริญญาโท มหาวิทยาลัย ภาควิชาวิทยาศาสตร์ทางทะเล บัณฑิตวิทยาลัย จุฬาลงกรณ์มหาวิทยาลัย. (กำลังดำเนินการวิจัย)
- อัจฉราภรณ์ เปี่ยมสมบุรณ์, ณิชฐารัตน์ ปภาวสิทธิ์, อิชฌมิกา พรหมทอง และวราพร ธารางกูร. 2545. ผลของการเปลี่ยนแปลงองค์ประกอบขนาดของแพลงก์ตอนพืชต่อการถ่ายทอดพลังงานในระบบนิเวศชายฝั่ง. ในประมวลงานวิจัยการประชุมวิชาการทรัพยากรและสิ่งแวดล้อมทางน้ำ เรื่องการจัดการและการใช้ประโยชน์อย่างบูรณาการ, หน้า I 81-I 89. 6-8 ธันวาคม 2544 ณ โรงแรมโลตัส ปางสวนแก้ว จังหวัดเชียงใหม่.
- อัมพร อึ้งปกรณ์แก้ว. 2540. เครื่องวิเคราะห์คาร์บอน ไฮโดรเจน ไนโตรเจน ซัลเฟอร์และออกซิเจน. วารสารศูนย์เครื่องมือวิจัยวิทยาศาสตร์และเทคโนโลยี 6(1): 51-71.
- อิชฌมิกา พรหมทอง. 2542. พลวัตและความหลากหลายของแพลงก์ตอนพืชในบริเวณปากแม่น้ำท่าจีน จังหวัดสมุทรสาคร. วิทยานิพนธ์ปริญญาโท มหาวิทยาลัย ภาควิชาวิทยาศาสตร์ทางทะเล บัณฑิตวิทยาลัย จุฬาลงกรณ์มหาวิทยาลัย.
- อุดมนิยมวิทยา, กรม. 2546. บันทึกข้อมูลทางอุดมนิยมวิทยารายเดือนของจังหวัดสมุทรสงคราม ตั้งแต่ปี พ.ศ. 2544-2545. ฝ่ายกรรมวิธีข้อมูล กรมอุดมนิยมวิทยา กระทรวงคมนาคม.

ภาษาอังกฤษ

- Agawin, N. S. R., Duarte, C. M. and Agusti, S. 2000. Nutrient and temperature control of the contribution of picoplankton to phytoplankton biomass and production. Limnol. Oceanogr. 45(3): 591-600.
- Allredge, A. L. and Madin, L. P. 1982. Pelagic tunicates: Unique herbivores in the marine plankton. BioScience. 32(8): 655-662.

- Almeida, M. A., Cunha, M. A. and Alcantara, F. 2001. Physiological responses of marine and brackish water bacterial assemblages in a tidal estuary (Ria de Aveiro, Portugal). Aquat. Microb. Ecol. 25 (2): 113-125.
- Alongi, D. M. (ed.), 1998. Coastal Ecosystem Processes. Florida: CRC Press.
- Andersen, R. A., Saunders, G. W., Paskind, M. P. and Sexton, J. P. 1993. Ultrastructure and 18S rRNA gene sequence for *Pelagomonas calceolata* gen. et sp. nov. and the description of a new class, the pelagophyceae classis nov. J. Phycol. 29: 701-715.
- Arar, E. J. and Collins, G. B. 1992. Method 445.0: In vitro determination of chlorophyll *a* and pheophytin *a* in marine and freshwater phytoplankton by epifluorescence. In USEPA Methods for the Determination of Chemical Substances in Marine and Estuarine Environmental Samples. Ohio, CA: U. S. Environmental Protection Agency.
- Berman, T., Azov, Y. and Townsend, D. 1984. Understanding oligotrophic oceans: Can eastern Mediterranean be a useful model?, pp. 101-112. In O. Holm-Hansen, L. Bolis and R. Gilles (eds.), Marine phytoplankton and productivity: lecture notes on coastal and estuarine studies vol. 8. Berlin: Springer-Verlage. Cited in Stockner, J. G. and Antia, N. J. Algal picoplankton from marine and freshwater ecosystems: A multidisciplinary perspective. Can. J. Fish. Aquat. Sci. 43: 2472-2503, 1986.
- Black, J. G. 1999. Characteristics of prokaryotic and eukaryotic cells. In Microbiology Principles and Explorations, pp. 71-75. New Jersey: Prentice-Hall.
- Brand, L. E., Sunda, W. G. and Guillard, R. R. L. 1986. Reduction of marine phytoplankton reproduction rates by copper and cadmium. J. Exp. Mar. Biol. Ecol. 96: 225-250. Cited in Vaquer, A., Troussellier, M., Courties, C. and Bibent, B. Standing stock and dynamics of picophytoplankton in the Thau Lagoon (northwest Mediterranean coast). Limnol. Oceanogr. 41(8):1821-1828, 1996.
- Butcher, R. W. 1967. An Introduction Account of the smaller algae of British coastal waters. Fishery investigation series IV. Part IV: Cryptophyceae. London: Her Majesty's Stationery Office. 54 pp.
- Calbet, A. and Landry, M. R. 1999. Mesozooplankton influences on the microbial food web: Direct and indirect trophic interactions in the oligotrophic open ocean. Limnol. Oceanogr. 44(6): 1370-1380.

- Campbell, L. and Vaultot, D. 1993. Photosynthetic picoplankton community structure in the subtropical North Pacific Ocean near Hawaii (Station Aloha). Deep-Sea Res. 40: 2043-2060.
- Caron, D. A., Peele, E. R., Lim, E. L. and Dennett, M. R. 1999. Picoplankton and nanoplankton and their trophic coupling in surface waters of the Sargasso sea south of Bermuda. Limnol. Oceanogr. 44(2): 259-272.
- Charpy, L. and Blanchot, J. 1998. Photosynthetic picoplankton in French Polynesian atoll lagoons: estimation of taxa contribution to biomass and production by flow cytometry. Mar. Eco. Prog. Ser. 162: 57-70.
- Cherrier, J., Bauer, J. E. and Druffel E. R. M. 1996. Utilization and turnover of labile dissolved organic matter by bacterial heterotrophs in eastern North Pacific surface waters. Mar. Eco. Prog. Ser. 139: 267-279.
- Chisholm, S. W., Olson, R. J., Zettler, E. R., Goericke, R., Waterbury, J. B. and Welschmeyer, N. A. 1988. A novel free-living prochlorophyte abundant in the oceanic euphotic zone. Nature 334: 340-343.
- Choi, K., Dobbs, F. C. and Cowen, R. K. 2001. Short-term temporal and spatial dynamics of bacterioplankton near Barbados in the Caribbean Sea. Aquat. Microb. Ecol. 25(1): 43-53.
- Clarke, K. R. and Gorley, R. N. 2001. PRIMER v5. [Computer program]. Plymouth Marine Laboratory: PRIMER-E Ltd.
- Conover, R. J. 1966. Feeding on large particles by *Calanus hyperboreus* (Kroyer). In H. Barnes (ed.), Some Contemporary Study in Marine Science. London: Allen & Unwin, pp. 187-194. Cited in D. L. Kirchman (ed.), Microbial ecology of the ocean, pp. 121-152. New York: WILEY-LISS, 2000.
- Daley, R. J. 1979. Direct epifluorescence enumeration of native aquatic bacteria : Uses, limitations, and comparative accuracy. In J. W. Costerton and R. R. Colwell (eds.), Native aquatic bacteria: enumeration, activity, and ecology , pp. 29-45. Philadelphia: American Society for Testing and Materials. Cited in Kirchman, D., Sigda, J. Kapuscinski, R. and Mitchell, R. Statistical analysis of the direct count method for enumerating bacteria. Appl. Environ. Microbiol. 44(3): 376-382, 1982.
- Day, J. W. Jr., Hall, C. A. S., Kemp, W. M. and Yanez-Arancibia, A. 1989. Estuarine Ecology. New York: Wiley & Son.

- Dehairs, F., Rao, R. G., Mohan, P. C., Raman, A. V., Marguillier, S. and Hellings, L. 2000. Tracing mangrove carbon in suspended matter and aquatic fauna of the Gautami-Godavari Delta, Bay of Bengal (India). HYDROBIOLOGIA 431(2-3): 225-241.
- Deibel, D. and Lee, S. H. 1992. Retention efficiency of sub-micrometer particles by the pharyngeal filter of the pelagic tunicate *Oikopleura vanhoeffeni*. Mar. Eco. Prog. Ser. 81: 25-30.
- Diez, B., Pedros-Alio, C. and Massana, R. 2001. Study of genetic diversity of eukaryotic picoplankton in different oceanic regions by small-subunit rRNA gene cloning and sequencing. Appl. Environ. Microbiol. 67(7): 2932-2941.
- Ducklow, H. W. and Carlson, C. A. 1992. Oceanic bacterial production. Adv. Microb. Ecol. 12: 113. Cited in D. M. Alongi (ad.), Coastal Ecosystem Processes. pp. 194-198. Florida: CRC Press. 1998.
- Ducklow, H. W., Purdie, D. A. Williams, P. J. L. and Davies, J. M. 1986. Bacterioplankton: A Sink for Carbon in a Coastal Marine plankton Community. Science 232: 865-867.
- Ducklow, H. W. and Shiah, F. K. 1993. Bacterial production in estuaries. In T. E. Ford (ed.), Aquatic Microbiology: An ecological approach, Boston: Blackwell, chap. 11. Cited in D. M. Alongi (ad.), Coastal Ecosystem Processes. pp. 194-198. Florida: CRC Press, 1998.
- Egan B. and Floodgate, G. D. 1985. Biological studies in the vicinity of shallow-sea tidal mixing front. II. The distribution of bacteria. Phil. Trans. R. Soc. Lond. B. 310: 435-444. Cited in Fogg, G. E. Some comments on picoplankton and its importance in the pelagic ecosystem. Aquat. Microb. Ecol. 9: 33-39, 1995.
- Fogg, G. E. 1986a. Picoplankton. In F. Megusar. and M. Gantar (eds.), Perspectives in microbial ecology. pp. 96-100. Ljubljana: Slovene Society for Microbiology. Cited in Fogg, G. E. Some comments on picoplankton and its importance in the pelagic ecosystem. Aquat. Microb. Ecol. 9: 33-39, 1995.
- Fogg, G. E. 1986b. Review Lecture: Picoplankton. Proc. R. Soc. Lond. B. 228: 1-30.
- Fogg, G. E. 1995. Some comments on picoplankton and its importance in the pelagic ecosystem. Aquat. Microb. Ecol. 9: 33-39.
- Fuhrman, J. A. and Noble, R. T. 1995. Virus and protists cause similar bacterial mortality in coastal seawater. Limnol. Oceanogr. 40(7): 1236-1242.

- Gin, K. Y. H., Lin, X. and Zhang, S. 2000. Dynamics and size structure of phytoplankton in the coastal of Singapore. Journal of Plankton Research 22(8): 1465-1484.
- Giovanoni, S. and Rappe, M. 2000. Evolution, diversity, and molecular ecology of marine prokaryotes. In D. L. Kirchman (ed.), Microbial Ecology of the Oceans, pp. 47-84. USA: Wiley-Liss.
- Goldman, J. C., Caron, D. A. and Dennett, M. R. 1987. Regulation of gross growth efficiency and ammonium regeneration in bacteria by substrate C:N ratio. Limnol. Oceanogr. 32: 1239-1252.
- Goldman, J. C. and Dennett, M. R. 1991. Ammonium regeneration and carbon utilization by marine bacteria grown on mixed substrates. Mar. Biol. 109: 369-378. Cited in Cherrier, J., Bauer, J. E. and Druffel E. R. M. Utilization and turnover of labile dissolved organic matter by bacterial heterotrophs in eastern North Pacific surface waters. Mar. Eco. Prog. Ser. 139: 267-279, 1996.
- Guillou, L., Chretiennot-Dinet, M., Boulben, S., Staay, S. Y. M. and Vaultot, D. 1999a. *Symbiomonas scintillans* gen. et sp. nov. and *Picophagus flagellatus* gen. et. sp. Nov. (Heterokonta): Two new heterotrophic flagellates of picoplanktonic size. Protist. 150: 383-398.
- Guillou, L., Chretiennot-Dinet, M., Medlin, L.K., Claustre, H., Goer, S. L. and Vaultot, D. 1999b. *Bolidomonas*: A new genus with two species belonging to a new algal class, the bolidophyceae (Heterokonta). J. Phycol. 35: 368-381.
- Guillou, L., Jacquet, S., Chretiennot-Dinet, M. J. and Vaultot, D. 2001. Grazing impact of two small heterotrophic flagellates on *Prochlorococcus* and *Synechococcus*. Aquat. Microb. Ecol. 26 (2): 201-207.
- Hamasaki, K., Satoh, F., Kikuchi, T., Toda, T. and Taguchi, S. 1999. Biomass and production of cyanobacteria in a coastal water of Sagami Bay, Japan. Journal of Plankton Research 21 (8):1538-1591.
- Healey, M. J., Moll, R. A. and Da'Állo, C. O. 1988. Abundance and distribution of bacterioplankton in the Gambia River, West Africa. Microbial Ecology 16: 291-310. Cited in Robertson, A. I. and Blaber, S. J. M. Plankton, Epibenthos and Fish Communities. In A. I. Robertson and D. M. Alongi (eds.), Coastal and Estuarine Studies; 41, pp. 173-224. American Geophysical Union: Washington DC, 1992.

- Herbland, A. and Le Bouteiller, A. 1981. The size distribution of phytoplankton and particulate organic matter in the equatorial Atlantic Ocean: Importance of ultraseston and consequences. Journal of Plankton Research 3: 659-673. Cited in Stockner, J. G. and Antia, N. J. Algal picoplankton from marine and freshwater ecosystems: A multidisciplinary perspective. Can. J. Fish. Aquat. Sci. 43: 2472-2503, 1986.
- Iriarte, A. and Purdie, D. A. 1994. Size distribution of chlorophyll *a* biomass and primary production in a temperate estuary (Southampton Water): the contribution of photosynthetic picoplankton. Mar. Eco. Prog. Ser. 115: 283-297.
- Jeffrey, S. W. and Hallgraeff, G. M. 1990. Phytoplankton ecology of Australian waters. In M. N. Clayton and R. J. King (eds.), The Biology of Marine Plants, pp. 310-348. Melbourne: Longman Cheshire. Cited in Jeffrey, S. W., and Veski, M. Introduction to marine phytoplankton and their pigment signatures. In S. W. Jeffrey, R. F. C. Mantoura, and S. W. Wright (eds.), Phytoplankton pigments in oceanography: guidelines to modern methods. pp. 37-84. Paris: UNESCO Publishing, 1997.
- Jochem, F. J. 2001. Morphology and DNA content of bacterioplankton in the northern Gulf of Mexico: analysis by epifluorescence microscopy and flow cytometry. Aquat Microb Ecol. 25: 179-194.
- Jochem, F. J. (n. d.). Bacterioplankton in the western Gulf of Mexico: Analysis by epifluorescence microscopy and flow cytometry. [Online]. Available from: <http://www.jochemnet.de/bact.html> [2002, December 12].
- Johnson, P. W. and Sieburth, J. McN. 1982. In-situ morphology and occurrence of eukaryotic phototrophs of bacterial size in the picoplankton of estuarine and oceanic waters. J. Phycol. 18: 318-327.
- Johnson, P. W., Xu, H. S. and Sieburth, J. McN. 1982. The utilization of chroococcoid cyanobacteria by marine protozooplankters but not by calanoid copepods. Ann. Inst. Oceanogr. Paris 58(S): 297-308. Cited in Stockner, J. G. and Antia, N. J. Algal picoplankton from marine and freshwater ecosystems: A multidisciplinary perspective. Can. J. Fish. Aquat. Sci. 43: 2472-2503, 1986.
- Kawachi, M., Atsumi, M., Ikemoto, H. and Miyachi, S. 2002. *Pinguiochrysis pyriformis* gen. et sp. nov. (Pinguiophyceae), a new picoplanktonic alga isolated from the Pacific Ocean. Phycological Research 50 (1): 49-56.

- Kirchman, D. L. 1990. Limitation of bacterial growth by dissolved organic matter in subarctic Pacific. Mar. Eco. Prog. Ser. 62: 47-54.
- Kirchman, D. L. and Williams, P. J. le B. 2000. Introduction In D. L. Kirchman (ed.), Microbial Ecology of the Oceans, pp. 1-12. USA: Wiley-Liss.
- Kirchman, D., Sigda, J. Kapuscinski, R. and Mitchell, R. 1982. Statistical analysis of the direct count method for enumerating bacteria. Appl. Environ. Microbiol. 44(3): 376-382.
- Kroer, N. 1993. Bacterial growth efficiency on natural dissolved organic matter. Limnol. Oceanogr. 38: 1282-1290.
- Kuylenstierna, M. and Karlson, B. (n.d.). Department of Marine Botany, Goteborg University, Sweden [Online]. Available from:
http://www.marbot.gu.se/sss/others/Synechococcus_sp.html [2002, December 12].
- Lalli, C. M. and Parsons, T. R. 1993. Biological Oceanography: An Introduction, pp. 22-43. New York : Pergamon Press.
- Lande, R. Li, W. K. W., Horn, E. P. W. and Wood, A. M. 1989. Phytoplankton growth rates estimated from depth profiles of cell concentration and turbulent diffusion. Deep Sea Res. 36: 1141-1159. Cited in Fogg, G. E. Some comments on picoplankton and its importance in the pelagic ecosystem. Aquat. Microb. Ecol. 9: 33-39, 1995.
- Larsson, U. and Hagstorm, A. 1982. Fractionated phytoplankton primary production, exudate Release and Bacterial Production in a Baltic Eutrophication Gradient. Limnol. Oceanogr. 67: 57-70.
- Lee, C. W., Kudo, I., Yanada, M. and Maita, Y. 2001. Bacterial abundance and production and their relation to primary production in Funika Bay. Plankton Biol. Ecol. 48(1): 1-9.
- Le Gall, S., Hassen, M. B. and Le Gall, P. 1997. Ingestion of bacterivorous ciliate by the oyster *Grassostrea gigas*: protozoa as a trophic link between picoplankton and benthic suspension-feeders. Mar. Eco. Prog. Ser. 152: 301-306.
- Lipej, L., Mozetic, P., Turk, V. and Malej, A. 1997. The trophic role of the marine cladoceran *Penilia avirostris* in the Gulf of Trieste. HYDROBIOLOGIA 360 (1-3): 197-203.
- Li, W. K. W., Subba Rao, D. V., Harrison, W. G., Smith, J. C., Cullen, J. J., Irwin, B. and Platt, T. 1983. Autotrophic picoplankton in the tropical ocean. Science 219: 292-295.

- Lovejoy, C., Legendre, L., Therriault, J. C., Tremblay, J. E., Klein, B. and Ingram, R. G. 2000. Growth and distribution of marine bacteria in relation to nanoplankton community structure. DEEP-SEA RESEARCH PART II-TROPICAL STUDIES IN OCEANOGRAPHY. 47(3-4): 461-487.
- Maclsaac, E. A. and Stockner, J. G. 1993. Enumeration of phototrophic picoplankton by Autofluorescence microscopy. In P. F. Kemp, B. F. Sherr, E. B. Sherr and J. J. Cole (eds.), Handbook of Methods in Aquatic Microbial Ecology, pp. 187-197. Boca Raton: Lewis Publishers.
- Magazzu, G. and Decembrini, F. 1995. Primary production, biomass and abundance of phototrophic picoplankton in the Mediterranean Sea: A review. Aquat. Microb. Ecol. 9: 97-104.
- Marine Biotechnology Institute Culture Collection (MBIC). (n.d.). Catalog (bacteria, algae) [Online]. Marine Biotechnology Institute Co., Ltd. Available from: <http://www.mbio.co.jp> [2002, December 12].
- Metzler, P. M., Gilbert, P. M., Gaeta, S. A. and Ludlam, J. M. 2000. Contrating effects of substrate and grazer manipulations on picoplankton in oceanic and coastal waters off Brazil. Journal of Plankton Reseaech 22(1): 77-90.
- Morel, A. and Bricaud, A. 1981. Deep-Sea Res. 28: 1375. Cited in Li, W. K. W., Subba Rao, D. V., Harrison, W. G., Smith, J. C., Cullen, J. J., Irwin, B. and Platt, T. Autotrophic picoplankton in the tropical ocean. Science 219: 292-295, 1983.
- Munk, P., Janekarn, V., Nielsen, T., Boonruang, P., Sawangarreruks, S., Satapoomin, S. and Limtrakulvong. 2002. Proceedings of the Regional Workshop on Ecology of Tropical Mesoplankton and Fish Larvae. Phuket Marine Biological Center Special Publication (27): pp. 44.
- Nakamura, Y., Suzuki, K., Suzuki, S. and Hiromi, J. 1997. Production of *Oikopleura dioica* (Appendicularia) following a picoplankton 'bloom' in eutrophic coastal area. Journal of Plankton Reseaech 19(1): 113-124.
- Neilson, A. H. and Larsson, T. 1980. The utilization of organic nitrogen for growth of algae: Physiological aspects. Physiol. Plant. 48: 542-553. Cited in Stockner, J. G. Phototrophic picoplankton: An overview from marine and freshwater ecosystems. Limnol. Oceanogr. 33(4,part 2): 765-775, 1988.

- Ning, X., Cloern, J. E. and Cole, B. E. 2000. Spatial and temporal variability of picocyanobacteria *Synechococcus* sp. In San Francisco Bay. Limnol. Oceanogr. 45 (3): 695-702.
- Noji, T. T. , Bathmann, U. V., Bodungen, B. V., Voss, M., Antia, A., Krumbholz, M., Klein, B., Peeken, I., Noji, C. I. M. and Rey, F. 1997. Clearance of picoplankton-sized particles and formation of rapidly sinking aggregates by the pteropod, *Limacina retroversa*. Journal of Plankton Research 19(7): 863-875.
- Odate, T. and Fukuchi, M. 1994. Surface distribution of picophytoplankton along the first leg of the JARE-33 cruise, from Tokyo to Fremantle, Australia. Bulletin of Plankton Society of Japan 41(2): 93-104.
- Palenik, B. P. 1999. Picophytoplankton seasonal cycle at the Sio pier, La Jolla, California. J. Phycol. 14(12): 53-53
- Parsons, T. R., Maita, Y. and Lalli, C. M. 1984. A Manual of Chemical and Biological Methods For Seawater Analysis. Oxford: Pergamon Press.
- Pedros-Alio, C., Calderon-Paz, J. I., Guixa-Boixereu, N., Estrada, M. and Gasol, J. M. 1999. Bacterioplankton and phytoplankton biomass and production during summer stratification in the northwestern Mediterranean Sea. Deep Sea Res.1 Oceanogr. Res. 46 (6): 985-1019.
- Péna, G. 1989. Sels nutritifs et micropolluants métalliques dans un écosystème lagunaire: l'étang de Thau. D. S. thesis, Univ. Montpellier II. 143 p. Cited in Vaquer, A., Troussellier, M., Courties, C. and Bibent, B. Standing stock and dynamics of picophytoplankton in the Thau Lagoon (northwest Mediterranean coast). Limnol. Oceanogr. 41(8): 1821-1828, 1996.
- Piumsomboon, T., Paphavasit, N. and Phromthong, I. 2003. Biological Mediated Carbon flowing Pelagic Ecosystem of Bandon Bay, Southern Thailand. NAT. HIST. J. CHULALONGKORN UNIV. (submitted)
- Platt, T., Subba-Rao, D. V. and Irwin, B. 1983. Photosynthesis of picoplankton in oligotrophic ocean. Nature (Lond.) 301: 702-704. Cited in Stockner, J. G. and Antia, N. J. Algal picoplankton from marine and freshwater ecosystems: A multidisciplinary perspective. Can. J. Fish. Aquat. Sci. 43: 2472-2503, 1986.
- Pomeroy, L. R. 1974. The Ocean's Food Web, A Changing Paradigm. BioScience. 24(9): 499-504.

- Pomeroy, L. R. 1974. The Ocean's Food Web, A Changing Paradigm. BioScience. 24(9): 499-504.
- Raven, J. A. 1998. The twelfth transley lecture. Small is beautiful: the picophytoplankton. Functional Ecology 12: 503-513.
- Robarts, R. D., Zohary, T., Waiser, M. J. and Yacobi, Y. Z. 1996. Bacterial abundance, biomass, and production in relation to phytoplankton biomass in the Levantine Basin of the southeastern Mediterranean Sea. Mar. Eco. Prog. Ser. 137: 273-281.
- Robertson, A. I. and Blaber, S. J. M. 1992. Plankton, Epibenthos and Fish Communities. In I. Robertson, and D. M. Alongi (eds.), Coastal and Estuarine Studies; 41, pp. 173-224. American Geophysical Union, Washington DC.
- Roff, J. C., Turner, J. T., Webber, M. K. and Hopcroft, R. R. 1995. Bacterivory by tropical copepod nauplii: extent and possible significance. Aquat. Microb. Ecol. 9(2): 165-175.
- Saijo, Y. and Takesue, K. 1965. Further studies on the size distribution of photosynthesizing phytoplankton in the Indian Ocean. J. Oceanogr. Soc. Jpn. 20: 264-271. Cited in Stockner, J. G. and Antia, N. J. Algal picoplankton from marine and freshwater ecosystems: A Multidisciplinary Perspective. Can. J. Fish. Aquat. Sci. 43: 2472-2503, 1986.
- Sanders, R. W., Berninger, U. G., Lim, E. L., Kemp, P. F. and Caron, D. A. 2000. Heterotrophic and mixotrophic nanoplankton predation on picoplankton in the Sargasso Sea and on Georges Bank. Mar. Eco. Prog. Ser. 192: 103-118.
- Selmer, J. S., Ferrier-Pages, C., Cellario, C. and Rassoulzadegan, F. 1993. New and regenerated production in relation to the microbial loop in the NW Mediterranean Sea. Mar. Eco. Prog. Ser. 100: 71-83. Cited in Iriarte, A. and Purdie, D. A. Size distribution of chlorophyll *a* biomass and primary production in a temperate estuary (Southampton Water): the contribution of photosynthetic picoplankton. Mar. Eco. Prog. Ser. 115: 283-297, 1994.
- Sherr, E. B., Caron, D. A. and Sherr, B. F. 1993. Staining of heterotrophic protists for visualization via epifluorescence microscopy. In P. F. Kemp, B. F. Sherr, E. B. Sherr and J. J. Cole (eds.), Handbook of Methods in Aquatic Microbial Ecology, pp. 213-227. Boca Raton: Lewis Publishers.
- Sherr, E. and Sherr, B. 2000. Marine microbes. In D.L. Kirchman (ed.), Microbial ecology of the oceans, pp. 13-46. New York: WILEY-LISS.

- Sheldon, R. W., Rassoulzadegan F., Azam, F., Berman, T. Bezanson, D. S., Bianchi, M., Bonin, D., Hagstrom, A., Laval-Peuto, M., Neveux, J., Raimbault, P., Rivier, A., Sherr, B., Sherr, E., Van Wambeke, F., Wikner, J., Wood, A. M. and Yentsch, C. M. 1992. Nano- and picoplankton growth and production in the Bay of Villefranche-sur-Mer (NW Mediterranean). HYDROBIOLOGIA 241: 91-106. Cited in Fogg, G. E. 1995. Some comments on picoplankton and its importance in the pelagic ecosystem. Aquat. Microb. Ecol. 9: 33-39.
- Shiah, F. K., Liu, K. K. and Gong, G. Ch. 1999. Temperature versus substrate limitation of heterotrophic bacterioplankton production across trophic and temperature gradients in the East China Sea. Aquat. Microb. Ecol. 17(3): 247-254.
- Shiomoto, A., Tadokoro, K. Monaka, K. and Nanba, M. 1997. Productivity of picoplankton compares with that of large phytoplankton in the subarctic region. Journal of Phytoplankton Research 19(7): 907-916.
- Sieburth, J. McN, Smetacek, V. and Lenz, J. 1978. Pelagic ecosystem structure: heterotrophic compartments of plankton and their relationship to plankton size fractions. Limnol. Oceanogr. 23: 1256-1263. Cited in Graham, J. M. Symposium introductory remarks: A brief history of aquatic microbial ecology. J. Protozool. 38(1): 66-69, 1991.
- Sieracki, M. E. and Wong, M. W. (n.d.). Image gallery [Online]. Bigelow laboratory for Ocean Sciences. Available from: http://www.bigelow.org/cytometry/Image_gallery/SYN.html [2002, December 12].
- Sin, Y., Wetzel, R. L. and Anderson, I. C. 2000. Seasonal variations of size-fractionated phytoplankton along the salinity gradient in the York River estuary, Virginia (USA). Journal of Phytoplankton Research 22(10): 1945-1960.
- Station Biologique de Roscoff. (n.d.). Picoplankton Gallery [Online]. Station Biologique de Roscoff. Available from: ([www.sb-roscoff.fr/Phyto/phyto fr.html](http://www.sb-roscoff.fr/Phyto/phyto_fr.html)) [2002, December 12].
- Stickland, J. D. H. and Parsons, T. R. 1972. A practical handbook of seawater analysis. Fisheries Research Board of Canada Bull. 167: 1-308.
- Stockner, J. G. 1988. Phototrophic picoplankton: An overview from marine and freshwater ecosystems. Limnol. Oceanogr. 33(4, part 2): 765-775.
- Stockner, J. G. and Antia, N. J. 1986. Algal picoplankton from marine and freshwater Ecosystems: A Multidisciplinary Perspective. Can. J. Fish. Aquat. Sci. 43: 2472-2503.

- Strom, S. L., Benner, R., Ziegler, S. and Dagg, M. J. 1997. Planktonic grazers are a potentially important source of marine dissolved organic carbon. Limnol. Oceanogr. 42(6): 1364-1374.
- Tada, K., Yamada, M., Takemura, A. and Nakano, Y. 1999. Size distribution of phytoplankton community in oligotrophic tropical coastal waters. La mer 36:139-145.
- Takahashi, M., Kikuchi, K. and Hara, Y. 1985. Importance of picocyanobacteria biomass (unicellular, blue-green algae) in the phytoplankton population of the coastal waters off Japan. Marine Biology 89: 63-69.
- Takahashi, M., Koike, I., Iseki, K., Bienfang, P. K. and Hattori, A. 1982. Phytoplankton species' responses to nutrient changes in experimental enclosures and coastal waters. In G. D. Grice and M. R. Reeve (eds.), Marine mesocosms. pp. 333-340. New York, Springer-Verlag. Cited in Fogg, G. E. Some comments on picoplankton and its importance in the pelagic ecosystem. Aquat. Microb. Ecol. 9: 33-39, 1995.
- Tanaka, T. and Taniguchi, A. 1996. Short-term variation in abundance of bacteria and heterotrophic nanoflagellates in summer observed in Onagawa Bay, Japan. Bulletin of Plankton Society of Japan 43(1): 21-29.
- Thingstad, T. F., Zweifel, U. L. and Rassoulzadegan, F. 1998. P limitation of heterotrophic bacteria and phytoplankton in the northwest Mediterranean. Limnol. Oceanogr. 43 (1): 88-94.
- Thomsen, H. A. 1986. A survey of the smallest eukaryotic organisms of the marine phytoplankton. Can. Bull. Fish. Aquat. Sci. 214: 121-158. Cited in Diez, B., Pedros-Alio, C., and Massana, R. Study of genetic diversity of eukaryotic picoplankton in different oceanic regions by small-subunit rRNA gene cloning and sequencing. Appl. Environ. Microbiol. 67(7): 2932-2941, 2001.
- Tomaru, Y., Kawabata, Z. and Nakano, S. 2000. Consumption of picoplankton by bivalve larvae of Japanese pearl oyster *Pinctada fucata martensii*. Mar. Eco. Prog. Ser. 192: 195-202.
- Tomas, C. R. 1997. Identifying marine phytoplankton. USA: Academic Press.
- Torretón, J. 1999. Biomass, production and heterotrophic activity of bacterioplankton in the Great Astrolabe Reef Lagoon (Fiji). Coral-Reefs 18(1): 43-53.

- Turley, C. M., Bianchi, M., Christaki, U., Conan, P., Harris, J. R. W., Psarra, S., Ruddy, G., Stutt, E. D., Tselepidis, A. and Van Wambeke, F. 2000. Relationship between primary producers and bacteria in an oligotrophic sea - the Mediterranean and biogeochemical implications. Mar. Ecol. Prog. Ser. 193: 11-18.
- Vaquer, A., Troussellier, M., Courties, C. and Bibent, B. 1996. Standing stock and dynamics of picophytoplankton in the Thau Lagoon (northwest Mediterranean coast). Limnol. Oceanogr. 41(8): 1821-1828.
- Waterbury, J. B. 1986. Introduction to the picoplankton: The prokaryotes. In T. Platt (ed.), Proceedings of the NATO Picoplankton Advanced Study Institute, Can. Bull. Fish. Aquat. Sci. Cited in Fogg, G. E. Review lecture: Picoplankton. Proc. R. Soc. Lond. B. 228: 1-30, 1986.
- Waterbury, J. B., Watson, S. W., Valois, F. W. and Franks, D. 1986. Biological and ecological characterization of marine unicellular cyanobacterium *Synechococcus*. In T. Platt, and W. K. W. Li (eds.), Photosynthetic picoplankton. Can. Bull. Fish. Aquat. Sci. 241: 71-120. Ontario: Canadian Government Publishing Center
- Weinbauer, M. G. and Peduzzi, P. 1995. Effect of virus-rich high molecular weight concentrates of seawater on the dynamics of dissolved amino acids and carbohydrates. Mar. Ecol. Prog. Ser. 127: 245-253. Cited in D. L. Kirchman (ed.), Marine Ecology of the ocean. New York: WILEY-LISS, 2000.
- Ypma, J. E. and Throndsen, J. 1996. Seasonal dynamics of bacteria, autotrophic picoplankton and small nanoplankton in the inner Oslofjord and the Skagerrak in 1993. SARSIA 81(1): 1-80.



ภาคผนวก

ศูนย์วิทยทรัพยากร
จุฬาลงกรณ์มหาวิทยาลัย

ภาคผนวก ก ปริมาณน้ำฝนในเขตจังหวัดสมุทรสงครามระหว่างปี พ.ศ. 2544 ถึงพ.ศ. 2545

ปี	เดือน												ทั้งปี
	ม. ค.	ก. พ.	มี. ค.	เม. ย.	พ. ค.	มิ. ย.	ก. ค.	ส. ค.	ก. ย.	ต. ค.	พ. ย.	ธ. ค.	
พ.ศ. 2544													
ปริมาณน้ำฝนเฉลี่ย (มิลลิเมตร)	5.7	0.0	135.4	45.9	117.4	71.4	96.7	56.9	209.1	200.5	9.6	7.5	956.1
จำนวนวันที่มีฝนตก (วัน)	1.5	0.0	8.5	2.0	12.0	8.5	12.0	12.0	14.5	22.0	3.5	2.5	99.0
ปริมาณน้ำฝนในวันที่ฝนตกมากที่สุด (มิลลิเมตร)	7.5	0.0	45.5	38.7	30.5	28.4	26.5	21.3	60.5	53.4	5.6	7.5	60.5
พ.ศ. 2545													
ปริมาณน้ำฝนเฉลี่ย (มิลลิเมตร)	1.2	0.0	4.9	2.3	215.1	147.0	76.5	79.8	142.0	213.3	107.2	19.4	1008.7
จำนวนวันที่มีฝนตก (วัน)	0.3	0.0	2.3	1.0	15.0	17.0	9.7	13.0	11.5	8.0	6.5	3.0	87.3
ปริมาณน้ำฝนในวันที่ฝนตกมากที่สุด (มิลลิเมตร)	3.6	0.0	6.5	2.7	80.5	32.5	30.5	26.0	60.5	120.5	55.0	12.0	120.5

หมายเหตุ: ปริมาณน้ำฝนเฉลี่ยได้จากการหาค่าเฉลี่ยของปริมาณน้ำฝนรวมในแต่ละเดือนจาก 3 สถานี ได้แก่ อำเภอเมือง อำเภออัมพวาและอำเภอบางคนที จังหวัดสมุทรสงคราม (กรมอุตุนิยมวิทยา, 2546)

ประวัติผู้เขียนวิทยานิพนธ์

นางสาววรพร ธารางกูร เกิดวันที่ 28 ธันวาคม พ.ศ. 2519 ที่อำเภอเมือง จังหวัดกระบี่ สำเร็จการศึกษาระดับมัธยมตอนต้นจากโรงเรียนอมาตย์พานิชนุกูล จังหวัดกระบี่ ในปีการศึกษา 2534 สำเร็จการศึกษาระดับมัธยมตอนปลายจากโรงเรียนภูเก็ตวิทยาลัย จังหวัดภูเก็ต ในปีการศึกษา 2537 สำเร็จการศึกษาระดับปริญญาวิทยาศาสตรบัณฑิต เกียรตินิยมอันดับสอง สาขาวิชาวิทยาศาสตร์ คณะทรัพยากรธรรมชาติ มหาวิทยาลัยสงขลานครินทร์ วิทยาเขตหาดใหญ่ จังหวัดสงขลา ในปีการศึกษา 2541 หลังจากนั้นได้เข้าศึกษาต่อในหลักสูตรปริญญาวิทยาศาสตรมหาบัณฑิต สาขาชีววิทยาทางทะเล ภาควิชาวิทยาศาสตร์ทางทะเล คณะวิทยาศาสตร์ จุฬาลงกรณ์มหาวิทยาลัย ในปีการศึกษา 2542



ศูนย์วิทยทรัพยากร
จุฬาลงกรณ์มหาวิทยาลัย