

REFERENCES

- Archarya, S., and Bussel, J.B. Hematologic toxicity of sodium valproate. *J. Pediatr. Hematol. Oncol.* 22(1)(2000) : 62-65.
- Amara, S.G., and Fontana, A.C. Excitatory amino acid transporters : keeping up with glutamate. *Neurochem. Int.* 41(5)(2002) : 313–318.
- Annegers, J.F. The epidemiology of epilepsy. In E. Wyllie (ed.). *The Treatment of Epilepsy*, pp.157–164. Lea and Febiger : Philadelphia, 1993.
- Aragón, C., and López-Corcuera, B. Structure, function and regulation of glycine neurotransmitters. *Eur. J. Pharmacol.* 479(1-3)(2003) : 249-262.
- Arzimanoglou, A., Hirsch, E., Nehlig, A., Castelnau, P., Gressens, P., and Vasconcelos, A.P. Epilepsy and neuroprotection : an illustrated review. *Epileptic Disord.* 4(3)(2002) : 173-182.
- Avanzini, G., and Franceschetti, S. Cellular biology of epileptogenesis. *Lancet Neurol.* 2(2003) : 33-42.
- Awad, I.A., Rosenfeld, J., Ahl, H., Hahn, J.F., and Luders, H. Intractable epilepsy and structural lesions of the brain : mapping, resection strategies, and seizure outcome. *Epilepsia* 32(1991) : 179-186.
- Baker, D.A., Xi, Z., Shen, H., Swanson, C.J., and Kalivas, P.W. The origin and neuronal function of in vivo nonsynaptic glutamate. *J. Neurosci.* 22(2002) : 9134–9141.
- Balcar, V.J. Molecular pharmacology of the Na^+ -dependent transport of acidic amino acids in the mammalian central nervous system. *Biol. Pharm. Bull.* 25(2002) : 291–301.
- Bardford, H.F. Glutamate, GABA and epilepsy. *Prog. Neurobiol.* 47(1995) : 477-511.
- Bashkatova, V., Narkevich, V., Vitskova, G., and Vanin, A. The influence of anticonvulsant and antioxidant drugs on nitric oxide level and lipid peroxidation in the rat brain

- during pentylenetetrazole-induced epileptiform model seizures. *Prog. Neuropsychopharmacol. Biol. Psychiatry.* 27(2003) : 487-492.
- Bazil, C.W., and Pedley, T.A. Advances in the medical treatment of epilepsy. *Annu. Rev. Med.* 45(1998) : 135–162.
- Beck, H., Steffens, R., Elger, C.E., and Heinemann, U. Voltage-dependent Ca^{2+} currents in epilepsy. *Epilepsy Res.* 32(1998) : 321–332.
- Bellissimo, M.I., Amado, D., Abdalla D.S., Ferreira, E.C., Cavalheiro, E.A., and Naffah-Mazzacoratti, M.G. Superoxide dismutase, glutathione peroxidase activities and hydroperoxide concentration are modified in the hippocampus of epileptic rats. *Epilepsy Res.* 46(2001) : 121-128.
- Ben-Ari, Y. Limbic seizure and brain damage produced by kainic acid : mechanisms and relevance to human temporal lobe epilepsy. *Neuroscience* 14(1985) : 375-403.
- Bertrand, S., Weiland, S., Berkovic, S.F., Steinlein O.K., and Bertrand D. Properties of neuronal nicotinic acetylcholine receptor mutants from humans suffering from autosomal dominant nocturnal frontal lobe epilepsy. *Br. J. Pharmacol.* 125(1998) : 751–760.
- Bezzi, P., Domercq, M., Vesce, S., and Volterra, A. Neuron–astrocyte cross-talk during synaptic transmission: physiological and neuropathological implications. *Prog. Brain Res.* 132(2001) : 255–265.
- Biervert, C., Schroeder, B.C., Kubisch, C., Berkovic S.F., Propping, P., Jentsch, T.J., and Steinlein, O.K. A potassium channel mutation in neonatal human epilepsy. *Science* 279(1998) : 403–406.
- Bittigau, P., ifringer, M., Genz, K., Reith, E., Pospischill, D., Govindarajalu, S., Dzietko, M., Pesditschek, S., Mai, I., Dikranian, K., Olney, J.W., and Ikonomidou, C. Antiepileptic drugs and apoptotic neurodegeneration in the developing brain. *Proc. Natl. Acad. Sci. USA* 99(23)(2002) : 15089-15094.

- Bonan, C.D., Walz, R., Pereira, G.S., Worm, P.V., Battastini, A.M.O., Cavalheiro, E.A., Izquierdo, I., and Sarkis, J.J.F. Changes in synaptosomal ectonucleotidase activities after status epilepticus induced by pilocarpine and kainic acid. *Epilepsy Res.* 39(2000) : 229–238.
- Bonfoco, E., Krainic, D., Ankarcrona, M., Nocotera, P., and Lipton, A.S. Apoptosis and necrosis : two distinct events induced, respectively, by mild and intense insults with *N*-methyl-*D*-aspartate or nitric oxide/superoxide in cortical cell cultures. *Proc. Natl. Acad. Sci. USA* 92(1995) : 7162–7166.
- Bradford, H.F. Glutamate, GABA and epilepsy. *Prog. Neurobiol.* 47(1995) : 477–511.
- Brodie, M.J. Do we need any more new antiepileptic drugs? *Epilepsy Res.* 45(2001) : 3–6.
- Bruce, A.J. and Baudry, M. Oxygen free radicals in rat limbic structures after kainic induced seizures. *Free Radic. Biol. Med.* 18(1995) : 993-1002.
- Bruhn, T., Cobo, M., Berg, M., and Diemer, N.H. Limbic seizure-induced changes in amino-acid levels in the hippocampal formation : a microdialysis study of freely moving rats. *Acta Neurol. Scand.* 86(1992) : 455-461.
- Bruton, C.J., Stevens, J.R., and Frith, C.D. Epilepsy, psychosis, and schizophrenia: clinical and neuropathologic correlations. *Neurology* 44(1994) : 34-42.
- Carlson, H., Ronne-Engström, E., Ungerstedt, U., and Hillered, I. Seizure related elevations of extracellular amino acids in human focal epilepsy. *Neurosci. Lett.* 140(1992) : 30-32.
- Cascino, G.D. Surgical treatment for epilepsy. *Epilepsy Res.* 60(2004) : 179-186.
- Cascino, G.D., Jack, Jr., and Parisi, J.E. MRI in the presurgical evaluation of patients with frontal lobe epilepsy and children with temporal lobe epilepsy : pathological correlation and prognostic importance. *Epilepsy Res.* 11(1992) : 51–59.

- Cavalheiro, E.A., Leite, J.P., Bortolotto, Z.A., Turski, W.L., Ikonomidou, C., and Turski, L. Long-term effect of pilocarpine in rats : structural damage of the brain triggers kindling and spontaneously recurrent seizures. *Epilepsia* 32(1991) : 778-782.
- Cavalheiro, E.A., Fernandes, M.T., Turski, L., and Naffah-Mazzacoratti, M.G. Spontaneous recurrent seizures in rats : amino acids and monoamines determination in the hippocampus. *Epilepsia* 35(1994) : 1-11.
- Chagnac-Amitai, Y., and Connors, B.W. Synchronized excitation and inhibition driven by intrinsically bursting neurons in neocortex. *J. Neurophysiol.* 62(1989) : 1149-1162.
- Chance, E., and Williams, G.R. The respiratory chain and oxidative phosphorylation. *Adv. Enzymol.* 17(1956) : 65-134.
- Chapman, V., and Dickenson, A.H. Time-related roles of excitatory amino acid receptors during persistent noxiously evoked responses of rat dorsal horn neurones. *Brain Res.* 703(1995) : 45-50.
- Chaudieu, I., Rondouin, G., Chicheportiche, M., and Chicheportiche, R. Presynaptic alteration of [³H] GABA transport in hippocampus by amygdala kindling. *Neurosci. Lett.* 76(1987) : 329-334.
- Clifford, D.B., Olney, J.W., Collins, R.C., and Zorumski, C.F. The functional anatomy and pathology of lithium-pilocarpine and high-dose of pilocarpine seizures. *Neuroscience* 23(1987) : 953- 967.
- Cock, H.R. The role of mitochondria and oxidative stress in neuronal damage after brief and prolonged seizures. *Prog. Brain Res.* 135(2002a) : 187-196.
- Cock, H.R., Tong, X., Hargreaves, I.P., Heales, S.J.R., Clark, J.B., Patsalos, P.N., Thom, M., Groves, M., Schapira, A.H.V., Shorvon, S.D., and Walker, M.C. Mitochondrial dysfunction associated with neuronal death following status epilepticus in rat. *Epilepsy Res.* 48(2002b) : 157-168.

Commission on Classification and Terminology of the International League Against Epilepsy.

Proposal for revised clinical and electroencephalographic classification of epileptic seizures. *Epilepsia* 22(1981) : 489-501.

Commission on Classification and Terminology of the International League Against Epilepsy.

Proposal for revised classification of epilepsies and epileptic syndromes. *Epilepsia* 30(1989) : 389-399.

Cooper, J.R., Bloom, F.E., and Roth, R.H. Amino acid transmitters. In J.R. Cooper, F.E. Bloom, and R.H. Roth (eds.). *The biochemical basis of neuropharmacology* (8th ed.), pp.105-150. New York : Oxford University Press, 2003.

Costa, M.S., Rocha, J.B.T., Perosa, S.R., Cavalheiro, E.A., and Naffah-Mazzacoratti, M.G. Pilocarpine-induced status epilepticus increases glutamate release in rat hippocampal synaptosomes. *Neurosci. Lett.* 356(2004) : 41-44.

Covolan, L., Smith, R.L., and Mello, L.E. Ultrastructural identification of dentate granule cell death from pilocarpine-induced seizures. *Epilepsy Res.* 41(2000) : 9-21.

D'Souza, S. W., McConnell, S. E., Slater, P., and Barson, A. J. Glycine site of the excitatory amino acid N-methyl-D-aspartate receptor in neonatal and adult brain. *Arch. Dis. Child.* 69(1993) : 212-215.

Dal-Pizzol, F., Klamt, F., Vianna, M.M.R., Schroder, N., Quevedo, J., Benfato, M., Moreira, J.C.F., and Walz, R. Lipid peroxidation in hippocampus early and late after status epilepticus induced by pilocarpine or kainic acid in wistar rats. *Neurosci. Lett.* 291(2000) : 179-182.

Danbolt, N.C. Glutamate uptake. *Prog. Neurobiol.* 65(2001) : 1-105.

DeLorenzo, R.J. The challenging genetics of epilepsy. *Epilepsy Res.* 4(Suppl.)(1991) : 3-17.

DeLorenzo, R.J., Hauser, W.A., Towne, A.R., Boggs, J.G., Pellock, J.M., Penberthy, L., Garnett, L., Fortner, C.A., Ko, D. A prospective, population-based epidemiologic study of status epilepticus in Richmond, Virginia. *Neurology* 46(1996) : 1029-1035.

- Dhillon, S., and Sander, J.W.A.S. Epilepsy. In R. Walker, and C. Edwards (eds.). *Clinical Pharmacy and Therapeutics 2nd ed.*, pp.435-451. Edinburgh : Churchill Livingstone, 1999.
- Draper, H.H., and Hadley, M. Malondialdehyde determination as index of lipid peroxidation. *Methods Enzymol.* 186(1990) : 421-431.
- Duchen, M.R. Ca⁺⁺-dependent changes in the mitochondrial energies in single dissociated mouse sensory neurons. *Biochem.* 283(1992) : 41-58.
- Dumoulin, A., Rostaing, P., Bedet, C., Levi, S., Isambert, M.F., Henry, J.P., Triller, A., and Gasnier, B. Presence of the vesicular inhibitory amino acid transporter in GABAergic and glycinergic synaptic terminal boutons. *J. Cell. Sci.* 112(1999) : 811-823.
- During, M.J., and Spencer, D.D. Extracellular hippocampal glutamate and spontaneous seizure in the conscious human brain. *Lancet* 341(1993) : 1607-1610.
- Escayg, A., MacDonald, B.T., Meisler M.H., Baulac, S., Huberfeld, G., An-Gourfinkel, I., Brice, A., LeGuern, E., Moulard, B., Chaigne, D., Buresi, C., and Malafosse, A. Mutations of SCN1A, encoding a neuronal sodium channel, in two families with GEFS⁺. *Nat. Genet.* 24(2000) : 343-354.
- Estabrook, R.W. Mitochondrial respiratory control and the polarographic measurement of ADP:O ratios. In S.P. Colowick, and N.O. Kaplan (eds.). *Methods in enzymology Vol. X.*, pp.41-47. New York : Academic Press, 1967.
- Finsterer, J. Mitochondriopathies *Eur. J.. Neurol.* 11(2004) : 163-186.
- Foldvary, N., Bingaman, W.E., and Wyllie, E. Surgical treatment of epilepsy. *Neurol. Clin.* 19(2001) : 491-515.
- Franceschetti, S., Guatteo, E., Panzica, F., Sancini, G., Wanke, E., and Avanzini, G. Ionic mechanisms underlying burst firing in pyramidal neurons : intracellular study in at sensorimotor cortex. *Brain Res.* 69(1995) : 6127-6139.

- Freitas, R.M., Sousa, F.C.F., Vasconcelos, S.M.M., Viana, G.S.B., and Fonteles, M.M.F. Pilocarpine-induced status epilepticus in rats : lipid peroxidation level, nitrite formation, GABAergic and glutamatergic receptor alterations in the hippocampus, striatum and frontal cortex. *Pharmacol. Biochem. Behav.* 78(2004) : 327-332.
- Fujikawa, D.G. The temporal evolution of neuronal damage from pilocarpine-induced status epilepticus. *Brain Res.* 725(1996) : 11-22.
- Fujikawa, D.G., Shinmei, S.S., and Cai, B. Kainic acid-induced seizures produced necrotic, not apoptotic, neurons with internucleosomal DNA cleavage : implications for programmed cell mechanisms. *Neuroscience* 98(2000) : 41-53.
- Gambardella, A., Annesi, G., De Fusco, M., Patrignani, A., Aguglia, U., Annesi, F., Pasqua, A.A., Spadafora, P., Oliveri, R.L., Valentino, P., Zappia, M., Ballabio, A., Casari, G., and Quattrone, A. A new locus for autosomal dominant nocturnal frontal lobe epilepsy maps to chromosome 1. *Neurology* 55(2000) : 1467-1471.
- Gilbert, J.C., and Sawas, A.H. ATPase activities and lipid peroxidation in rat cerebral cortex synaptosomes. *Arch. Int. Pharmacodyn. Ther.* 263(1983) : 189-196.
- Green, D.R., and Reed, J.C. Mitochondria and apoptosis. *Science* 281(1988) : 1309-1310.
- Grillner, S., Ekeberg, O., Manira, A.E.I., Lansner, A., Parker, D., Tegner, J., and Wallen, P. Intrinsic function of a neuronal network-a vertebrate central pattern generator. *Brain Res. Rev.* 26(1998) : 184-197.
- Gupta, R.C., and Dettbarn, W-D. Prevention of kainic acid seizures-induced changes in levels of nitric oxide and high-energy phosphates by 7-nitroindazole in rat brain regions. *Brain Res.* 981(2003) : 184-192.
- Haerer, A.F., Anderson D.W., and Schoenberg, B.S. Prevalence and clinical features of epilepsy in a biracial United States population. *Epilepsia* 27(1)(1986) : 66-75.

- Hamberger, A.C., Chiang, G.H., Nylen, E.S., Scheff, S.W., and Cotman, C.W. Glutamate as a CNS transmitter I, evaluation of glucose and glutamine as precursors for the synthesis of preferentially released glutamate. *Brain Res.* 168(1979) : 513-530.
- Hamilton, S.E., Loose, M.D., and Qi, M. Disruption of the m1 receptor gene ablates muscarinic receptor-dependent M current regulation and seizure activity in mice. *Proc. Natl. Acad. Sci. USA* 94(1997) : 13311-13316.
- Haskew, R.E., Mongin, A.A., and Kimelberg, H.K. Peroxynitrite enhances astrocytic volume-sensitive excitatory amino acid release via a src tyrosine kinase-dependent mechanism. *J. Neurochem.* 82(2002) : 903-912.
- Hayasaka, K., Takahashi, I., Kobayashi, Y., Iinuma, K., Narisawa, K., and Tada, K. Effects of valproate on biogenesis and function of liver mitochondria. *Neurology* 36(3)(1986) : 351-356.
- Heinemann, U., Buchheim, K., Gabriel, S., Kann, O., Kovacs, R., and Schuchmann, S. Coupling of electrical and metabolic activity during epileptiform discharges. *Epilepsia* 43(2002) : 168-173.
- Henry, T.R. Therapeutic mechanisms of vagus nerve stimulation. *Neurology* 59(Suppl.4)(2002) : S3-S14.
- Hirao, T., Morimoto, K., Yamamoto, Y., Watanabe, T., Sato, H., Sato, K., Sato, S., Yamada, N., Tanaka, K., and Suwaki, H. Time-dependent and regional expression of GABA transporter mRNAs following amygdala-kindled seizures in rats. *Mol. Brain Res.* 54(1998) : 49-55.
- Hirose, S., Okada, M., Yamakawa, K., Sugawara, T., Fukuma, G., Ito, M., Kaneko, S., and Mitsudome, A. Genetic abnormalities underlying familial epilepsy syndromes. *Brain Dev.* 24(2002) : 211-222.
- Holmes, G.L. Seizure-induced neuronal injury. *Neurology* 59(suppl.5)(2002) : S3-S6.

Jenthet, S. *Effect of N-(2-propylpentanoyl) urea on GABA_A receptor in acutely dissociated rat hippocampal neurones.* Master's Thesis, Interdepartment Program of Physiology, Graduate School, Chulalongkorn University, 2002.

Johannessen, C.U. Mechanisms of action of valproate : a commentary. *Neurochem. Int.* 37(2000) : 103-110.

Johnson, J.W., and Ascher, P. Glycine potentiates the NMDA response in cultured mouse brain neurons. *Nature* 325(1987) : 529-531.

Kaura, S., Bradford, H.F., Young, A.M.J., Croucher, M.J., and Hughes, P.D. Effect of amygdaloid kindling on the content and release of amino acids from the amygdaloid complex : in vivo and in vitro studies. *J. Neurochem.* 65(1995) : 1240-1249.

Kleckner, N.W., and Dingledine, R. Requirement for glycine in activation of NMDA receptors expressed in *Xenopus* oocytes. *Science* 214(1988) : 843-848.

Khan, G.M., Smolders, I., Lindekens, H., Manil, J., Ebinger, G., Michotte, Y. Effects of diazepam on extracellular brain neurotransmitters in pilocarpine-induced seizures in rats. *Eur. J. Pharmacol.* 373(1999) : 153-161.

Khongsombat, O. *Effect of N(2'-propylpentanoyl)-2-pyrrolidinone and N-(2-propylpentanoyl) urea on neurons of cerebral cortex and cerebellar Purkinje cells in rats.* Master's Thesis, Interdepartment Program of Physiology, Graduate School, Chulalongkorn University, 1997.

Kramer, G. Epilepsy in the elderly : some clinical and pharmacotherapeutic aspects, *Epilepsia* 42(Suppl.3)(2001) : 55-59.

Kromer, G., and Reed, J.C. Mitochondrial control of cell death. *Nat. Med.* 6(2000) : 513-519.

Kudin, A.P., Kudina, T.A., Seyfried, J., Vielhaber, S., Beck, H., Elger, C.E., and Kunz, W. Seizure-dependent modulation of mitochondrial oxidative phosphorylation in rat hippocampus. *Eur. J. Neurosci.* 15(2002) : 1105-1114.

- Kunz, W.S., Kudin, A.P., and Vielhaber, S. Mitochondrial complex I deficiency in the epileptic focus of patients with temporal lobe of epilepsy. *Ann. Neurol.* 48(2000) : 766–773.
- Kwan, P., Sills, G.J., and Brodie, M.J. The mechanisms of action of commonly used antiepileptic drugs. *Pharmacol. Ther.* 90(2001) : 21–34.
- Kwan, P., and Brodie, M.J. Early identification of refractory epilepsy. *N. Eng. J. Med.* 342(2000) : 314–319.
- Lallement, G., Carpentier, P., Collet, A., Pernot-Marino, I., Baubichon, D., Blanchet, G. Effects of soman-induced seizures on different extracellular amino acid levels and on glutamate uptake in rat hippocampus. *Brain Res.* 563(1991) : 234-240.
- Laube, B., Hirozaku, H., Sturgess, M., Betz, H. and Kuhse, J. Molecular determinants of agonist discrimination by NMDA receptor subunits : analysis of the glutamate binding site on the NR2B subunit. *Neuron* 18(3)(1997) : 493–503.
- Lee, M-C., Kim, G-M., Woo, Y-J., Kim, M-K., Kim, J-H., Nam, S-C., Suh, J-J., Chung, W-K., Lee, J-S., Kim, H-I., Choi, H-Y., and Kim, S.U. Pathogenic significance of neuronal migration disorders in temporal lobe epilepsy. *Hum. Pathol.* 32(2001) : 643–648.
- Legendre, P. The glycinergic inhibitory synapse. *Cell. Mol. Life Sci.* 58(2001) : 760–793.
- Lehmann, A., Hagberg, H., Jacobson, I., and Hamberger, A. Effects of status epilepticus on extracellular amino acids in the hippocampus. *Brain Res.* 359(1985) : 147-151.
- Leite, J.P., Garcia-Cairasco, N., and Cavalheiro, E.A. New insights from the use of pilocarpine and kainate models. *Epilepsy Res.* 50(2002) : 93-103.
- Lemos, T., and Cavalheiro, E.A. Suppression of pilocarpine-induced status epilepticus and the late development of epilepsy in rats. *Exp. Brain Res.* 102(3)(1995) : 423-428.
- Liang, L.P., Ho, Y.S., and Patel, M. Mitochondrial superoxide production in kainate-induced hippocampal damage. *Neuroscience* 101(2000) : 563–570.

- Lipicky, R.J., Gilbert, D.L., and Stillman, I.M. Diphenylhydantoin inhibition of sodium conductance in squid giant axon. *Proc. Natl. Acad. Sci. USA* 69(1972) : 1758–1760.
- Litchfield, J.T., and Wilcoxon, F. A simplified method of evaluating dose-effect experiments. *J. Pharmacol. Exp. Ther.* 96(1949) : 99-113.
- Logothetis, J., and Bovis, M. Free amino acid changes in the spinal fluid of patients with CNS degeneration and chronic epileptic disorders. *World Neurol.* 3(1962) : 366-414.
- Lombardo, A.J., Kuzniecky, R., Powers, R.E., and Brown, G.B. Altered brain sodium channel transcript levels in human epilepsy. *Mol. Brain Res.* 35(1996) : 84–90.
- Loscher, W. Animal models of epilepsy for the development of antiepileptogenic and disease-modifying drugs : A comparison of the pharmacology of kindling and post-status epilepticus models of temporal lobe epilepsy *Epilepsy Res.* 50(2002) : 105-123.
- Lowry, O.H., Rosebrough, N.J., Farr, A.L., and Randell, R.J. Protein measurement with Folin phenol reagent. *J. Biol. Chem.* 193(1951) : 265-275.
- Mantegazza, M., Franceschetti, S., and Avanzini, G. Anemone toxin (ATXII) induced increase in persistent sodium current : effects on the firing properties of rat neocortical pyramidal neurones. *J. Physiol.* 507(1998) : 105–116.
- Maragakis, N.J., and Rothstein, J.D. Glutamate transporters in neurologic disease. *Arch. Neurol.* 58(2001) : 365–370.
- Maslansky, J.A., Powelt, R., and Diemengiant, C. Assessment of the muscarinic receptor subtype involved in pilocarpine-induced seizure in mice. *Neurosci. Lett.* 168 (1994) : 225-228.
- Matherne, G.W., Mendoza, D., Lozada, A., Pretorius, J.K., Desnes, Y., Danbolt, N.C., Nelson, N., Leite, J.P., Chimelli, L., Born, D.E., Sakamoto, A.C., Assirati, J.A., Fried, I., Peacock, W.J., Ojemann, G.A., Adelson, P.D. Hippocampal GABA and glutamate transporter immunoreactivity in patients with temporal lobe epilepsy. *Neurology*

52(1999) : 453–472.

Matsumoto, H., and Ajmone-Marsan, C. Cortical cellular phenomena in experimental epilepsy, Interictal manifestations. *Exp. Neurol.* 9(1964) : 286–304.

McAuley, J.W., Biederman, T.S., Smith, J.C., and Moore, J.L. Newer Therapies in the drug treatment of epilepsy. *Ann. Pharmacother.* 36(1)(2002) : 119-129.

McCabe, P.H. New anti-epileptic drugs for the 21st century. *Expert. Opin. Pharmacother.* 1(2000) : 633–674.

McNamara, J.O. Emerging insights into the genesis of epilepsy. *Nature* 399(1999) : A15-A22.

McNamara, J.O. Drugs effective in the therapy of the epilepsies. In J.G. Hardman, and L.E. Limbird (eds.). *Goodman & Gilman's the pharmacological basis of therapeutics*, pp.521-547. U.S.A. : McGraw-Hill, 2001.

Meldrum, B.S. GABAergic mechanisms in the pathogenesis and treatment of epilepsy. *Br. J. Pharmacol.* 27(suppl.1)(1989) : 3-11.

Meldrum, B.S., and Chapman, A.G. Excitatory amino acid receptors and antiepileptic drug development. *Adv. Neurol.* 79(1999) : 965–978.

Mello, L.E., Cavalheiro, E.A., Tan, A.M., Kupfer, W.R., Pretorius, J.K., Babb,m T.L., and Finch, D.M. Circuit mechanisms of seizures in the pilocarpine model of chronic epilepsy : cell loss and mossy fiber sprouting. *Epilepsia* 34(1993) : 985-995.

Men, S., Lee, D.H., Barron, J.R., and Munoz, D.G. Selective neuronal necrosis associated with status epilepticus : MR findings. *Am. J. Neuroradiol.* 21(2000) : 1837–1840.

Millan, M.H., Chapman, A.G., and Meldrum, B.S. Extracellular amino acid levels in hippocampus during pilocarpine-induced seizures. *Epilepsy Res.* 14(1993) : 139-148.

- Minamoto, Y., Itano, T., Tokuda, M., Matsui, H., Janjua, N.A., Hosokawa, K., Okada, Y., Murakami, T.H., Negi, T. and Hatase, O. In vivo microdialysis of amino acid neurotransmitters in the hippocampus in amygdaloid kindled rat. *Brain Res.* 573 (1992): 345–348.
- Morimoto, K., Sato, H., Sato, K., Sato, S., and Yamada, N. BW1003C87, phenytoin and carbamazepine elevate seizure threshold in the rat amygdala-kindling model of epilepsy. *Eur. J. Pharmacol.* 339(1997) : 11–15.
- Mosewich, R.K., So, E.L., and O'Brien, T.J. Factors predictive of the outcome of frontal lobe epilepsy surgery *Epilepsia* 41(2000) : 843–849.
- Mulder, A.H., and Snyder, S.H. Potassium-induced release of amino acids from cerebral cortex and spinal cord slices of the rat. *Brain Res.* 76(1974) : 297-308.
- Mulley, J.C., Scheffer, I.E., Petrou, S., and Berkovic, S.F. Channelopathies as a genetic cause of epilepsy. *Curr. Opin. Neurol.* 16(2003) : 171-176.
- Najm, I., Ying, Z., and Janigro, D. Mechanisms of epileptogenesis. *Neurol. Clin.* 19(2)(2001) : 237-50.
- Nanri, M., Yamamoto, J., Miyake, H., and Watanabe, H. Protective effect of GTS-21, a novel nicotinic receptor agonist, on delayed neuronal death induced by ischemia in gerbils. *Jpn. J. Pharmacol.* 76 (1998) : 23-29.
- Ni, J-W., Matsumoto, K., Li, H.B., Murakami, Y., and Watanabe, H. Neuronal damage and decrease of central acetylcholine level following permanent occlusion of bilateral common carotid arteries in rat. *Brain Res.* 673(1995) : 290-296.
- Ng, S.K., Hauser, W.A., Brust, J.C., and Susser, M. Hypertension and the risk of new-onset unprovoked seizures. *Neurology* 43(1993) : 425–428.
- O'Shea, R.D. Roles and regulation of glutamate transporters in the central nervous system. *Clin. Exp. Pharmacol. Physiol.* 29(2002) : 1018–1023.

Ortiz, J.G., Moshé, S.L., Sperber, E.F., Velisek, L., Ferchmin, P., and Claudio, O.I. Plasticity of excitatory amino acid transporters in experimental epilepsy. *Epilepsia* 41(2000) : 104–110.

Oxbury, J.M., and Whitty, W.M. Causes and consequences of status epilepticus in adults. *Brain* 94(1971) : 733–744.

Patchamart, W. Hepatotoxicity of N-(2-propylpentanoyl) urea in rats. Master's Thesis, Interdepartment Program of Pharmacology, Graduate School, Chulalongkorn University, 1996.

Patel, M. Mitochondrial dysfunction and oxidative stress : cause and consequence of epileptic seizures. *Free. Rad. Biol. Med.* 37(2004) : 1951-1962.

Patel, M., Liang, L.P., and Roberts, L.J. Enhanced hippocampal F2-isoprostane formation following kainate-induced seizures. *J. Neurochem.* 79(2001) : 1065–1070.

Paxinos, G., and Watson, C. *The rat brain in stereotaxic coordinates*, 2nd ed. San Diego : Academic Press, 1986.

Pena, F., and Tapia, R. Seizures and neurodegeneration induced by 4-aminopyridine in rat hippocampus in vivo : role of glutamate-and GABA-mediated neurotransmission and of ion channels. *Neuroscience* 101(2000) : 547–561.

Perry, T.L., and Hansen, S. Amino acid abnormalities in epileptogenic foci. *Neurology* 37(7)(1981) : 872-876.

Perucca, E. Pharmacological and therapeutic properties of valproate : a summary after 35 years of clinical experience. *CNS Drugs* 16(10)(2002) : 695-714.

Phillips, H.A., Scheffer, I.E., Berkovic, S.F., Hollway, G.E., Sutherland, G.R., and Mulley, J.C. Localization of a gene for autosomal dominant nocturnal frontal lobe epilepsy to chromosome 20q13.2. *Nat. Genet.* 10(1995) : 117–118.

Plum, M.C. Free amino acid levels in CSF of normal humans and their variation in cases of epilepsy and Spielmeyer-Vogt-Batten disease. *J. Neurochem.* 23(1974) : 595-600.

Ponsup, P. Effects of *N-(2-propylpentanoyl) urea (VPU)* on NMDA receptors subtype *NR1A/NR2B* expressed in *Xenopus laevis* oocytes. Master's Thesis, Chulalongkorn University, 2003.

Priel, M., and Albuquerque, E.X. Short-term effects of pilocarpine on rat hippocampal neurons in culture. *Epilepsia* 43(2002) : 40-46.

Prince, D.A. Physiological mechanisms of focal epileptogenesis. *Epilepsia* 26(suppl.)(1985) : S3-S14.

Puranam, R.S., and McNamara, J.O. Seizure disorders in mutant mice : relevance to human epilepsies. *Curr. Opin. Neurobiol.* 9(1999) : 281-287.

Racine, R.J. Modification of seizure activity by electrical stimulation II motor seizures. *Electroencephalogr. Clin. Neurophysiol.* 32(1972) : 295-299.

Radhakrishnan, K., So, E.L., and Silbert, P.L. Predictors of outcome of anterior temporal lobectomy for intractable epilepsy : a multivariate study. *Neurology* 51(1998) : 465-471.

Ragsdale, D.S., McPhee, J.C., Scheuer, T., and Catterall, W.A. Common molecular determinants of local anaesthetic, antiarrhythmic and anticonvulsant block of voltage-gated Na⁺ channels. *Proc. Natl. Acad. Sci. USA* 93(1996) : 9270-9275.

Rakic, P. Molecular and cellular mechanisms of neuronal migration : relevance to cortical epilepsies. *Adv. Neurol.* 84(2000) : 1-14.

Regesta, G., and Tanganelli, P. Clinical aspects and biological bases of drug-resistant epilepsies. *Epilepsy Res.* 34(1999) : 109-122.

Rosebrough, N.J., Farr, A.L., and Randell, R.J. Protein measurement with folin phenol reagent. *J. Biol. Chem.* 193(1951) : 265-275.

Rosenthal, R.E., Hamud, F., Fiskum, G., Varghese, P.J., and Sharpe, S. Cerebral ischemia and reperfusion : prevention of brain mitochondria and injury by lidoflazine. *J. Cereb. Blood Flow Metab.* 7(1987) : 752-758.

Rumbach, L., Warter, J.M., Rendon, A., Marescaux, C., Micheletti, G., and Waksman, A.

Inhibition of oxidative phosphorylation in hepatic and cerebral mitochondria of sodium valproate-treated rats. *J. Neurol. Sci.* 61(3)(1983) : 417-423.

Saisorn, B., Patarapanich, C., and Janwitayanuchit, W. Synthesis of monouric analogues of valproic acid. *Thai J. Pharm. Sci.* 16(1992) : 145-150.

Sato, K., Iwai, M., Zhang, W.R., Kamada, H., Ohta, K., Omori, N., Nagano, I., Shoji, M., and Abe, K. Highly polysialylated neural cell adhesion molecule (PSA-NCAM) positive cells are increased and change localization in rat hippocampus by exposure to repeated kindled seizures. *Acta. Neurochir. Suppl.* 86(2003) : 575-579.

Sawas, A.H., and Gilbert, J.C. Lipid peroxidation as a possible mechanism for the neurotoxic and nephrotoxic effects of a combination of lithium carbonate and haloperidol. *Arch. Int. Pharmacodyn. Ther.* 276(2)(1985) : 301-312.

Sayin, U., Osting, S., Hagen, J., Rutecki, P. and Sutula, T. Spontaneous seizures and loss of axo-axonic and axo-somatic inhibition induced by repeated brief seizures in kindled rats. *J. Neurosci.* 23(2003) : 2759-2768.

Schuchmann, S., Buchheim, K., Meierkord, H., and Heinemann, U. A relative energy failure is associated with low Mg⁺⁺ but not with 4-aminopyridine induced seizure-like events in entorhinal cortex. *J. Neurophysiol.* 81(1999) : 399-402.

Schulz, J.N., Henshaw, D.R., Siwek, D., Jenkins, B.G., Ferrante, R.J., Cipolloni, P.B., Kowall, N.W., Rosen, B.R., and Beal, M.F. Involvement of free radicals in excitotoxicity *in vivo*. *J. Neurochem.* 64(1995) : 2239-2247.

Seal, R.P., and Amara, S.G. Excitatory amino acid transporters: a family in flux. *Annu. Rev. Pharmacol. Toxicol.* 39(1999) : 431-456.

Sherwin, A.L. Neuroactive amino acids in focally epileptic human brain. *Neurochem. Res.* 24(1999) : 1387-1395.

- Shigeri, Y., Seal, R.P., and Shimamoto, K. Molecular pharmacology of glutamate transporters, EAATs and VGLUTs. *Brain Res . Brain Res. Rev.* 45(3)(2004) : 250-265.
- Shorvon, S. D. *Handbook of epilepsy treatment*. Oxford : Malden, MA, Blackwell Science, 2000.
- Sloviter, R.S., and Dempster, D.W. 'Epileptic' brain damage is replicated qualitatively in the rat hippocampus by central injection of glutamate or aspartate but not by GABA or acetylcholine. *Brain Res. Bull.* 15(1)(1985) : 39-60.
- Smolders, I., Belle, K.V., Ebinger, G., and Michotte, Y. Hippocampal and cerebellar extracellular amino acids during pilocarpine-induced seizures in freely moving rats. *Eur. J. Pharmacol.* 319(1997) : 21-29.
- Soderpalm, B. Anticonvulsants : aspects of their mechanisms of action. *Eur. J. Pain* 6(suppl.A)(2002) : 3-9.
- Sooksawate, T. *Anticonvulsant effects of N-(2-propylpentanoyl) urea*. Master's Thesis, Interdepartment Program of Pharmacology, Graduate School, Chulalongkorn University, 1995.
- Sordahl, L.A., Johnson, C., Blailock, Z.R., and Schwartz, A. The mitochondrion. In A. Schwartz (ed.). *Methods in Pharmacology*, New York : Merdith Corporation, 1971.
- Steinlein, O., Mulley, J.C., Propping, P., Wallace, R.H., Phillips, H.A., Sutherland G.R., Scheffer, I.E., and Berkovic S.F. A missense mutation in the neuronal nicotinic acetylcholine receptor α 4 subunit is associated with autosomal dominant nocturnal frontal lobe epilepsy. *Nat. Genet.* 11(1995a) : 201-203.
- Steinlein, O., Schuster, V., Fischer, C., and Haussler, M. Benign familial neonatal convulsions: confirmation of genetic heterogeneity and further evidence for a second locus on chromosome 8q. *Human Genet.* 95(1995b) : 25-29.

- Straub, H., Köhling, R., Frieler, A., Grigat, M. and Speckmann, E.J. Contribution of L-type calcium channel to epileptiform activity in hippocampal and neocortical slices of guinea-pigs. *Neuroscience* 95(2000) : 63–72.
- Sutula, T.P., Hagen, J., and Pitkanen, A. Do epileptic seizures damage the brain?. *Curr. Opin. Neurol.* 16(2003) :189–195.
- Tanaka, K., Watase, K., Manabe, T., Yamada, K., Watanabe, M., Takahashi, K., Iwama, H., Nishikawa, T., Ichihara, N., Kikuchi, T., Okuyama, S., Kawashima, N., Hori, S., Takimoto, M. and Wada, K. Epilepsy and exacerbation of brain injury in mice lacking the glutamate transporter GLT-1. *Science* 276(5319)(1997) : 1699–1702.
- Tantisira, B., Tantisira, M.H., Patarapanich, C, Sooksawate, T., and Chunngam, T. Preliminary evaluation of the anticonvulsant activity of a valproic acid analog : N-(2-propylpentanoyl) urea. *Res. Commun. Mol. Pathol. Pharmacol.* 97(2)(1997) : 151-164.
- Temkin, N.R. Antiepileptogenesis and seizure prevention trials with antiepileptic drugs : meta-analysis of controlled trials. *Epilepsia* 42(2001) : 515-524.
- Thom, M. Neuropathological findings in epilepsy. *Curr. Diagnos. Pathol.* 10(2004) : 93-105.
- Treiman, D.M. GABAergic mechanisms in epilepsy. *Epilepsia* 42(2001) : 8-17.
- Trist, D.G. Excitatory amino acid agonists and antagonists: pharmacology and therapeutic applications. *Pharmaceutica. Acta. Helvetiae.* 74(2000) : 221-229.
- Turski, W.A., Cavalheiro, E.A., Coimbra, C., da Penha Berzaghi, M., Ikonomidou-Turski, C., and Turski, L. Only certain antiepileptic drugs prevent seizures induced by pilocarpine. *Brain Res.* 434(1987) : 281-305.
- Turski, W.A., Cavalheiro, E.A., Schwarz, M., Czuczwar, S.J., Kleinrok, Z., and Turski, L. Limbic seizures produced by pilocarpine in rats: behavioral, electroencephalographic and neuropathological study. *Behav. Brain Res.* 9(1983) : 315-335.

- Turski, L., Ikonomidou, C., Turski, W.A., Bortolotto, Z.A., and Cavalheiro, E.A. Review: cholinergic mechanisms and epileptogenesis. The seizures induced by pilocarpine: a novel experimental model of intractable epilepsy. *Synapse* 3(1989) : 154-159.
- Ueda, Y. and Tsuru, N. Simultaneous monitoring of the seizure-related changes in extracellular glutamate and gamma-aminobutyric acid concentration in bilateral hippocampi following development of amygdaloid kindling. *Epilepsy Res.* 20(1995) : 213-219.
- Ueda, Y., Yokoyamaa, H., Niwa, R., Konaka, R., Ohya-Nishiguchi, H., and Kamada, H. Generation of lipid radicals in hippocampal extracellular space during kainic acid-induced seizures in rats. *Epilepsy Res.* 26(1997) : 329-333.
- Van den Pol, A.N. Obrieten, K., and Belousov, A. Glutamate hyperexcitability and seizure-like activity throughout the brain and spinal cord upon relief from chronic glutamate receptor blockade in culture. *Neuroscience* 74(1996) : 653-674.
- Wallace, R.H., Wang, D.W., Singh, R., Scheffer, I.E., George, A.L., Phillips, H.A., Saar, K., Reis, A., Johnson, E.W., Sutherland, G.R., Berkovic, S.F., and Mulley, J.C. Febrile seizures and generalized epilepsy associated with a mutation in the Na^+ -channel beta1 subunit gene SCN1B. *Nat. Genet.* 19(1998) : 366-370.
- Walton, N.Y., Gnawan, S., and Treiman, D.M. Brain amino acid concentration changes during status epilepticus induced by lithium and pilocarpine. *Exp. Neurol.* 108(1990) : 61-69.
- Wang, J.F., Azzam, J.E., and Young, L.T. Valproate inhibits oxidative damage to lipid and protein in primary cultured rat cerebrocortical cells. *Neuroscience* 116(2003) : 485-489.
- Watanabe, T., Morimoto, K., Hirao, T., Suwaki, H., Watase, K. and Tanaka, K., Amygdala-kindled and pentylenetetrazole-induced seizures in glutamate transporter GLAST-deficient mice. *Brain Res.* 845(1999) : 92-96.

- Werman, R., Davidoff, R.A., and Aprison, M.H. Inhibition of motoneurones by iontophoresis of glycine. *Nature* 214(1967) : 681–683.
- White, H.S. Clinical significance of animal seizure models and mechanism of action studies of potential antiepileptic drugs. *Epilepsia* 38(1997) : S9-S17.
- White, H.S. Comparative anticonvulsant and mechanistic profile of established and newer antiepileptic drugs. *Epilepsia* 40(suppl.5)(1999) : S2–S10.
- Wiebe, S., Blume, W.T., Girvin, J.P., and Eliasziw, M. Effectiveness and efficiency of surgery for temporal lobe epilepsy study group, a randomized, controlled trial of surgery for temporal-lobe epilepsy. *N. Engl. J. Med.* 345(2001) : 311-318.
- Williams, K., Chao, J., Kashiwagi, K., Masuko, T., and Igarashi, K. Activation of *N*-methyl-D-aspartate receptors by glycine : role of an aspartate residue in the M3–M4 loop of the NR1 subunit. *Mol. Pharmacol.* 50(1996) : 701–708.
- Wilson, C.L., Maidment, N.T., Shomer, M.H., Behnke, E.J., Ackerson, L., Fried, I. and Engel Jr., J., Comparison of seizure related amino acid release in human epileptic hippocampus versus a chronic, kainate rat model of hippocampal epilepsy. *Epilepsy Res.* 26(1996) : 245–254.
- Zabara, J. Peripheral control of hypersynchronous discharge in epilepsy. *Electroencephalogr. Clin. Neurophysiol.* 61(1985) : S162.
- Zhang, W.Q., Hudson, P.M., Sobotka, T.J., Hong, J.S. and Tilson, H.A., Extracellular concentrations of amino acid transmitters in ventral hippocampus during and after the development of kindling. *Brain Res.* 540 (1991) : 315–318.

BIOGRAPHY

NAME Miss Onrawee Khongsombat

DATE OF BIRTH 7th March 1970

PLACE OF BIRTH Singburi, Thailand

INSTRUCTIONS ATTENDED Mahidol University (1988-1991)

Bachelor of Nursing

Chulalongkorn University (1994-1997)

Master of Science (Physiology)

Chulalongkorn University (2000-2004)

Ph.D. candidate (Physiology)

POSITION & OFFICE 1992-1994 : Ramathipbodi Hospital

Thailand

1997-present: Department of Physiology

Faculty of Medical science

Naresuan University, Thailand

