

REFERENCES

- [1] Da Fonseca, M.G.; De Oliveira, M.M.; and Arakaki, L.N.H. Removal of cadmium, zinc, manganese and chromium cations from aqueous solution by a clay mineral. *J. Hazard. Mater.* B134 (2006): 288-292.
- [2] Manohar, D.M., Anoop Krishnan, K., and Anirudhan T.S. Removal of mercury(II) from aqueous solutions and chlor-alkali industry wastewater using 2-mercaptobenzimidazole-clay. *Water Res.* 36 (2002): 1609-1619.
- [3] Chiron, N., Guilet, R., and Deydier, E. Adsorption of Cu(II) and Pb(II) onto a grafted silica: isotherms and kinetic models. *Water Res.* 37 (2003): 3079-3086.
- [4] Mercier, L.; and Pinnavaia, T.J. Heavy metal ion adsorbents formed by the grafting of thiol functionality to mesoporous silica molecular sieves: factors affecting Hg²⁺ uptake. *Environ. Sci. Technol.* 32 (1998): 2749-2754.
- [5] Tonle, I.K.; Ngameni, E.; Njopwouo, D.; Carteret, C.; and Walcarius, A. Functionalization of natural smectite-type clays by grafting with organosilanes: physico-chemical characterization and application to mercury(II) uptake. *Phys. Chem. chem. Phys.* 5 (2003): 4951-4961.
- [6] Bourlinos, A.B.; Jiang, D.D.; and Giannelis, E.P. Clay-organosiloxane hybrids: a route to cross-linked clay particles and clay monoliths. *Chem. Mater.* 16 (2004): 2404-2410.
- [7] Erdemoglu, M.; Erdemoglu, S.; Sayilkan, F.; Akarsu, M.; Sener, S.; and Sayilkan, H. Organo-functional modified pyrophyllite: preparation, characterization and Pb(II) ion adsorption property. *Appl. Clay Sci.* 27 (2004): 41-52.
- [8] Krishna, B.S.; Murty, D.S.R.; and Jai Prakash, B.S. Surfactant-modified clay as adsorbent for chromate. *Appl. Clay Sci.* 20 (2001): 65-71.

- [9] Lin, S.H.; and Juang, R.S. Heavy metal removal from water by sorption using surfactant modified montmorillonite. J. Hazard. Mater. 92 (2002): 315 – 326.
- [10] Mercier, L.; and Detellier, C. Preparation, characterization, and application as heavy metals sorbents of covalently grafted thiol functionalities on the interlamellar surface of montmorillonite. Environ. Sci. Technol. 29 (1995): 1318-1323.
- [11] Wypych, F.; and Satyanarayana, K.G. Clay surface: Fundamentals and Applications. NewYork., USA., Elsevier Academic Press, Inc., 2004.
- [12] Bohn, H. L.; McNeal, B. L.; and O'Connor, G. A. Soil Chemistry. 3rd ed. NewYork., USA., JOHN WILEY & SONS, Inc., 2001.
- [13] Moore, D. M.; and Reynolds, R. C. X-Ray Diffraction and the Identification and Analysis of Clay Minerals. NewYork., USA., Oxford University Press, Inc., 1989.
- [14] Bradl, H. B. Heavy metals in the environment. NewYork., USA., Elsevier Academic Press, Inc., 2005.
- [15] BUILDING THE PHYLLOSILICATES[Online]. (n.d.). Available from: <http://pubpages.unh.edu/~harter/crystal.htm>[2008, March 15]
- [16] Michael, R.; Grace, S. T.; and Vishnu, K. The many ways of marking anionic clays. Proc. Indian Acad. Sci. (Chem. Sci.) 113 (2001): 671-680.
- [17] Baskaralingam, P.; Pulikesi, M.; Ramamurthi, V.; and Sivaneson, S. Equilibrium studies for the adsorption of acid dye onto modified hectorite. J. Hazard. Mater. B136 (2006): 989-992.
- [18] Jean, F. L.; and Georges, P. Acidity in pillared clays: origin and catalytic manifestations. Top. Catal. 4 (1997): 43-56.
- [19] Varma, R. S. Clay and clay-supported reagents in organic synthesis. Tetrahedron 58 (2002): 1235-1255.

- [20] Sayilkan, H.; Erdemoglu, S.; Sener, S.; Sayilkan, F.; Akarsu, M.; and Erdemoglu, M. Surface modification of pyrophyllite with amino silane coupling agent for the removal of 4-nitrophenol from aqueous solutions. J. Colloid. Interf. Sci. 275 (2004): 530-538.
- [21] Jiang, J. Q.; Cooper, C.; and Ouki, S. Comparison of modified montmorillonite adsorbents: Part I. Preparation, characterization and phenol adsorption. Chemosphere 47 (2002): 711-716.
- [22] Wu, P. X.; Lion, Z. W.; Zhang, H. F.; and Guo, J. G. Adsorption of phenol on ionorganic-organic pillared montmorillonite in polluted water. Environ. Int. 26 (2001): 401-406.
- [23] Liu, P.; and Guo, J. Polyarylamide grafted attapulgite (PAM-ATP) via surface-initiated atom transfer radical polymerization (SI-ATRP) for removal of Hg(II) ion and dyes. Colloids Surf., A 282-283 (2006): 498-503.
- [24] Salih, B.; Denizli, A.; Kavakli, C.; Say, R.; and Piskin, E. Adsorption of heavy metal ions onto dithizone-anchored poly (EGDMA-HEMA) microbeads. Talanta 46 (1998): 1205-1213.
- [25] Water Quality standards[Online]. Pollution control Department, Ministry of Natural Resources and Environment. Available from: http://www.pcd.go.th/Info_serv/reg_std_water.html[2008, March 15]
- [26] Zhang, F. S.; Nriagu, J. O.; and Itoh, H. Mercury removal from water using activated carbons derived from organic sewage sludge. Water Res. 39 (2005): 389-395.
- [27] Knocke, W. R.; and Hemphili, L. H. Mercury(II) sorption by waste rubber. Water Res. 15 (1981): 275-282.

- [28] Herrera. I.; Gardea-Torresdey. J. L.; Tiemann, K. J.; Peralta-Videa. J. R.; Armendariz. V.; and Parsons. J. G. Binding of silver(I) ions by Alfalfa biomass (*Medicago Sativa*): batch pH, time, temperature, and ionic strength studies. J. Hazard. Sub. Res. 4 (2003): 1-14.
- [29] Begum, S. Silver removal from aqueous solution by adsorption on concrete particles. Turk. J. Chem. 27 (2003): 609-617.
- [30] Ho, T. L. Hard and Soft Acids and Bases Principle in Organic Chemistry. NewYork., USA., Academic Press, Inc., 1977.
- [31] Eckenfelder, W. W. Application of adsorption to wastewater treatment. Tennessee., Enviro Press., 1981.
- [32] Goldberg, S.; Criscenti, L. J.; Turner, D. R.; Davis, J. A.; and Cantrell, K. J. Adsorption-desorption processes in subsurface reactive transport modeling. Vadose Zone J. 6 (2007): 407-435.
- [33] Mellah, A.; and Chegrouche, S. The removal of zinc from aqueous solutions by natural bentonite. Water Res. 31 (1997): 621-629.
- [34] Vengris, T.; Binkiene, R.; and Sveikauskaite, A. Nickel, copper and zinc removal from waste water by a modified clay sorbent. Appl. Clay Sci. 18 (2001): 183-190.
- [35] Park, M.; Shim, I. K.; Jung, E. Y.; and Choy, J. H. Modification of external surface of laponite by silane grafting. J. Phys. Chem. Solids 65 (2004): 499-501.
- [36] Shanmugharaj, A. M.; Rhee, K. Y.; and Ryu, S. H. Influence of dispersing medium on grafting of aminopropyltriethoxysilane in swelling clay materials. J. Colloid. Interf. Sci. 298 (2006): 854-859.
- [37] Evangelista, S. M.; DeOliveira, E.; Castro, G. R.; Zara, L. F.; and Prado, A. G. S. Hexagonal mesoporous silica modified with 2-mercaptothiazoline for removing mercury from water solution. Surf. Sci. 601 (2007): 2194-2202.

- [38] Mercier, L.; Pinnavaia, T. J. A functionalized porous clay heterostructure for heavy metal ion (Hg^{2+}) trapping. *Microporous Mesoporous Mater.* 20 (1998): 101-106.
- [39] Akhond, M.; Absalan, G.; Sheikhan, L.; Eskandari, M. M.; and Sharghi, H. Di (n-propyl) thiuram disulfide bonded on silica gel as a new sorbent for separation, preconcentration, and measurement of silver ion from aqueous samples. *Sep. Purif. Technol.* 52 (2006): 53-59.
- [40] Kavaklı, C.; Malcı, S.; Tuncel S. A.; and Salih, B. Selective adsorption and recovery of precious metal ions from geological samples by 1,5,9,13-tetrathiacyclohexadecane-3,11-diol anchored poly(p-CMS-DVB) microbeads. *React. Funct. Polym.* 66 (2006): 275-285.
- [41] Choudhury, S.B.; Ray, D.; and Chakravorty, A. Sulfur-ligated nickel oxidation states. Chemistry of a family of $\text{Ni}_z\text{S}_2\text{N}_4$ ($z = +2, +3, +4$) complexes incorporating hexadentate thioether-imine-oxime binding. *Inorg. Chem.* 30 (1991): 4354-4360.
- [42] Greenberg, A. E.; Clescerl, L. S.; and Eaton, A. D. STANDARD METHODS FOR THE EXAMINATION OF WATER AND WASTEWATER. 18th ed. NewYork., USA., EPS Group, Inc., 1992.
- [43] Spagnuoloa, M.; Marti'neza, C. E.; Jacobsona, A. R.; Baveyea, P.; McBridea, M. B.; and Newton, J. Coprecipitation of trace metal ions during the synthesis of hectorite. *Appl. Clay Sci.* 27 (2004): 129-140.
- [44] Madejova, J. FTIR techniques in clay mineral studies. *Vib. Spectrosc.* 31 (2003): 1-10.
- [45] Cariati, F.; Erre, L.; Micera, G.; Piu, P.; and Gessa, C. Water molecules and hydroxyl groups in montmorillonites as studied by near infrared spectroscopy. *Clays and Clay Miner.* 29 (1981): 157-159.

- [46] Carrado, K. A.; Xu, L.; Csencsits, R.; and Muntean, J. V. Use of organo- and alkoxysilanes in the synthesis of grafted and pristine clays. Chem. Mater. 13 (2001): 3766-3773.
- [47] Ma, J.; Xu, Y.; Fan, B.; and Liang, B. Preparation and characterization of sodium carboxymethylcellulose/poly(N-isopropylacrylamide)/clay semi-IPN nanocomposite hydrogels. Eur. Polym. J. 43 (2007): 2221-2228.
- [48] Silverstein, R. M.; Bassler, G. C.; and Morrill, T. C. Spectrometric Identification of Organic Compounds. New York., USA., John Wiley & Sons, Inc., 1991.
- [49] Tiwari, R. R.; Khilar, K. C.; and Natarajan, U. Synthesis and characterization of novel organo-montmorillonites. Appl. Clay Sci. 38 (2008): 203-208.
- [50] He, H.; Duchet, J.; Galy, J.; and Gerard, J. F. Grafting of swelling clay materials with 3-aminopropyltriethoxysilane. J. Colloid. Interf. Sci. 288 (2005): 171-176.
- [51] Pearson, R. G. Hard and Soft Acids and Bases. Stroudsville, PA., Dowden, Hutahinson and Ross., 1973.
- [52] Elastic network of lamellar particles: formation and rheological characterization [Online]. Majzik, A.; and Tombacz, E. Available from: http://www.rheofuture.de/papers2004/150404_hu_01.pdf[2008, April 9]
- [53] Baskaralingam, P.; Pulikesi, M.; Ramamurthi, V.; and Sivaneson, S. Modified hectorites and adsorption studies of a reactive dye. Appl. Clay Sci. 37 (2007): 207-214.
- [54] Shukla, A.; Zhang, Y. H.; Dubey, P.; Margrave, J. L.; and Shukla, S. S. The role of sawdust in the removal of unwanted materials from water. J. Hazard. Mater. B95 (2002): 137-152.
- [55] Lagergren, S. About the theory of so-called adsorption of soluble substances. K. Sven. Vetenskapsakad. Handl. 24 (1898): 1-39.

- [56] Aksu, Z.; and Tezer, S. Equilibrium and kinetic modelling of biosorption of Remazol Black B by *Rhizopus arrhizus* in a batch system: effect of temperature. Process Biochem. 36 (2000): 431-439.
- [57] Hameed, B. H. Equilibrium and kinetic studies of 2,4,6-trichlorophenol adsorption onto activated clay. Colloids Surf., A 307 (2007): 45-52.
- [58] Wan Ngah, W. S.; Kamari, A.; and Koay, Y. J. Equilibrium and kinetic studies of adsorption of copper(II) on chitosan and chitosan/PVA beads. Int. J. Biol. Macromol. 34 (2004): 155-161.
- [59] Ho, Y.S.; and McKay, G. The kinetics of sorption of divalent metal ions onto sphagnum moss peat. Water Res. 34 (2000): 735-742.
- [60] Hayes, K. F.; Papelis, C.; and Leckie, J. O. Modeling ionic strength effects on anion adsorption at hydrous oxide/solution interfaces. J. Colloid. Interf. Sci. 125 (1988): 717-726.
- [61] Singh, S. P.; Ma, L. Q.; and Hendry, M. J. Characterization of aqueous lead removal by phosphatic clay: Equilibrium and kinetic studies. J. Hazard. Mater. B136 (2006): 654-662.
- [62] Skoog, D. A.; West, D. M.; and James Holler, F. Fundamentals of Analytical Chemistry. 7th ed. USA., Tomson Learning, Inc., 1996.
- [63] Sarkar, D.; Essington, M.E.; and Misra, K. C. Adsorption of Mercury(II) by Kaolinite. Soil Sci. Soc. Am. J. 64 (2000): 1968-1975.
- [64] Dean, J. A. LANGE'S HANDBOOK OF CHEMISTRY. USA., McGraw-Hill, Inc., 1999.
- [65] Özcan, A. S.; and Özcan, A. Adsorption of acid dyes from aqueous solutions onto acid-activated bentonite. J. Colloid. Interf. Sci. 276 (2004): 39-46.
- [66] Vadivelan, V.; and Kumar, K. V. Equilibrium, kinetics, mechanism, and process design for the sorption of methylene blue onto rice husk. J. Colloid. Interf. Sci. 286 (2005): 90-100.

- [67] Filho, N. L. D.; Gushikem, Y.; Polito, W. L.; and Moreira, J. C. Sorption and preconcentration of metal ions in ethanol solution with a silica gel surface chemically modified with benzimidazole Ehirim. *Talanta* 42 (2004): 1625-1630.
- [68] Ngeontae, W.; Aeungmaitrepirom, W.; and Tuntulani, T. Chemically modified silica gel with aminothioamidoanthraquinone for solid phase extraction and preconcentration of Pb(II), Cu(II), Ni(II), Co(II) and Cd(II). *Talanta* 71 (2007): 1075-1082.
- [69] Bhattacharyya, K. G.; and Gupta, S. S. Pb(II) uptake by kaolinite and montmorillonite in aqueous medium: influence of acid activation of the clays. *Colloids Surf., A* 277 (2006): 191-200.
- [70] Alcantara, E. F. C.; Faria, E. A.; Rodrigues, D. V.; Evangelista, S. M.; DeOliveira, E.; Zara, L. F.; Rabelo, D.; and Prado, A. G. S. Modification of silica gel by attachment of 2-mercaptobenzimidazole for use in removing Hg(II) from aqueous media: A thermodynamic approach. *J. Colloid. Interf. Sci.* 311 (2007): 1-7.
- [71] Padilha, P. M.; Gomes, L. A. D.; Padilha, C. C. F.; Moreir, J. C.; and Dias Filho, N. L. Determination of metal ions in natural waters by flame-AAS after preconcentration on a 5-Amino-1,3,4-thiadiazole-2-thiol modified silica gel. *Anal. Lett.* 32 (1999): 1807.
- [72] Cestari, A. R.; Vieira, E. F. S.; Lopes, E. C. N.; and Da Silva, R. G. Kinetics and equilibrium parameters of Hg(II) adsorption on silica-dithizone. *J. Colloid. Interf. Sci.* 272 (2005): 271-276.
- [73] Pérez-Quintanilla, D.; Del Hierro, I.; Carrillo-Hermosilla, F.; Fajardo, M.; and Sierra, I. Adsorption of mercury ions by mercapto-functionalized amorphous silica. *Anal. Bioanal. Chem.* 384 (2006): 827.

- [74] Pérez-Quintanilla, D.; Del Hierro, I.; Fajardo, M.; and Sierra, I. Mesoporous silicafunctionalized with 2-mercaptopyridine: Synthesis, characterization and employment forHg(II) adsorption. Microp. Mesop. Mater. 89 (2006): 58-68.
- [75] Pérez-Quintanilla, D.; Del Hierro, I.; Fajardo, M.; and Sierra, I. 2-Mercapto-thiazoline modified mesoporous silica for mercury removal from aqueous media. J. Hazard. Mater. B134 (2006): 245-256.
- [76] Dias Filho, N. L.; Ribeiro do Carmo, D. Study of an organically modified clay: Selective adsorption of heavy metal ions and voltammetric determination of mercury(II). Talanta 68 (2006): 919-927.
- [77] Duc. M.; Gaboriaud. F.; and Thomas. F. Sensitivity of the acid-base properties of clays to the methods of preparation and measurement 1. Literature review. J. Colloid. Interf. Sci. 289 (2005): 139-147.
- [78] Sajidu, S. M. I.; Persson, I.; Masamba, W. R. L.; Henry, E. M. T.; and Kayambazinrhu, D. Removal of Cd^{2+} , Cr^{3+} , Cu^{2+} , Hg^{2+} , Pb^{2+} and Zn^{2+} cations and AsO_4^{3-} anions from aqueous solutions by mixed clay from Tundulu in Malawi and characterisation of the clay. Water SA 32 (2006): 519-526.
- [79] Ho, Y. S.; and Mckay, G. A comparison of chemisorptions kinetic models applied to pollutant removal on various sorbents. Trans. IChemE. 76 (1998): 332-340.

VITA

Mr. Theeradit Phothonthimongkol was born on July 19, 1983 in Bangkok, Thailand. He received his Bachelor degree of Science in Environmental Resource Chemistry, King Mongkut's Institute of Technology Ladkrabang in 2004. After that, he has been a graduate student at the Department of Chemistry, Chulalongkorn University and a member of Environmental Analysis Research Unit. He finished his postgraduate study with the Master degree of Science in 2007.