

## REFERENCES

- Anonymous. 1999. SPSS for windows Release 10.0, Standard version [Computer software], Chicago, SPSS Inc.
- Barrington, D.S. 1985. The present evolutionary and taxonomic status of the fern genus *Polystichum*: The 1984 botanical society of America pteridophyte section symposium. *Am. Fern J.* 75: 22-28.
- Bartolozzi, F., Warburton, M.L., Arulsekhar, S., and Gradziel T.M. 1998. Genetic characterization and relatedness among Californian almond cultivars and breeding lines detected by randomly amplified polymorphic DNA (RAPD) analysis. *J. Amer. Soc. Hort. Sci.* 123: 381-387.
- Benzing, D.H., 1990. *Vascular Epiphytes*. Cambridge University Press, Cambridge.
- Boonkerd, T. and H.P. Nooteboom. 2001. A new species of *Microsorum* (Polypodiaceae) from Thailand. *Blumea* 46: 581-583.
- Boonkerd, T., Pechsri, S. and Baum, B.R. 2005. A phenetic study of *Cassia* sensu lato (Leguminosae-Caesalpinoideae: Cassieae: Cassiinae) in Thailand. *Pl. Syst. Evol.* 252: 153-165.
- Boonkerd, T. 2006. A New Species of *Microsorum* (Polypodiaceae) from Thailand. *Blumea*. 51(1): 143-145.
- Bosman, M.T.M. 1991. *A monograph of the fern genus Microsorum (Polypodiaceae)*. Leiden, Rijksherbarium/Hortus Botanicus.
- Bosman, M.T.M., Hovenkamp, P.H. and Nooteboom, H.P. 1998. *Microsorum* Link. In: C. Kalkman, H.P. Nooteboom (eds.), *Flora Malesiana*. Ser. II, 3: 90-131. Leiden, Rijksherbarium/Hortus Botanicus.
- Botstein, D., White, R.L., Skolnick, M. and Davis, R. 1980. Construction of a genetic linkage map in man using restriction fragment length polymorphisms. *Am. J. Hum. Genet.* 32: 314-331.
- Budke, J.M., Hickey, R.J. and Heafner, K.D. 2005. Analysis of morphological and anatomical characteristics of *Isoetes* using *Isoetes tennesseensis*. *Brittonia* 57: 167-182.
- Buczkowska, K. 2004. The genus *Calypogeia* Raddi (Jungermanniales, Hepaticae) in Poland, biometrical analysis of morphological and anatomical variation. *Nova Hedwigia*. 78: 121-146.

- Casiva, P.V., Saidman, B.O., Vilardi, J.C. and Cialdella, A.M. 2002. First comparative phenetic studies of Argentinean species of *Acacia* (Fabaceae), using morphometric, isozymal, and RAPD approaches1. *Am. J. Bot.* 89:843-853.
- Chen, C.-C. and Chen, Y.-R. 2005. Study on laminar hydathodes of *Ficus formosana* (Moraceae) I. Morphology and ultrastructure. *Bot. Bull. Acad. Sin.* 46: 205-215
- Conceição, A.S., Queiroz, L.P., Lambert, S.M., Pereira, A.C.S. and Borba, E.L. 2008. Biosystematics of *Chamaecrista* sect. *Absus* subsect. *Baseophyllum* (Leguminosae-Caesalpinoideae) based on allozyme and morphometric analyses. *Pl. Syst. Evol.* 270: 183-207.
- Dallwitz, M.J., Paine, T.A. and Zurcher, E.J. 1993. User's Guide to the DELTA System: a General System for Processing Taxonomic Descriptions. 4th edition, available on the internet at <http://biodiversity.uno.edu/delta/>.
- Dellaporta, S.L., Wood, J. and Hicks, J.B. 1983. A plant DNA minipreparation: Version II. *Plant Mol. Biol. Rep.* 1: 19.
- Dice, L.R., 1945: Measurement of the amount of ecological association between species. *Ecology.* 26: 295-302.
- Dillon, W.R. and Goldstein, M. 1984. *Multivariate analysis. methods and applications.* New York, John Wiley and Sons.
- Dirlewanger, E., Duha, S., Viruel, M.A., and Saunier, R. 1998. Identification of peach varieties using molecular markers. *Acta. Hort.* 465: 69-77.
- Doyle, J.J. and Doyle J.L. 1987. A rapid DNA isolation procedure from small quantities of fresh leaf tissue. *Phytochem. Bull.* 19: 11-15.
- Felsenstein, J. 1985. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 39: 783-791.
- Geesink, R and Kornet, D.J. 1989. Speciation and Malesian Leguminosae. In L.B. Holm-Nielsen et al. *Tropical forests: botanical dynamics, speciation and diversity.* 135-151. London.
- Ghiselin, M.T. 1975. A radical solution to the species problem. *Syt. Zool.* 23: 536-544.
- Giussani, L.M. 2000. Phenetic similarity patterns of dioecious species of *Poa* from argentina and neighboring countries. *Ann. Missouri Bot. Gard.* 87:203-233.
- Gower, J.C. 1971. A general coefficient of similarity and some of its properties. *Biometrics.* 27: 857-871.

- Hanafy, M.F., Henry, T.W. and Domier, L.L. 1996. Use of random amplified polymorphic DNA (RAPD) for identification of *Gaeumannomyces* species. *Soil Bio and Biochem.*. 28: 703-710.
- Hasebe, M. et al. 1995. Fern phylogeny based on rbcL nucleotide sequences. *Am. Fern J.* 85, 134–181.
- Hilton, A.C., Banks, J.G., and Penn, P.C. 1997. Optimization of RAPD for fingerprinting *Salmonella*. *Appl. Microbiol.* 24: 243-248.
- Hogbin, P.M. and Crisp, M.D. 2003. Evolution of the coastal neospecies *Zieria prostrata* (Rutaceae) and its relationship to the *Zieria smithii* species complex. *Australian Syst. Bot.* 16: 515-525.
- Holmgren, P.K. and N.H. Holmgren. 1998 [continuously updated]. Index Herbariorum: A global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. <http://sweetgum.nybg.org/ih/>
- Holtum, R.E. 1949. *A revised flora of Malaya II: Ferns of Malaya*. Govt. Printer. Singapore.
- Holtum, R.E. 1954. (revised 1965). *Ferns of Malaya. A revised flora of Malaya II*. Incl. Appendix by I. Manton, Cytological notes on 100 species of Malayan ferns. Govt. Printer. Singapore.
- Hull, D.L. 1978. A matter of individuality. *Philosophy. Sci.* 45: 335-360.
- Janssen, T., Kreier, H.-P., and Schneider, H. 2007. Origin and diversification of African ferns with special emphasis on Polypodiaceae *Brittonia*, 59: 159-181.
- Kidyue, M., Boonkerd, T., Thaithong, O. and Seelanan, T. 2005. Numerical Taxonomy of the *Hoya parasitica* (Asclepiadaceae) complex in Thailand. *Nat. His. J. Chulalongkorn Univ.* 5:47-59.
- Kim, C., Shin, H. and Choi, H. 2003. A phenetic analysis of *Typha* in Korea and far east Russia. *Aquat. Bot.* 75: 33-43.
- Korner, J.S. and Livak, K.J. 1989. Mutation detection using nucleotide analogs that alter electrophoretic mobility. *Nucleic Acids Res.* 17: 7779-7783.
- Kreier, H.-P., and Schneider, H. 2006. Phylogeny and biogeography of the staghorn fern genus *Platycerium* (Polypodiaceae, Polypodiidae). *Am.J. Bot.* 93: 217–225.
- Lachenbruch, P.A. and Goldstein, M. 1979. Discriminant analysis. *Biometric.* 35: 69-85.
- Li, W. and Haufler, C.H. 1999. Genetic variation, breeding system, and pattern of diversification in Hawaiian *Polypodium* (Polypodiaceae). *Syst. Bot.* 24: 339-355.

- Lisek, A., Korbin, M. and Rozpara, E. 2006. Using simply generated RAPD markers to Distinguish between sweet cherry (*Prunus avium* L.) cultivars. *Fruit Ornam. Plant Res.* 14: 53-59
- Masood, A. and Abraham, G. 2006. Physiological response of *Azolla pinnata* plants to salinity stress. *Roum. Biotechnol. Lett.* 11: 2841-2844.
- Motley, T.F. and Morden, C.W., 2001. Utility of RAPD markers in evaluating the status of the Hawaiian tree fern *Cibotium xheleniae*. *Pacific Sci.* 55: 145-155.
- Nei, M. and Li, W.-H. 1979. Mathematical model for studying genetic variation in terms of restriction endonucleases. *Proc. Nat. Academy of Sci., USA.* 76: 5269-5273.
- Nilsson, J., Svensson, B., Ekelund, K. and Christiansson, A. 1998. A RAPD-PCR method for large-scale typing of *Bacillus cereus*. *Appl. Microbiol.* 27: 168-172.
- Nooteboom, H.P. 1997. The microsoroid ferns (Polypodiaceae). *Blumea*. 42: 261-395.
- Ortiz A., Renaud R., Calzada I., and Ritter E. 1997. Analysis of plum cultivars with RAPD markers. *J. Hort. Sci.* 72: 1-9.
- Pangua, E., Quinilla, L.G. and Pajaron, S., 2006. Taxonomic relevance of the gametophytic generation in a strictly rupicolous fern group: *Asplenium seelosii* s.l. *Bot. J. Linn. Soc.* 151: 375–386.
- Paris, C.A. 1989. Cryptic species, species delimitation and taxonomic practice in the homosporous ferns. *Am. Fern J.* 79: 46-54.
- Piesschaert, F., Jansen, S., Jaimes, I., Robbrecht, E. and Smets, E. 2001. Morphology, anatomy, and taxonomic position of *Pagameopsis* (Rubiaceae-Rubioideae). *Brittonia*. 53:490-504.
- Pryer, K.M., Smith, A.R. and Skog, J.E., 1995. Phylogenetic relationships of extant ferns based on evidence from morphology and rbcL sequences. *Am. Fern J.* 85: 202–282.
- Pryer, K.M. et al. 2001. Horsetails and ferns are a monophyletic group and the closest living relatives to seed plants. *Nature*. 409: 618–622.
- Regalado, L. and Sánchez, C. 2001. Spore morphology as a taxonomic tool in the delimitation of three *Asplenium* L. species complexes (Aspleniaceae: Pteridophyta) in Cuba. *Grana* 41: 107-113.
- Reudink, M.W., Joshua, P.S., Xu, B., Cunkelman, A. and Ronald, A.B. 2005. A Comparison of Physiological and Morphological Properties of Deciduous and Wintergreen Ferns in Southeastern Pennsylvania. *Am. Fern J.* 95: 45-56.

- Rohlf, F.J. 1998. *NTSYS-pc version 2.0k Numerical taxonomy and multivariate analysis system*. Setauket, N. Y.: Software.
- Rossello, J.A., Cebrian, M.C., and Mayol, M. 2002. Testing Taxonomic and Biogeographical Relationships in a Narrow Mediterranean Endemic Complex (*Hippocratea balearica*) using RAPD Markers. *Ann. Bot.* 89(3):321-327.
- Saiki, R.K. et al., 1988. Primer-directed enzymatic amplification of DNA with a thermostable DNA polymerase. *Science*. 239: 487-91.
- Salgado, E. 2004. Taxonomic and momenclature notes on Phillipine ferns II: three new names in *Asplenium*. *Edinburgh J. Bot.* 61: 185-195.
- Sano, R., Takamiya, M., Kurita, S., Ito, M. and Hasebe, M. 2000. *Diplazium subsinuatum* and *Di. tomitaroanum* should be Moved to *Deparia* According to Molecular, Morphological, and Cytological Characters. 113: 157-163.
- Schneider, H. et al. 2003. Unraveling the phylogeny of polygrammoid ferns (Polypodiaceae and Grammitidaceae): exploring aspects of the diversification of epiphytic plants. *Mol. Phylogen. Evol.* 31: 1041-1063.
- Seeprasert, T.. 2006. *Molecular taxonomic studies on Polygonum L. in Northeastern Thailand*. Master Degree Thesis. Department of Biologu, Khon Kaen University.
- Shimada T., Hayama H., Haji T., Yamaguchi M., and Yoshida, M. 1999. Genetic diversity of plums characterized by random amplified polymorphic DNA (RAPD) analysis. *Euphytica*. 109: 143-147.
- Small R.L. and Hickey R.J. 2001. Systematics of the Northern Andean *Isoëtes karstenii* Complex. *Am. Fern J.* 91: 41-69.
- Smith, A.R. and Hoshizaki, B.J., 2000. *Microsorum whiteheadii* (Polypodiaceae), an attractive new species from Sumatra. *Novon*. 10: 411-414.
- Sneath, P.H.A. and Sokal, R.R. 1973. *Numerical taxonomy: The principles and practice of numerical classification*. San Francisco, W. H. Freeman and Company.
- Starr, J.R. and Ford, B.A. 2001. The taxonomic and phylogenetic utility of vegetative anatomy and fruit epidermal silica bodies in *Carex* section *Phyllostachys* (Cyperaceae). *Can. J. Bot.* 79: 362–379.
- Swartz, L.M. and Brunsfeld, S.J. 2002. The morphological and genetic distinctness of *Botrychium minganense* and *B. crenulatum* as Assessed by Morphometric analysis and RAPD markers. *Am. Fern J.* 92: 249-269.
- Swofford, D.L. 1998. *PAUP\*: phylogenetic analysis using parsimony*. Sinauer Associates, Sunderland, Massachusetts, USA.

- Szwejkowski, J., Buczkowska, K., and Odrzykoski, I. J. 2005. *Conocephalum salebrosum* (Marchantiopsida, Conocephalaceae) – a new Holarctic liverwort species. *Pl. Syst. Evol.* 253:133-158
- Tagawa, M. and Iwatsuki, K. 1989. Polypodiaceae. In: T. Smitinand and K. Larsen (eds.), *Flora of Thailand*. 3(1): 486-580. Bangkok, Chutima Press.
- Tanksley, S. D., Young, N. D., Paterson, A. H., and Bonierbale, M. W. 1989. RFLP mapping in plant breeding – new tools for an old science. *Bio Technol.* 7: 257-264.
- Tenhuunen, J. D., Lange, O.L., Harley, P.C., Beyschlag, W. and Meyer, A. 1985. Limitations due to water stress on leaf net photosynthesis of *Quercus coccifera* in the Portuguese evergreen scrub. *Oecologia*. 67: 23-30.
- Thomson, J.A. 2000. Morphological and Genomic Diversity in the Genus *Pteridium* (Dennstaedtiaceae). *Annals of Botany* 85:77-99.
- Thomson, J.A., Chikuni, A.C., and McMaster, C.S. 2005. The taxonomic status and relationships of bracken ferns (*Pteridium*: Dennstaedtiaceae) from sub-Saharan Africa. *Bot J Linn Soc.* 148:311–321.
- Wang, T., SU, Y., LI, X., ZHENG, B., CHEN, G., and ZENG, G. 2004. Genetic structure and variation in the relict populations of *Alsophila spinulosa* from southern China based on RAPD markers and cpDNA atpB-rbcL sequence data. *Hereditas*. 140: 8-17.
- Warburton, M.L. and Bliss, F.A. 1996. Genetic diversity in peach (*Prunus persica* L. Batch) revealed by randomly amplified polymorphic DNA (RAPD) markers and compared to inbreeding coefficients. *J. Amer. Soc. Hort. Sci.* 121: 1012-1019.
- Ward, J. M. 1993. Systematics of New Zealand *Inuleae* (Compositae-Asteraceae) 1. A numerical phenetic study of the species of *Raoulia*. *New Zealand J. Bot.* 31: 21-28.
- Wiley, E.O. 1981. *Phylogenetics, The theory and practice of phylogenetic systematics*. John Wiley and Sons, New York.

## APPENDIX

## APPENDIX

**Table 4.4** Means and standard deviation of 23 quantitative characters of the eight clustering groupings of the *M. punctatum* (L.) Copel. complex.

Character	I		II		III		IV		V		VI		VII		VIII	
	mean	± SE	mean	± SE	mean	± SE	mean	± SE	mean	± SE	mean	± SE	mean	± SE	mean	± SE
LMW	37.63	3.33	16.30	2.34	72.64	26.81	78.10	11.75	68.93	10.83	39.33	10.58	96.53	16.45	58.94	21.12
LML	130.62	3.11	304.54	52.37	536.25	184.31	970.13	157.08	386.16	62.33	314.65	70.38	831.55	156.73	747.08	222.75
STL	40.68	2.31	34.19	1.05	138.66	6.54	19.23	1.71	64.84	4.44	62.34	4.61	0.47	0.12	101.8	7.53
STD	2.49	0.11	2.92	0.18	4.75	0.97	5.31	1.28	5.84	0.34	1.54	0.36	5.75	1.04	4.55	0.96
PDL	3.19	0.39	3.41	0.32	2.67	0.88	5.87	1.45	4.63	0.45	3.28	1.15	6.13	1.25	4.48	1.45
RHD	3.63	0.32	4.93	0.25	5.56	1.47	7.48	1.56	5.71	0.35	4.99	0.88	6.96	1.44	5.82	1.38
RDL	4	0.20	1.63	0.19	2.83	1.32	11.51	2.80	5.11	0.54	3.48	1.64	12.45	2.99	8.57	2.56
SCL	2.6	0.27	3.02	0.20	4.42	1.34	4.01	0.64	4.23	0.75	4.03	1.03	3.64	0.40	3.31	0.63
SCW	1.24	0.06	1.15	0.08	1.46	0.36	1.25	0.18	2.78	0.39	1.04	0.19	1.42	0.22	1.20	0.27
ALA	19.33	0.82	7.33	1.41	14.25	4.36	18.88	3.57	18.4	2.97	16.71	5.53	23.57	5.84	13.86	4.67
ALB	2.5	0.55	1.78	0.83	2.02	1.09	6.01	1.56	8.00	1.00	1.88	1.75	19.34	6.88	4.88	3.94
DSO	1.37	0.04	1.31	0.06	1.93	0.39	1.85	0.41	1.37	0.10	1.33	0.13	1.39	0.15	1.28	0.17
DSR	0.18	0.01	0.18	0.01	0.19	0.02	0.19	0.02	0.18	0.01	0.16	0.03	0.16	0.01	0.17	0.15
NAC	20.83	0.75	28.33	1.80	19.41	1.76	19.78	0.97	26.00	2.00	21.56	1.37	18.90	0.75	19.25	1.00
SPL	51.25	1.37	45.22	1.89	58.75	8.14	57.97	4.68	51.5	2.85	50.64	4.16	47.71	4.89	51.67	4.13
SPW	33.75	1.37	31.94	3.70	36.60	5.57	32.53	3.94	36.00	2.24	33.27	3.98	33.33	3.62	31.38	3.10
DSL	12	1.26	19.11	2.37	9.87	3.78	37.34	4.63	17.00	1.87	25.97	7.92	39.71	7.13	38.70	10.93
PLS	59.44	2.81	138.91	17.19	196.78	113.64	517.69	144.79	124.61	16.03	156.98	43.55	430.14	137.82	383.64	120.66
PHS	11.22	1.15	5.01	1.75	16.79	6.07	17.66	7.57	16.57	2.41	16.13	7.84	19.33	4.85	14.35	6.63
NSR	3.33	0.52	7.22	0.67	2.24	0.46	4.01	0.87	5.40	0.55	4.10	0.88	8.08	1.47	9.29	2.01
DBSV	7.1	0.64	16.22	0.63	10.03	2.36	8.66	1.68	11.64	1.75	9.05	1.87	13.27	1.95	13.45	2.84
APS	29.5	1.52	49.78	1.72	21.51	3.57	29.10	4.84	41.00	2.65	44.73	6.04	27.74	3.60	33.98	5.46
PAW	1.46	0.13	1.09	0.11	4.14	1.97	3.11	1.73	1.55	0.29	2.16	0.59	4.35	1.52	1.10	0.55

**Table 4.5** Summary of canonical discriminant function of eight clustering groups based on 23 quantitative characters.

Function	Eigenvalue	% of Variance	Cumulative %	Canonical Correlation	Wilks' Lambda	Chi-square	df	Sig.
1	43.628	77.6	77.6	.989	.000	6290.994	140	.000
2	7.141	12.7	90.3	.937	.005	3662.521	114	.000
3	2.761	4.9	95.2	.857	.041	2211.437	90	.000
4	1.469	2.6	97.8	.771	.154	1294.813	68	.000
5	.791	1.4	99.2	.665	.380	669.409	48	.000
6	.297	.5	99.8	.478	.681	266.016	30	.000
7	.133	.2	100.0	.342	.883	86.148	14	.000

**Table 4.6** Summary of canonical discriminant function of 21 taxa based on 23 quantitative characters.

Function	Eigenvalue	% of Variance	Cumulative %	Canonical Correlation	Wilks' Lambda	Chi-square	df	Sig.
1	46.351	75.4	75.4	.989	.000	7195.815	380	.000
2	7.616	12.4	87.8	.940	.001	4549.504	342	.000
3	3.267	5.3	93.1	.875	.011	3072.092	306	.000
4	1.656	2.7	95.8	.790	.048	2076.731	272	.000
5	.949	1.5	97.3	.698	.129	1406.509	240	.000
6	.686	1.1	98.4	.638	.251	948.607	210	.000
7	.368	.6	99.0	.519	.423	590.102	182	.000
8	.288	.5	99.5	.473	.579	374.949	156	.000
9	.108	.2	99.7	.312	.746	201.275	132	.000
10	.062	.1	99.8	.242	.826	130.810	110	.086
11	.044	.1	99.9	.206	.878	89.536	90	.494
12	.028	.0	99.9	.166	.916	59.853	72	.846
13	.019	.0	99.9	.136	.942	40.787	56	.937
14	.016	.0	100.0	.126	.960	28.017	42	.952
15	.011	.0	100.0	.104	.975	17.048	30	.972
16	.008	.0	100.0	.088	.986	9.582	20	.975
17	.004	.0	100.0	.061	.994	4.239	12	.979
18	.002	.0	100.0	.046	.998	1.647	6	.949
19	.000	.0	100.0	.016	1.000	.175	2	.916

**Table 4.7** Cross-validation summary with the linear discriminant function of the *M. punctatum* complex. ( I = *Microsorum siamense*, II = *M. thailandicum*, III = *M. membranaceum*, IV = *M. glossophyllum*, V = *M. steerei* var. *whiteheadii*, VI = *M. steerei* var. *steerei*, VII = *M. musifolium*, and VIII = *M. punctatum*

From taxa	Classified in taxa							
	I	II	III	IV	V	VI	VII	VIII
<i>M. siamense</i>	6	0	0	0	0	0	0	0
<i>M. thailandicum</i>	0	9	0	0	0	0	0	0
<i>M. membranaceum</i>	0	0	86	0	0	0	0	0
<i>M. glossophyllum</i>	0	0	0	68	0	0	0	0
<i>M. steerei</i> var. <i>whiteheadii</i>	0	0	0	0	5	0	0	0
<i>M. steerei</i> var. <i>steerei</i>	0	0	0	0	0	39	0	0
<i>M. musifolium</i>	0	0	0	0	0	0	48	0
<i>M. punctatum</i>	0	0	0	0	0	0	0	446
<b>Total correctly classified</b>	<b>100%</b>	<b>100%</b>	<b>100%</b>	<b>100%</b>	<b>100%</b>	<b>100%</b>	<b>100%</b>	<b>100%</b>

## BIOGRAPHY

Mr. Sahanat Petchsri was born on May 11, 1978, in Phatthalung Province. He was graduated in Biology from Faculty of Science, Prince of Songkla University in 1999. In 2002, he received his Master of Science in Botany from the Department of Botany, Faculty of Science, Chulalongkorn University, then continued his study in Biological Science Ph.D. Program, Faculty of Science, Chulalongkorn University from 2003-2008.