



## REFERENCES

- Ackard, M.D et.al. (2001). Health Care Information Sources for Adolescents: Age and Gender Differences on Use, Concerns, and Needs. **Journal of Adolescent Health**. 29(3),170 –176.
- Adolescent health and school health (1998): it's time to meet the challenge. (1998). **Health & Social Work**. 23(1), 3-7.
- Alfieri, T., Ruble, D. N., & Higgins, E. T. (1996). Gender stereotypes during adolescence: Developmental changes and the transition to junior high school. *Developmental Psychology*, 32, 1129-1137.
- Andrews University. (2002). Risk Behaviorus. Retrieved 20 September 2002, From,[http://www.andrews.edu/IPA/education/adolescent\\_health/risk\\_behaviour](http://www.andrews.edu/IPA/education/adolescent_health/risk_behaviour)
- Anjira Assavanonda. (2000, June, Sunday 4). **Bangkok post**. (newspaper).
- Anonymous.(1995). Health of young people [news].**World Health Forum** 16, 108–111.
- Aten, M.J., Siegel, D.M. and Roghmann. K.J. (1996). Use of health services by urban youth: a school-based survey to assess differences by grade level, gender, and risk behaviour. **Journal of Adolescent Health** 19, 258–266.
- Bailey and MacNally, K. (2003). Community partnerships. [Online]. Retrieved February 2003, from <http://www.partnerships.org.uk/guide/theory.html>.
- Baker, S. et al. (2000). **Sexual behavior among Thai college students**. The proceeding of The 2002 Thai National Symposium on Population Studies. Chulalongkorn University. Thailand.
- Bangkok Metropololitan Administration [BMA], Klong Toey District Office. (1999). **Klong Toey District Report 1999**. Bangkok: BMA.
- Bangkok Metropolitan Administration [BMA]. Department of Health. (2000). **Attitude and HIV risk behaviors surveillance survey**. BMA, Depart. of Health.
- Bangkok Metropolitan Administration [BMA] Department of Health. (2001).**Bangkok Metropolitan Administration report year 2001**. Bangkok: Department of Health
- Bangkok Metropolitan Administration.[BMA]. (2002). **Bangkok Metropolitan Administration report year 2002**. Bangkok: BMA.
- Bangkok Metropolitan Administration [BMA].( 2003). Medical Services and Health Development Project. Retrieved from [http://www.bma.go.th/bmaeng/body\\_project4.html](http://www.bma.go.th/bmaeng/body_project4.html)

- Bangkok Slum (2001). Bangkok Slum. Retrieved 18 May 2001, from [www.eapap.unep.org/reports/soe/bangkok\\_toc.pdf](http://www.eapap.unep.org/reports/soe/bangkok_toc.pdf).
- Barnes, M.P and Jenney, E. M. M. (2002). **Measuring Quality of Life**, 12, 467-480.
- Barkin, L.S.et al. (2003). Health care utilization among homeless adolescents and young adults. **Journal of Adolescent Health**. 32(4), 253- 256.
- Barton, J, and William, J.P.(2002). Adolescence. In: Detels, Roger. **Oxford Textbook of Public Health Vol. 3**, (pp.1623-1635). Oxford: Oxford University Press.
- Bearman, P. S. et al. (1999). Foregone Health Care Among Adolescents. *JAMA*, 282(23), 2227-2234.
- Beker, D. Z. (2002). Quality of Life Indicators: Adolescents' perceptions. From Dissertation Abstracts International.
- Bencha Yoddumnern. Attig. [et.al]. (1993). **Qualitative methods for population and health research**. Nakhon Pathom: Institute of Population and Social Research. Mahidol University.
- Bennett, A. (1993). **Do population studies need qualitative research? Qualitative methods for population and health research**. Nakhon Pathom: Institute for Population and Social Research, Mahidol University.
- Birmaher et al., (1996). Childhood and adolescent depression: a review of the past 10 years: Part I. **J. Am. Acad. Child Adolesc. Psych.** 35, 1427–1439.
- Bradford, R. Rutherford, L. D and John, A. (2002). Quality of life (QoL) in young people: ratings and factor structure of the QoL Profile – Adolescent version. **Journal of Adolescence**, 25, 261-274.
- Brindis, C. & Van Landeghem, K. (1999). **Adolescents and the State Children's Health Insurance Programme (CHIP): Healthy Options for Meeting the Needs of Adolescents**. San Francisco: Policy Information and Analysis Center for Middle Childhood and Adolescence & National Adolescent Health Information Center. University of California.
- Booth, L.M et. al. (2004). Access to health care among Australian A young people's perspectives and their sociodemographic distribution. **Journal of Adolescents Health**. 43(1), 97-103.
- Brooks, SJ. & Kutcher S. (2001) Diagnosis and measurement of adolescent depression: a review of commonly utilized instruments. **Journal of Child and Adolescent Psychopharmacology**. 11(4), 341-76.
- Carr., J.A & Higginson J. I. (2001). Measuring quality of life: Are quality of life measures patient centred?. **British Medical Journal**. 322, 1357-1360
- Centers for Disease Control and Prevention [CDC]. Mortality**. Retrieved 20 June 2002, from [www.cdc.gov/ncipc/default.htm](http://www.cdc.gov/ncipc/default.htm)

- Chai Podhisita and U. Pattaranich. (1995). **Youth in contemporary Thailand: Result from the family and youth survey**. Cited in Pimpawan Boonmongkon et al. (2000). Nakhon Prathom: Institute for population and social research, Mahidol University.
- Chanpen Chooprapawan, et.al. (1996). **The Thailand health status survey year 1987-1988**. Nonthaburi: Health Research Institute, MoPH. Thailand.
- Chanpen Chooprapawan, et.al. (2000). **The Thailand health status survey year**. Nonthaburi: Health Research Institute, MoPH. Thailand.
- Chavis, D.M.(1995). Building community capacity to prevent violence through coalitions and partnerships. **Journal of health care for the poor and underserved**, 6 (2), 234-245.
- Chitr Sitthi-amorn. (2000). **Summary of Asian Consultation Knowledge Management and Resourcing Public Health: Clarification of concepts and enabling national processes**. [Bangkok]: The College of Public Health, Chulalongkorn University. (Proceeding of Resourcing Public Health Workshop. Bangkok.)
- Chitr Sitthi-amorn. (2004). **Medicines as Emerging Technologies: Unprecedented Opportunities & Continuing Disparity**. [unpublished paper.]
- Chitr Sitthi-amorn and Ratana Somrongthong (2000). Multi-disciplinary research in public health. **J. of Research Methodology**. 13(3),1-21.
- Cheryl, P et al.(2000). The Minnesota Drug Abuse Resistance Education (DARE PLUS) Project: Creating community partnerships to prevent drug use and violence. **Journal of School Health**, 70(3), 84-89.
- Choe, M.K. et al. (2004). Substance use and premarital sex among adolescents in Indonesia, Nepal, the Philippines and Thailand. **Asia- Pacific Population Journal**.19(1), 5-26.
- Clark, J.P. ( 2000). Balance qualitative & quantitative methodology in health services research. **Retrieved 7 August 2004**, from <http://www.cehip.org>
- Devellis. F. R, (1991). **Scale Development: Theory and Applications**. New York: SAGE.
- Dew, T and E. Scott Huebner, E. S. (1994). Adolescents' perceived quality of life: An exploratory investigation. **Journal of School Psychology**, 32(2), 185-199.
- Duang Prateep Foundation (2000). **Duang Prateep Foundation Report 2000**. Bangkok: Foundation.
- Duang Prateep Foundation (2001). Duang Prateep Foundation. Retrieved 30 March 2001, from [www.dpf.or.th](http://www.dpf.or.th).
- Durex. (1999). **Global Sex Survey: Youth Perspective**. [S.l: S.n].

- Edman, J. L. (1996). Depressive symptom factors among Filipino American adolescents: An analysis of the Center for Epidemiologic Studies Depression Scale (CES-D). Paper presented at the conference for Pacific Islander & Asian American Biomedical Research, Honolulu, HI.
- Edwards, C.T. et al. (2001). Quality of life and health-risk behaviours among adolescents. **Journal of Adolescents Health**. 29(6), 426-435.
- Edward, LS. and Stern, FR. (1998) Building and Sustaining Community Partnerships for Teen Pregnancy Prevention. Retrieved 2 June 2002, from <http://www.aspe.hhs.gov/hsp/teenp/teenpreg/teenpreg.htm>.
- Executive Office for Weed and Seed, World Health Organization (WHO). (1996). OSAP; National Clearinghouse on Families and Youth unpublished Adolescent Health and Development. [S.l]: WHO.
- Family Planning and Population Division. Department of Health Ministry of Public Health [MOPH],. (1998). **Thailand National Family Planning Program**. Nonthaburi: Division.
- Fischhoff, B and Willis, H. (2002). Adolescent vulnerability: measurement and priority setting. **Journal of Adolescent Health**. 131(2), 58-75.
- Franklin, J.D. (2003). Adolescent Depression. Retrieved October 31, 2003 ,from [www.luphealth.com/health/adolescent\\_depression\\_inof.htm](http://www.luphealth.com/health/adolescent_depression_inof.htm) (Psychology Information Online).
- Family Planning Needs of and Services for the Underprivileged Urban: The Case of Bangkok. (1984). Proceedings of the Seminar on, January 26 – 28, Wongse Amataya Hotel, Chonburi.
- Fayers, M.P. and Machin D.( 2000). **Quality of life: Assessment, Analysis and Interpretation**. England: John Wiley & Son.LTD. England.
- Ford, Bearman, P.S. and Moody, J. (1999). Foregone health care among adolescents. **Journal of the American Medical Association**. 282, 2227–2234.
- Ford, C. A., Millstein, S.G., and Halpern-Felscher, B.L. (1997). Influence of physician confidentiality assurances on adolescents' willingness to disclose information and seek future health care. **Journal of the American Medical Association**. 278, 1029–1034.
- Freed, H.L et al. (1998). Determinants of Adolescents' Satisfaction With Health Care Providers and Intentions to Keep Follow-Up Appointments. **Journal of Adolescent Health**. 22(6), 475-479.
- Geenen, S.J., Powers, L.E., Sells, W. (2003) Understanding the role of health care providers during the transition of adolescents with disabilities and special health care needs. **Journal of Adolescents Health**. 32(3), 225-33.

- Gill T.M, and Feinstein A.R. ( 1994). A Critical appraisal of the Quality of Life Measurements. **Journal of the American Medical Association.** 272, 619-626.
- Gillifes, P. (1998). Effectiveness of alliances and partnerships for health promotion. Health promotion International. Alliances and partnerships for health promotion. Print in Great Britain. Oxford University Press.
- Gleeson, R. C., Robinson, B M, Neal D. R.( 2002) A review of teenagers' perceived needs and access to primary health care: implications for health services. **Journal of Primary Health Care Research and Development.** 3,184-193.
- Gonzales, C. et al. (1985). Adolescent health care: improving access by school-based service in Bangkok. [Online]. **Journal of Family Practice.** 21, 263-270. Retrieved from <http://db.jhuccp.org/dbtw-wpd/exec>
- Goodburn, E. Ross A, D. (2000). Young people's health in developing countries: a neglected problem and opportunity. **Health Policy and Planning;** 15(2),137-144.
- Goodman, E et.al. (2003). A multilevel analysis of the relation of socioeconomic status to adolescent depressive symptoms: does school context matter? **The Journal of Pediatrics,** 143(4), 451-456.
- Guzman, D. A. (1999). **Adolescent Sexual Health in Klong Toey Slum Community, Bangkok, Thailand.** Master's thesis of Public Health, University of Washington, Dept. of Health Services.
- Hallfors, D.D et. al., (2004). Adolescent depression and suicide risk Association with sex and drug behavior. **American Journal of Preventive Medicine.**27,(3), 224-231.
- Hardon, A. et.al ( 1995). Applied Health Research Manual Anthropology of Health and Health Care :A joint initiative of Mahidol University, University of Amsterdam. [Online]. Retrieved from <http://aspe.hhs.gov/hsp/teep/teenpreg/teenpreg.htm>
- Havisto, A. et al. (2004). Factors associated with depressive symptoms among 18-year-old boys: a prospective 10-year follow-up study **Journal of Affective Disorders,** 83(2/3),143. Retrieved August 25, 2004, from EBSCO host database.
- Health of young people** [news]. (1995). World Health Forum. 16: 108 –111.
- Hesketh, T., Ding, J Q., and Tomkins, M A.(2003) Health and health care-seeking behavior of adolescents in urban and rural China. **Journal of Adolescent Health.**33.(4), 271-274.
- Hi-class. (2000). 72 hours in Klong Toey Slum. (June 2000). **Hi-class.** p 82-86.(in Thai)
- Hiebert, B. (1998). A multi-site exploration of adolescent health –related needstudent and adult perceptions.[Online]. PhD dissertation. University of Calgary. Retrieved from **Dissertation Abstracts International Database.**
- Irwin, E.C, Duncan, M.P. (2002) Health Futures of Youth II: pathways to adolescent health, executive summary and overview. **Journal of Adolescent Health.** 31(6), 82-89.

- Khammapirad B., et. al. (1984). Family Planning Needs of and Services for the Underprivileged Urban: The Case of Bangkok: Proceedings of the Seminar on, January 26 – 28, Wongse Amataya Hotel, Chonburi.
- Khobkul Sudsawad. (1998). **Quality of life of the elderly in Nakhon Si Thammarat Province**. Master Thesis of Sociology. Faculty of Political Sciences. Chulalongkorn University, Bangkok.
- King, A. et. al. (1996). **The Health of Youth: a Cross- National Survey**. [Geneva: WHO]. WHO Regional Publication, European Series No. 69.
- King, and Keith A. (2001). Tri- level Suicide Prevention Covers it all. **Education Digest**. 67(1), 55-62.
- Kirby, D. (1997). **No Easy Answer: Research Findings on Programs to Reduce Teen Pregnancy**. [Online]. Washington, D.C.: the National Campaign to Reduce Teen Pregnancy. Retrieved from <http://www.aspe.hhs.gov/hsp/teenp/teenpreg/teenpreg.htm>.
- Klein, J.D. et al. ( 1992) Access to health care for adolescents: A position paper of the Society from Adolescent Medicines.[Online].**Journal of Adolescent Health**. 13:162-170. Retrieved August 25, 2004, from [http:// www. Adolescenthealth.org/access/html](http://www.Adolescenthealth.org/access/html)
- Klein, J.D. [et al.]. (1999). Access to medical care for adolescents: results from the 11997 Commonwealth Fund Survey of the Health of Adolescent Girls. **Journal of Adolescents Health**.25 (2), 120-130.
- Klein, J.D. (2000). Adolescents, health services, and access to care. **Journal of Adolescents Health**.27(5), 293-294.
- Kodjo,C., Auinger, P., and Ryan, S. (2002) Barriers to adolescents accessing mental health services. **J. of Adolescent Health**. 30(2), 101-102.
- Kovacs. M.(1981). Rating scale to assess depression in school aged children. **Acta Paedopsychiatr**. 46, 305–315.
- Lau, Joseph T. F. [et al.]. (2000). Studies on common illnesses and medical care utilization patterns of adolescents in Hong Kong. **Journal of Adolescents Health**. 27(6), 443-452.
- Lee, D. (2001). **Strategies for Enhancing Access to Medicines (SEAM) Conference 2001, Targeting Improved Access: Proceeding**, November 27-29, 2001. Washington DC.
- Levine, S.B and Coupey, S.M. (2003). Adolescent Substance Use, Sexual Behaviour, and Metropolitan Status: Is " urban" a risk factor?. **Journal of Adolescent Health**. 32:350-355
- Litt, IF (1998). Satisfaction with health care: The adolescents' perspective. **Journal Adolescent Health** 23, 59–60.

- Litt I.F. (Ed.). (2003). Focusing on focus groups. **Journal of Adolescent Health**. 32(5), 329-330.
- Mahidol University. Asian Institute for Health Development. (2002). **The behaviour surveillance survey of 7 target group in Bangkok**. Nakhon Pathom: Asian Institute for Health Development, 15-27.
- Mane, P. and McCaully P. A. (2000). **Impact of STIs including AIDS on Adolescents: A Global Perspective**. Proceeding from the International Conference on Adolescent Reproductive Health. Mumbai, India 1- 4 November.
- Manote Lortrakul. (1998). Suicide trends in Thailand: categorized by age and gender. **Journal of the Psychiatric Association of Thailand**. 43,1.
- Marc, J R. [et al.]. (2001). **Getting Health Reform Right**. [S.l: S.n].
- Marcell, V.A. [et al.]. (2002). Male adolescent use of health care services: where are the boys? **Journal of Adolescents Health**. 30(1), 35-43.
- Marcotte D, (1999). Adolescent Depression. Retrieved 10 January 2004, from <http://www.healthscout.com/ency/article/001518.htm>.
- Masatu, C. M et al. (2001). Use of health services and reported satisfaction among primary school adolescents in Arusha, Tanzania. **Journal of Adolescents Health**. 28(4), 278-287.
- Matsumoto S. (1999). Health needs of adolescents and sex education. [Online]. [S.l: S.n]. Retrieved from: <http://www.db.jhuccp.org/dbtw-wpd/exec>.
- McDowell, I. and Newell, C. (1996). **Measuring health: A guide to rating scales and questionnaires**. Oxford: Oxford University Press.
- McCraken, R.J. and Narayan, D. (1998). **Participation & Social Assessment**. USA: The International Bank for Reconstruction and Development, The World Bank.
- McKnight, G et.al. (2004). Perceived quality of life: A neglected component of adolescent health assessment and intervention. **Journal of Adolescents Health**. 34(4), 270-278.
- Medsearch. (2004). Retrieved October 1, 2004, from [http://www.gulflink.osd.mil/medsearch/glossary/glossary\\_d.shtml](http://www.gulflink.osd.mil/medsearch/glossary/glossary_d.shtml).
- Mensch B, Bruce J, Green M. (1999). **The Uncharted Passage: Girl's adolescence in the developing world**. USA: The Population Council.
- Mental Health. (2003). Mental Health: Depression. Retrieved 20 October 2003, from [www.gulflink.osd.mil/medsearch](http://www.gulflink.osd.mil/medsearch).
- Millstein, G.S et al. (1992). Health –Risk Behaviour and Health Concerns Among Young Adolescents. **Pediatrics**. 3(March).

- Millstein SG, Peterson AC, Nigthingale EO (1993). **Promoting the health of adolescents: New Direction for 21st. Century.** New York: Oxford University Press.
- Ministry of public Health [MoPH]. Department of Health. Family Planning and Population Division. (1998). **Thailand National Family Planning Program.** Nonthaburi: Division.
- Ministry of Public Health [MoPH]. (1999). Factsheet. Retrieved 23 May 2004, from: <http://www.anamai.moph.go.th/factsheet/school3>.
- Ministry of Public Health [MoPH]. (2000). **Ministry of Public Health report.** Nontaburi: MoPH.
- Ministry of Public Health [MoPH]. (2001). **Ministry of Public Health report.** Nontaburi: MoPH.
- Ministry of Public Health.[MoPH]. Department of Mental Health. (2002a). **The Thailand Health Profile.** Nonthaburi: Deapartment,.
- Ministry of Public Health [MoPH]. Department of Health. (2002b). **Proceeding of the workshop on Friend Corner.** Rimkok Resort, Chiang Rai, 18 February.
- Ministry of Public Health.[MoPH]. Department of Mental Health. (2003). **Child Development and Mental Health Assessment Form.** Nonthaburi: Department,.
- Newachec, P. W. [et al.]. ( 2003). Disparities in adolescent health and health care: does socioeconomic status matter?. Health Services Research. Retrieved October, 2003), from [http://www.findarticles.com/p/articles/mi\\_m4149/is\\_5\\_38/ai\\_110461719](http://www.findarticles.com/p/articles/mi_m4149/is_5_38/ai_110461719)
- Newman, I. and Innadda, S. (2000). Buddhism and adolescent Alcohol use. [Online]. Ph.D. Dissertation, University of Nebraska- Lincoln. Retrieved from <http://apha.confex.com/apha/128a/techprogrm>
- Nowarat Suwannapong and Chaveewan Boonsuya (2001). Evaluation and writing public health proposal. Thailand.: [S.n].
- Oana Xinhua. (2001, Jan 12). **Survey on children's day.** Bangkok Post Newspaper.
- Oberg, C., et., al. (2002). Health are access, sexually transmitted diseases, and adolescents: Identifying barriers and creating solutions 2002. **Current Problems in Pediatric and Adolescent Health Care**, 32(9), 320-339.
- Office of the Prime Minister, the National Statistical Office, the Social Statistics Division, (1998). **The report of the National Children and Youth Survey 1997.** Bangkok: Division,.
- Orasom (2000), "Dek Phan Mai Wai X" (New Generation Children X). 2000.
- Panee Panuwatsuk, Ratana Somrongthong. (1999). Sexual behaviour and the opinions about sexuality of adolescents in a slum community. **AIDS Journal**, 11(14): 208-216. (Thai version)



- Panee Panunvathsuk. et. al. (2002). Health illness and factors influencing the opinion about utilization of Banna primary health center of Muang district, Chumporn Province. Regional 11, **Medical Journal**. 16(1): 1-3.
- Pencharisky (2001). **The Proceeding of the International conference on Strategies for Enhancing Access to Medicines (SEAM) Conference 2001, Targeting Improved Access**: Proceeding, November 27-29, Washington DC.
- Penchan Charoensutthipan, ( 2001, June 12,). Adolescent. **Bangkok post**. (newspaper).
- Pennapa Hongthong. (2002, Jan 11). Youth Sell Drugs. **The Nation** (newspaper).
- Peter M. Barnes.(2003). **Measuring quality of life**. [Online]. [S.l: S.n]. Retrieved January 8, 2003.
- Pimpawan Boonmongkon. (1997). **Adolescent Sexual Health: Lesson Learned and New Direction**. Paper present in the Seminar on Future Needs of Thailand in Population and Reproductive Health, 17 –18 April, Melia Hotel. Prachuab Khirikan.
- Pimpawan Boonmongkon et al. (2000). **Thai Adolescent Sexuality and Reproductive Health: Implications for Developing Adolescents' Health Programs**. Nakhom Pathom: Center for Health Policy Studies, Faculty of Social Sciences and Humanities, Mahidol University.
- Poole, Dennis L. (1997). The safe project: Community –driven partnerships in health, mental health, and education. **Health & Social Work**. 22(4), 282 –230.
- Population Reference Bureau. (2004). Youth. [Online]. Retrieved 15 August 2004, from [www.prb.org/template.cfm](http://www.prb.org/template.cfm)
- Population report. (2001). **UCLA [Newsletter]**, xxix (3). (Series L, Number 12 issues in World Health: Why so vulnerable?).
- Prapaipan Supachaturaj. (1984). **The Role of the Bangkok Metropolitan Administration in Family Planning and Maternal and Child Health Care in the Slum Areas**. A proceeding of the workshop on Family Planning Needs of and Services for the Underprivileged Urban: The Case of Bangkok, January 26 – 28, ,Wongse Amataya Hotel, Chonburi.
- Prawase Wasi. (2000). **Triangle that moves the mountain and health systems reform movement in Thailand**. Nonthaburi: Health System Research Institute [HSRI], Ministry of Public Health [MoPH].
- Prescott, J.J. et al. (1998). Carlton, Prediction of major depression and dysthymia from CES-D scores among ethnic minority adolescents. **J. Am. Acad. Child Adolesc. Psychiatr**. 37, 495–503.
- Price, Richard H., Coici, Madalyn. (1993). Webs of influence: school and community programs that enhance adolescent health and education. **Community and Schoolteacher College Record**. 94(3), 478.

**Proceedings of the Seminar on Family Planning Needs of and Services for the Underprivileged Urban: The Case of Bangkok (1984).** January 26 – 28, Wongse Amataya Hotel, Chonburi.

Radloff, L S. (1991). The use of the Center for Epidemiologic Studies Depression Scale in adolescents and young adults. **J Youth Adolesc.** 20, 149–166.

Rajaratnam A, Jeyalakshmi S, Greed A, Vijayakumari E. (1997). **Adolescence.** India: Christian Medical College and Hospital. Rajan Offset Printer.

Raphael, D et al.(1996). The Quality of Life. **Public Health**, 34(2). 1996, 443-446.

**Raphael, Renwick, Brown.**(1998). Health Promotion;. **Journal of School Health**, 68(7), 297-300.

Raphael, Rukholm et al. (1996). The quality of life profile-Adolescent version: Background, Description and initial validation. **Journal of Adolescents Health.** 9, 366-375.

Ratana Somrongthong, & Chitr Sitti-amorn. (2000). Existing health needs and related health services for adolescents in slum community in Thailand. **International Journal of Adolescent Medicine and Health.** Vol.12, No2-3 No,191-202.

Ratana Somrongthong [et al.]. (2003). Sexual behaviours and opinions on sexuality of adolescents in a slum community in Bangkok. **Southeast Asian J Trop Med.** 34(2), 443-446.

Raty, L. K. A et al. (2003). Quality of life in youth: a comparison between adolescents and young adults with uncomplicated epilepsy and healthy controls. **Journal of Adolescent Health.** 33(4), 252-258.

Rivera, PM (2000). The relationship between exposure to violence and quality of life of urban adolescents. Dissertation Abstracts International. University of Missouri, Kansas, USA.

Robert, J.M et.al. (2001). **Getting Health Reform.** [S.l: S.n]

Roberts, RE et.al. ( 1990).Assessment of depression in adolescents using the Center for Epidemiologic Studies Depression Scale. *Psychological Assessment.* 1990;2(2):122.

Rushton, M. et.al(1999). Epidemiology of depressive symptoms in the National Longitudinal Study of Adolescent Health. **J A Acad Child Adolesc Psychiatry.** 41, 199–205.

Rutherford. et al.(1999). Health profession students as adolescent mentors in the Seattle school. **Journal of School Health.** 69,2: 77-82.

Rutter. (1986). The developmental psychopathology of depression: Issues and perspectives. In: M. Rutter, C.E. Izard and P.B. Read, (Eds). **Depression in Young People: Developmental and Clinical Perspectives,** (pp. 3–30). New York: Guilford.

- Sakondhavat et.al. (1991). Health education and services in human reproductive contraception for adolescent. In: Pimpawan Boonmongkol, (2000). **Thai Adolescent Sexuality and Reproductive Health: Implications for Developing Adolescents' Health Programs**. Nakhom Pathom: Center for Health Policy Studies, Faculty of Social Sciences and Humanities, Mahidol University
- Samai Arpapurum, (1997). Adolescent Sexuality: Heath System Research Institute, Ministry of Public Health. (Thai version)
- Sanhueza, XA et al.(2000). **Guide for monitoring equitable access to basic health services**. Draft research proposal. [Unpublished document, cited in WHO].
- Save the Children (UK). (2001). **Breaking Trough the Clouds. A Participatory Action Research (PAR)**. Project with Migrant Children and Youth Along the Borders of China, Myanmar and Thailand.
- Schensul. J.J. (1998). Community-based risk prevention with urban youth. **School Psychology Review**. 27(2), 233-245.
- Selener, D. (1997). **Participatory Action Research and Social Change**. USA: Cornell University.
- Senderowitz, J. (1997). Health facility programs on reproductive health for young adults. FOCUS on young adults research series.[Online]. Retrieved from **www.pathfind.org/RPPPS-paper/health**.
- Sentelli, J. et al. (1998). Managed care, school health prams and adolescent health services: opportunities for health promotion. **Journal of School Health**. 68(10), 434 – 441.
- Service de medicine pour adolescents** [Online]. Retrieved 28 October 2002. [Federation de pediatic, hospital Bicetre, 78, rue du General-Leclerc, 94257, Le Kremlin-Bicetre cedex, France ]
- Service Gaps (2002). Service Gaps. Retrieved 20 July 2002, from:[http://www.bihartimes.com/anant\\_pandey.html](http://www.bihartimes.com/anant_pandey.html).
- Sharma, R. (2003). **Contraceptive use among university undergraduate female students**. Master degree of Primary Health Care Management. Mahidol University.
- SIL International. (1999). What is a felt needs? Retrieved 20 June 2004, from [www.sil.org.lingualinkik/literacy](http://www.sil.org.lingualinkik/literacy)..
- Sikes, J A.(1996). Approach to adolescent reproductive health. **International Journal of Health Promotion and Education**. III(3),15-17.
- Simeoni, C.M et al. (2001). Health-related quality of life reported by French adolescents: A predictive approach of health status? **Journal of Adolescents Health**. 28(4), 288-294.

- Sirikul Bunnag, (2002, September 5). Report of Ms.Orsom's study (Dek Phan Mai Wai X). **Bangkok Post**. (newspaper).
- Slap, G. (1998). **Finding: Focus group discussion**. [S.l: S.n]. (unpublished document.)
- Smith, D et al. (2003). Socio-cultural contexts of adolescent sexual behavior in rural Hanover, Jamaica. **Journal of adolescent health**. 133(1), 41-48.
- Social Statistics Division, National Statistical office, Office of the Prime Minister. (1998). **The report of the children and youth survey 1997**. Bangkok: Division.
- Sosa, P and Suciion (1993). Paraguay, Centro de Estudios de publication (CEPCP),131-135. Retrieved from [www.unescobkk.org/ips/arh-web/PDF/Thailand](http://www.unescobkk.org/ips/arh-web/PDF/Thailand)
- Students abortions increasing. (2001, August 9 ). **The Nation**. (newspaper).
- Subhatra Bhumiprabhas. (2001, October 25). Juvenile. **The Nation**. (newspaper).
- Suwat, M et.al. (1997). **Comparison of the WHOQoL-100 and the WHOQoL-BREF**. Nonthaburi: Department of Mental Health, MoPH. Thailand.
- Tamura, M., Nopporn Howteerakul and Nawarat Suwanapong. ( 2000). HIV/AIDS Prevention Behaviour among Adolescents in Klong Toey Community Congested Area. **Journal of Public Health** 2000: 32(2):87-99.
- Tanida Siroratrtanakul, (1998, June, Monday 1). Adolescent and smoking. **Bangkok Post**. (newspaper).
- Think PN. (2001). **Assessing accessibility to primary care for households in the urban area of Danang City in Vietnam**. Master of Public Health, College of Public Health, Chulalongkorn University, Thailand.
- Topolski, D.T. et al. (2001). Quality of life and health-risk behaviors among adolescents. **Journal of Adolescent Health**. 129(6), 426-435.
- The National Adolescent Health Information Center, USA ( 2002). Adolescent. Retrieved June 2002, from [www.youth.ucsf.edu/nahic/img](http://www.youth.ucsf.edu/nahic/img).
- The Partnership for Child Development. (1999). **School health & nutrition: A Situation Analysis, a participatory approach to building programs that promote health, nutrition and learning in schools**. [S.l: S.n].
- The Quality of Life Center, Denmark. (2001). What is QoL?. Retrieved 30 May 2001. From, <http://www.fclk.dk/english/index.asp>.
- The School of Nursing, Laurentian University of Toronto, Canada (2002). Quality of Life Profile–Adolescents' Version. Retrieved 30 April 2002, from <http://www.utoronto.ca.qol.adolescents.htm>.
- The University of Washington ( ). The Youth Quality of Life Instrument - Research Version (YQOL-R). Retrieved 1 October 2001, from [http://depts.washington.edu/yqol.about\\_Us.html](http://depts.washington.edu/yqol.about_Us.html).

- The Wonder CDC (Centers for Disease Control and Prevention), USA. (2001). Mortality 1999. Retrieved 7 April 2002, from [www.wonder.cdc.gov](http://www.wonder.cdc.gov).
- The World Bank Group. (2004). Tools and data. Retrieved 2 January 2004, from <http://www.worldbank.org/poverty/impact/methods/indepth.html> 11 Apr 04
- The World's Youth- 1996 Population Reference Bureau, Inc. (1996). Cited in the *Adolescence: the Critical phase*, WHO/SERO1997. p4.
- Tiwari, S. (2003). **Access to and Utilization of Health Services in the Areas With and Without Civil Conflict in Nepal: A Case for Tuberculosis Services Through District Health Facilities**. Philosophy of Public Health, The College of Public Health, Chulalongkorn University.
- Topolski TD et al. (2001). Quality of life and health-risk behaviours among adolescents. *Journal of Adolescent Health*, 29 (6), 426-435.
- Trochim W. (2002). Qualitative Measures. Retrieved 2 August 2004, from <http://www.socialresearchmethods.net/kb/qual.htm>.
- Ueno, K. (2004). **The effects of friendship networks on adolescent depressive symptoms**. [S.l: S.n]
- Ulrike Ravens-Siebere (2000). CAT-SCREEN- CD Rom. Ulrike Ravens-Siebere, Denmark, Website [www.catscreen.de](http://www.catscreen.de).
- Umaporn Trungkasombaddi, Vachira Lapboonsap, and Piyalumporn Hawanon. (1998). Application of CES-D for Adolescents' Depression Screening. *Journal of Mental Health Association of Thailand*. 42(1)
- UNESCO. (1999a). Reproductive Health [Online]. Retrieved 15 July 2002, from <http://www.unescobkk.org/ips/arh-web/PDF/Thailand>.
- UNICEF.(1999). Monitoring and Statistics [Online]. Retrieved 10 December 2002, from [http://www.unicef.org/statis/country\\_1page174html](http://www.unicef.org/statis/country_1page174html)
- UNICEF (2001). Community participation. Retrieved 10 December 2002, from [www.unicef.org/programme/gpp/communi/commparti.html](http://www.unicef.org/programme/gpp/communi/commparti.html)
- United Nations [UN]. (1998). The World Programme of Action for Youth to the Year 2000 and Beyond: Guiding the future of humanity: Strengthening United Nations commitment to youth. Retrieved 10 April 2002, from <http://www.un.org/events/youth98/backinfo/ywpa2000.htm>.
- University of Maryland, Baltimore. USA. (1995). Project SUCCESS: A model for university- school-community partnerships. *Social Work in Education*, 17(4), 256-262.
- University of Toronto,(2001). Adolescents' Quality of Life [Online]. Retrieved August, 31 2004, from [www.utoronto.ca/Qol/adolescents.htm](http://www.utoronto.ca/Qol/adolescents.htm).

- Unsafe sex. (2001, June 8.). **The Nation**. (newspaper)
- Venus Udomprasertgul, et al. (2002). Student behavior towards playing computer games at secondary school level. **The Thai J. Health Research**, 16(2), 149–158.
- Virginia Gil-Rivas., et al.( 2003).Understanding depressed mood in the context of a family-oriented culture. **Adolescence**. 38 (149), 93-7
- Vogel, Julie S.,et al.(2003). The relationship between depression and smoking in adolescents. **Adolescence**.38(149).
- Wade, J. Cairney and D.J. Pevalin, (2002). Emergence of gender differences in depression during adolescence: national panel results from three countries, **J. Am. Acad. Child Adolesc. Psych**. 41, 190–198.
- Wage, T.J. [et al]. (2002). Emergence of gender differences in depression during adolescence: national panel results from three countries, **J. Am. Acad. Child Adolescence. Psych**. 41,190–198.
- Walsh, M and Kimberly A. et al. (1999). School counselors in school-community partnerships: Opportunities and challenges. **Professional School Counseling**.,2(5), 349-357.
- Wang, M.C., Haertel, G.D., & Walberg, J.H. (1995). The effectiveness of collaborative school-link services. In L. C. Rigsby, M.C. Reynolds, & M.C. Wang (Eds.). **School- community connections: Exploring issues for research and practice** (pp.283-309). San Francisco: Jossey Bass.
- Ware, J. [et al]. (1993). The short form SF-36. In: Ian McDowell & Claire Newell (Eds). (1996). **Measuring Health: a guide to rating scales and questionnaires**. New York: Oxford University Press.
- Wasana Pattanakamjorn (1999). Factors associated with Amphetamine use in students. **Journal of Psychiatrist Association**. 43(4), 345- 357.
- Weiser, Sarah Peyton. (19--). **Adolescents' use of health services**. [Online]. Dissertation Abstracts International, Ph.D.dissertation. Purdue University. Retrieved from [http:// thailis.db.car.chula.ac.th/dao/detail.nsp](http://thailis.db.car.chula.ac.th/dao/detail.nsp)
- Whyte, S. (2000). **Community Participation in Essential National Health Research**. [S.l ]: Council on Health Research for Development (COHRED).(Document 200.5 June 2000).
- Wilcox, D. (2000).Create Partnerships. Retrieved 10 July 2002, from [www.partnerships.org.uk/AZP/part.html](http://www.partnerships.org.uk/AZP/part.html).
- Winslow, W.B.(2003). Family care giver's experiences with community services: A qualitative analysis. **Journal of Public Health Nursing**. 20(5), 333-335.
- Wodarski, J S., Wodarski L A., & Dulmus, C N. (2003). Adolescent Depression and Suicide: A Comprehensive Empirical Intervention for Preventionand Treatment. [S.l].: Springfield, IL: Charles C Thomas,

- World Health Organization [WHO]. (1989). **The health of youth**. Geneva: WHO.
- World Health Organization [WHO]. (1993a). **The urban health crisis-Report of the technical discussions at the 44<sup>th</sup>.World Health Assembly**. Geneva.: WHO.
- World Health Organization [WHO]. (1993b). **WHOQOL**. Geneva.: WHO.
- World Health Organization [WHO]. (1996a). **Integration of health care delivery**. Geneva: WHO
- World Health Organization [WHO]. (1996b). WHOQOL-BREF: Introduction, Administration, Scoring and Generic Version of the Assessment: Field Trial Version. Geneva: WHO. Retrieved from [www.who.int/msa/qol/ql4.hmt](http://www.who.int/msa/qol/ql4.hmt), WHOQoL Group,
- World Health Organization [WHO]. (1996c). **Qualitative Research for Health Programmes**. Geneva: Division of Mental Health and Prevention of Substance Abuse.
- World Health Organization [WHO]. (1997). **Adolescence: The critical phase**. Delhi.: SEARO.
- World Health Organization [WHO]. (1998a). Strategies for adolescent health and development in South East Asia Region. Retrieved from <http://w3.whosea.org/adolescent/presentations.htm> (WHO/SEARO, 26-29 May).
- World Health Organization [WHO]. (1998b). **Improving Adolescent Health and Development. Adolescent Health and Development Programme, Family and Reproductive Health**. Geneva: WHO.
- World Health Organization [WHO]. (1998c). Health Promotion in the 21<sup>st</sup> century: an era of partnerships to achieve health for all. Retrieved 1 June 2002. from <http://www.who.int/inf-pr-1998/en/pr98-47.html>.
- World Health Organization [WHO]. (1999). **Programming for Adolescent Health Development: Report of a WHO/UNFPA/UNICEF Study Group on Promoting for Adolescent Health**. Geneva: WHO.
- World Health Organization [WHO]. (2000). **Equity in access to public health: Report and documentation of the technical discussions held in conjunction with the 37th meeting of CCPDM, New Delhi, 31 August**. India: WHO.
- World Health Organization [WHO]. (2001). **Regional Framework for Introducing Life skills Education to Promote the Health in Adolescents**. [India]: WHO.
- World Health Organization [WHO]. (2002a). Adolescent Health and Development Programme, Family and Reproductive Health. [S.l]: WHO. Retrieved from: <http://www.who.int/reproductive-health/adolescent>.
- World Health Organization [WHO]. (2002b). Retrieved from [http://www.who.int/child-adolescent-health/HIV/HIV\\_adolescents.htm](http://www.who.int/child-adolescent-health/HIV/HIV_adolescents.htm).

World Health Organization [WHO]. (2003). **World Health Organization and International Agency for the Prevention of Blindness**. [Geneva: WHO].

World Health Organization [WHO]. Quality of Life. Retrieved June 2003, from [www.who.int/msa/qol/q14hmt](http://www.who.int/msa/qol/q14hmt)

Yang, H.J et.al. (2004). Using the CES-D in a two-phase survey for depressive disorders among no referred adolescents in Taipei: a stratum-specific likelihood ratio analysis. **Journal of Affective Disorders**, Retrieved July 1, 2004 (In Press)


Yu Feng. (2003). **Attitude toward adolescent pregnancy, induced abortion and supporting health services among high school students in Nakorn Pratum Province**. Master of Primary Health Care Management, Mahidol University.

Zhang, K. et al. (2000). Adolescent Sexuality and Sexual Health in China. Community-Based Programs for adolescent sexual health and domestic violence against women. In: Pimpawun Boonmongkol, Anjana Suvarnananda. [Eds.]



## **APPENDICES**

## Appendix A Community Map



Source: Bunyit, Et al. 2002

## Appendix B

### Adolescent Questionnaire

- Part 1. General Information  
 Part 2. Health Need and health services  
 Part 3. QoL  
 Part 4. Depression

#### Part 1 General Information

Please tick  in the bracket ( ) or fill in the space .....

1. Your birth date. Day...../month...../year.....
2. Sex
  1. ( ) male
  2. ( ) female
3. Where were you born?
  1. ( ) Bangkok
  2. ( ) other province (please specify.....)
4. If born in other province, how long you have been living in Bangkok?  
 .....months, .....years.
5. You are now living in (name of sub- community) .....
6. Marital status
  1. ( ) single
  2. ( ) married
  3. ( ) separate
  4. ( ) divorced
  5. ( ) widowed
7. The number of persons living in your house.  
 ..... persons.
8. You are the .....child of the family.  
 1 ( )1<sup>st</sup>    2 ( )2<sup>nd</sup>    3 ( )3<sup>rd</sup>    4 ( )4<sup>th</sup>    5 ( )5<sup>th</sup>    6 ( )6<sup>th</sup>    7 ( )7<sup>th</sup> or over
9. Parents' living status
  1. ( ) living together
  2. ( ) separated
  3. ( ) divorced
  4. ( ) either father or mother passed away
  5. ( ) both of father and mother passed away

10. Who is your present guardian?

1.  father and mother
2.  father
3.  mother
4.  other (please specify.....)

11. Your current guardians' occupation?

1.  laborer
2.  seller
3.  farmer
4.  government or privatization officer
5.  other (please specify.....)

12. What is your highest education you received?

12.1  none

12.2  I am now studying in  Primary school, year 4  5  6   
 High school, year 1  2  3   
 College/University, year 1  2   
 3  4

12.3  graduated in  Primary school, year 4  5  6   
 High school, year 1  2  3   
 College/University 1  2  3  4

13. Are you currently working?

1.  no ( go to # 15)
2.  yes – part time
3.  yes – full time

14. If working, monthly income of yourself (in Baht)

1.  none
1.  less than 2,000
2.  2,000-4,999
3.  5,000-6,999
4.  7,000-9,999
5.  10,000-13,000
6.  over 13,000

15. The monthly income of your family (in Baht)

7.  less than 5,000
8.  5,000-9,999
9.  10,000-14,999
10.  15,000-19,999
11.  over 20,000
12.  don't know

16. Your household expenditure's circumstance

1.  able to save some
2.  enough
3.  sometime enough
4.  not enough and on debt
5.  don't know

17. How would you rate loving and caring of your family?

1.  very high
2.  fair
3.  less
4.  none

18. Whom would you turn to when facing with health problem?

1.  friend
2.  brother/ sister
3.  father/ mother
4.  other relative
5.  teacher
6.  other (please specify.....)

19. How do you rate the friends that you have?

- |   |   |
|---|---|
| 1 <input type="checkbox"/> not at all satisfied | 2 <input type="checkbox"/> not very satisfied |
| 3 <input type="checkbox"/> satisfied            | 4 <input type="checkbox"/> Very satisfied     |
| 5 <input type="checkbox"/> Extremely satisfied  |   |

20. Do you have a close boy/girl friend?

- |                                 |                                |
|---------------------------------|--------------------------------|
| 1. <input type="checkbox"/> Yes | 2. <input type="checkbox"/> No |
|---------------------------------|--------------------------------|

21. Please response truly the following sentences regarding family and peer relationships by tick  only one

### Family relationships & Peer relationships

Items	Rare or not at all	some or a little of the time	occasionally or moderate	most of the time
<b>Family relationships</b>				
● Getting along with my family				
● Consult with parent/relative when I have problem				
● Having things to do with family (recreation, chat etc.)				
● Quarrel with family				
<b>Peer relationships</b>				
● Getting along with peer				
● Consult with peer when I have problem				
● Having things to do with peer (recreation, chat etc.)				
● Quarrel with peer				

**Part 2 Adolescent's health and the need of health service**

22. What is your opinion of sex education in Thailand?  
 1.  enough  
 2.  not enough (reason.....)
23. Do you have any welfare, health care/30 Baht card, student health cards, or any cards relating to health service access which you can use to discount or cover your health service charge when accessing governmental or private health service?  
 1.  yes, the card is.....  
     a.  Welfare card    b.  30 Baht card    c.  Student card  
     d.  Private insurance card    e.  other specify.....  
 2.  no (go to #22)
24. Have you ever used the card to cover your health service?  
 1.  yes (in what matter.....)  
 2.  never (reason.....)
25. Do you have any chronic illness?  
 1.  yes .... Specify .....
2.  no  
 3.  not sure
26. During the past 2 months, have you been sick?  
 1.  yes  
 2.  no  
 3.  cannot remember
27. Your first sexual intercourse's experience  
 1  Never ( go to # 29)  
 2  Yes, when aged.....years.
28. How often do you or sex partner wear condom?

**Your partners**  
*(Can tick  $\checkmark$  more than one)*

**Frequency of use of a condom**  
*(Tick  $\checkmark$  only one)*

- |   | always                   | sometime                 | never                    |
|---|--------------------------|--------------------------|--------------------------|
| 1. <input type="checkbox"/> girlfriend/ boyfriend | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 1. <input type="checkbox"/> acquaintance          | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 2. <input type="checkbox"/> sex worker            | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 3. <input type="checkbox"/> bar girl, waitress    | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |
| 4. <input type="checkbox"/> other (.....)         | <input type="checkbox"/> | <input type="checkbox"/> | <input type="checkbox"/> |

29. Please tick the following list which one is sexual transmitted diseases (STDs)? **Can tick  $\surd$  more than one**
1. ( ) Gonorrhoea
  2. ( ) Syphilis
  3. ( ) Chancroid
  4. ( ) Prostate Gland Cancer
  5. ( ) Candidiasis
  6. ( ) Cystitis
  7. ( ) Herpes at genital area
  8. ( ) Cervical Cancer
  9. ( ) Trichomoniasis
  10. ( ) Condiloma
30. Please see the further next 2 pages, there are a set of questions ask about **yours opinion** towards adolescents' health problems and how they manage those problems by gender.
31. Please see the next 2 pages, there are a set of questions ask about **your health problems** and how do you manage those problems by gender

**FEMALE ONLY**

( your opinion towards other adolescents)

30 The following items in column (a), I would like to ask you opinion..... in column (b), (c), and (d) , please answer as many choices as you think are appropriated

Column b : Do you think it is a problem for other people of your age?

Column c : If YES, Who do you think they would talk to in their family about the problem?

Column d : If YES, Who do you think who they would talk to outside their family about the problem?

Column e : Do you think what health care facility they would like to visit about the problem?

**Table 2**

No.	(a) Items	(b) Do you think it is a problem for them ?	(c) Who would they talk to?							(e) What health care facility they would like to visit?					
			(c) People in their family			(d) People outside their family				Visit to					
			parent	sibling	peer	youth leader	community leader	community health volunteer	housewife group	gov. hospital	private hospital	health center	private clinic	DPF	other
<b>1</b>	<b>Sexuality and reproductive health</b>														
1.1	<i>Development of sexual organs ; e.g. breast enlargement</i>	Y N													
1.2	<i>Acne</i>	Y N													
1.3	<i>Having period</i>	Y N													
1.4	<i>Hygienic care for genital organ</i>	Y N													
1.5	<i>Dysmenorrhea</i>	Y N													
1.6	<i>Irregular period</i>	Y N													
1.7	<i>Masturbation</i>	Y N													
1.8	<i>Vaginal itching</i>	Y N													
1.9	<i>Having sexual intercourse</i>	Y N													
	-with boy friend	Y N													
	-with other for gift or money	Y N													
	-with acquaintance	Y N													
	- with girl friend	Y N													
1.10	<i>Having unsafe sex</i>	Y N													
1.11	<i>Family planning</i>	Y N													
	-using contraceptive pills for birth control	Y N													
	-condom use of sex partner	Y N													
	- others specify _____	Y N													
1.12	<i>Unintended pregnancy</i>	Y N													
1.13	<i>Abortion</i>	Y N													



**FEMALE ONLY**

( your opinion towards *other* adolescents)

30 The following items in column (a), I would like to ask you opinion..... in column (b), (c), and (d), please answer as many choices as you think are appropriated

Column b : Do you think it is a problem for other people of your age?

Column c : If YES, Who do you think they would talk to *in their family* about the problem?

Column d : If YES, Who do you think who they would talk to *outside their family* about the problem?

Column e : Do you think what health care facility they would like to visit about the problem?

Table 2 (cont)

No.	(a) Items  PROBLEMS	(b) Do you think it is a problem for them?		(c) Who would they talk to?						(d) Who would they talk to?						(e) What health care facility you would like to visit?					
				(c) People in their family		(d) People outside their family						Visit to									
				parent	sibling	peer	youth leader	community leader	community health volunteer	housewife group	gov. hospital	private hospital	health center	private clinic	DPF	other					
1.14	-burning urination	Y	N																		
1.15	- pus /discharge at vagina																				
1.16	HIV/AIDS infected	Y	N																		
1.17	Body and shape (too fat or too thin)	Y	N																		
1.18	sexual harassment	Y	N																		
2	<b>Injury and Violence</b>	Y	N																		
2.1	Road accident	Y	N																		
2.2	Home violence	Y	N																		
2.3	Fighting	Y	N																		
2.4	Rape	Y	N																		
3	<b>Mental health</b>	Y	N																		
3.1	Depression	Y	N																		
3.2	Stress	Y	N																		
3.3	Lonely	Y	N																		
3.4	Sad	Y	N																		
4	<b>Substances abuse</b>																				
4.1	Drinking alcohol	Y	N																		
4.2	Smoking	Y	N																		
4.3	Drug addict ( amphetamine)	Y	N																		
4.4	Drug addict ( heroin, marijuana, glue , etc..)	Y	N																		

**FEMALE ONLY**

31 The following items in column (a), I would like to ask you opinion..... in column (b), (c) and (d) , please answer as many choices as you think are appropriated

Column b : Whether you have had experienced with .....

Column c : Whether you have problems ?

Column d : Who would you talk to *in your family* about the problem? ( can answer more than 1 )

Column e : Who you would talk to *outside of your family* about the problem? ( can answer more than 1 )

Column f : What health care facility you would like visit about the problem? ( can answer more than 1 )

Table 1

No.	(a) Items	(b) Do you ever have experience with.... ?	(c) Do you have the problem?	Who would they talk to?							(f) What health care facility you would like to visit?						
				(d) People in your family		(e) People outside your family					Visit to						
				parent	sibling	peer	youth leader	community leader	community health volunteer	housewife group	gov. hospital	private hospital	health center	private clinic	DPF	other	
1	<b>Sexuality and reproductive health</b>																
1.1	<i>Development of sexual organs ; e.g. breast enlargement</i>	Y N	Y N														
1.2	<i>Acne</i>	Y N	Y N														
1.3	<i>Having period</i>	Y N	Y N														
1.4	<i>Hygienic care for genital organ</i>	Y N	Y N														
1.5	<i>Dysmenorrhea</i>	Y N	Y N														
1.6	<i>Irregular period</i>	Y N	Y N														
1.7	<i>Masturbation</i>	Y N	Y N														
1.8	<i>Vaginal Itching</i>	Y N	Y N														
1.9	<i>Having sexual Intercourse</i>	Y N	Y N														
	-with boy friend	Y N	Y N														
	-with other for gift or money	Y N	Y N														
	-with acquaintance	Y N	Y N														
	- with girl friend	Y N	Y N														
1.10	<i>Having unsafe sex</i>	Y N	Y N														
1.11	<i>Family planning</i>	Y N	Y N														
	-using contraceptive pills for birth control	Y N	Y N														
	-condom use of sex partner	Y N	Y N														
	- others specify _____	Y N	Y N														
1.12	<i>Unintended pregnancy</i>	Y N	Y N														
1.13	<i>Abortion</i>	Y N	Y N														

**FEMALE ONLY**

31 The following items in column (a), I would like to ask you opinion..... in column (b), (c), (d) and (f), please answer as many choices as you think are appropriated

Column b : Whether you have had experienced with .....

Column c : Whether you have problems ?

Column d : Who would you talk to in your family about the problem? ( can answer more than 1)

Column e : Who you would talk to outside of your family about the problem? ( can answer more than 1 )

Column f : What health care facility you would like visit about the problem? ( can answer more than 1 )

Table 1 (cont)

No.	(a) Items	(b) Do you ever have experience with.... ?		(c) Do you have the problem?		Who would they talk to?						(f) What health care facility you would like to visit?						
						(d) People in your family			(e) People outside your family			Visit to						
						parent	sibling	peer	youth leader	community leader	community health volunteer	housewife group	gov. hospital	private hospital	health center	private clinic	DPF	other
1.14	-burning urination	Y	N	Y	N													
1.15	- pus /discharge at vagina	Y	N	Y	N													
1.16	HIV/AIDS infected	Y	N	Y	N													
1.17	Body and shape (too fat or too thin)	Y	N	Y	N													
1.18	sexual harassment	Y	N	Y	N													
2	<b>Injury and Violence</b>	Y	N	Y	N													
	2.1 Road accident	Y	N	Y	N													
	2.2 Home violence	Y	N	Y	N													
	2.3 Fighting	Y	N	Y	N													
	2.4 Rape	Y	N	Y	N													
3	<b>Mental health</b>	Y	N	Y	N													
	3.1 Depression	Y	N	Y	N													
	3.2 stress	Y	N	Y	N													
	3.3 lonely	Y	N	Y	N													
	3.4 sad	Y	N	Y	N													
4	<b>Substances abuse</b>																	
	4.1 Drinking alcohol	Y	N	Y	N													
	4.2 Smoking	Y	N	Y	N													
	4.3 Drug addict ( amphetamine)	Y	N	Y	N													
	4.4 Drug addict ( heroin, marijuana, glue , etc..)	Y	N	Y	N													

**Part 3 Accessibility****Geographic accessibility and utilization**

32. Please tick (  ) any of the following services that are available in your community
- |  |  |
|--|--|
| 1 ( <input type="checkbox"/> ) Government hospital | 2 ( <input type="checkbox"/> ) Private hospital    |
| 3 ( <input type="checkbox"/> ) Private clinic      | 4 ( <input type="checkbox"/> ) Health Center       |
| 5 ( <input type="checkbox"/> ) Drug store          | 6 ( <input type="checkbox"/> ) Fitness & Beauty    |
| 7 ( <input type="checkbox"/> ) Hot line counseling | 8 ( <input type="checkbox"/> ) Others specify..... |
33. When you have had health problem, where did you usually go? (you can check one or more)
- |  |   |
|--|---|
| 1 ( <input type="checkbox"/> ) Drug store          | 2 ( <input type="checkbox"/> ) Health Center              |
| 3 ( <input type="checkbox"/> ) Private Clinic      | 4 ( <input type="checkbox"/> ) Private Hospital           |
| 5 ( <input type="checkbox"/> ) Government hospital | 6 ( <input type="checkbox"/> ) Traditional healer         |
| 7 ( <input type="checkbox"/> ) DPF                 | 8 ( <input type="checkbox"/> ) Community health Volunteer |
34. How long does it take from your location to the facility ( that you answered in # )
- |  |   |
|--|---|
| 1 ( <input type="checkbox"/> ) less than 30 mins | 2 ( <input type="checkbox"/> ) 30 mins – 1 hour |
| 3 ( <input type="checkbox"/> ) 1 – 2 hour        | 4 ( <input type="checkbox"/> ) over 2 hrs.      |

**For each of the services list below, which you last visited, please answer the following questions.**

Items	Place					
	Health center	Private Clinic	Gov Hospital	Private Hospital	DPF	Others
35. Where did you last visit a health center facility						
A Past 1 month						
B Past 2 months						
C Past 3 months						
D Past 4 months						
E Past 12 months						
36. On your last visit, How long did it take						
To travel to health center						
A less than 30 minutes						
B 30-60 minutes						
C 60-120 minutes						
D more than 120 minutes						
37. On your last visit, How would you rate						
Difficulty of utilization						
A Very easy						
B Easy						
C Moderate						
D Difficult						
E Very difficult						





**Part 4 QoL**

52. Please read each question, assess your feeling, and circle the number on the scale for each question that gives the best answer for you

No	Items	Very good	Poor	Neither poor nor good	Good	Very good
52.1	How would you rate your quality of life	1	2	3	4	5

No	Items	Very dissatisfied	Dissatisfied	Neither satisfied nor dissatisfied	satisfied	Very satisfied
52.2	How satisfied are you with your health?	1	2	3	4	5

No	Items	Not at all	A little	A moderate amount	Very much	An extreme amount
52.3	To what extent do you feel that physical pain prevents you from doing what you need to do?	1	2	3	4	5
52.4	How much do you need any medical treatment to function in your daily life?	1	2	3	4	5
52.5	How much do you enjoy life?	1	2	3	4	5
52.6	To what extent do you feel your life to be meaningful?	1	2	3	4	5
52.7	How well are you able to concentrate?	1	2	3	4	5
52.8	How safe do you feel in your daily life?	1	2	3	4	5
52.9	How healthy in your physical environment?	1	2	3	4	5
52.10	Do you have enough energy for everyday life?	1	2	3	4	5
52.11	Are you able to accept your bodily appearance?	1	2	3	4	5
52.12	How you enough money to meet your needs?	1	2	3	4	5
52.13	How available to you is the information that you need in your day to-day life?	1	2	3	4	5
52.14	To what extent do you have the opportunity for leisure activities?	1	2	3	4	5

The following question ask about how satisfied you have experienced certain things in the last two weeks

No	Items	Very dissatisfied	Dissatisfied	Neither satisfied nor dissatisfied	satisfied	Very satisfied
52.15	How well are you able to get around?	1	2	3	4	5
53.16	How satisfied are you with your sleep?	1	2	3	4	5
52.17	How satisfied are you with your ability to perform your daily living activities?	1	2	3	4	5
52.18	How satisfied are you with your capacity for work?	1	2	3	4	5
52.19	How satisfied are you with yourself?	1	2	3	4	5
52.20	How satisfied are you with your personal relationships?	1	2	3	4	5
52.21	How satisfied are you sex life?	1	2	3	4	5
52.22	How satisfied are you with the support you get from your friend?	1	2	3	4	5
52.23	How satisfied are you with the conditions of your living place?	1	2	3	4	5
52.24	How satisfied are you with your access to health services?	1	2	3	4	5
52.25	How satisfied are you with your transportation?	1	2	3	4	5
52.26	How often do you have negative feelings, such as blue mood, despair , anxiety, depression?	1	2	3	4	5



## **Part 5 Depression**

53. Instructions for questions:

Below is a list of the ways you might have felt or behaved, Please tell me **how often** you felt this way during the past week

<b>During past week</b>	<b>Rarely or none of the time  (less than 1 day)</b>	<b>Some or a little of the time  (1-2 days)</b>	<b>Occasionally or a moderate amount of the time (3 - 4 days)</b>	<b>Most or all of the time  (5 -7days)</b>
53.1 I was bothered by things that usually don't bother me				
53.2 I did not feel like eating; my appetite was poor				
53.3 I felt that I could not shake off the blues				
53.4 I felt that I was just as good as other people				
53.5 I had trouble keeping my mind on what I was doing				
53.6 I felt depressed				
53.7 I felt that everything I did was an effort				
53.8 I felt hopeful about the future				
53.9 I thought my life had been a failure				
53.10 I felt fearful				
53.11 My sleep was restless				
53.12 I was happy				
53.13 I talked less than usual				
53.14 I felt lonely				
53.15 People were unfriendly				
53.16 I enjoyed life				
53.17 I had crying spells				
53.18 I felt sad				
53.19 I felt that people dislike me				
53.20 I could not get "going"				



## \*\*สำหรับวัยรุ่นชาย\*\*

แบบสอบถามเลขที่

### โครงการศึกษาปัญหาสุขภาพและความต้องการบริการด้านสุขภาพในกลุ่มวัยรุ่น

โครงการนี้มีวัตถุประสงค์เพื่อศึกษาถึงปัญหาด้านสุขภาพและการไปใช้บริการสุขภาพของวัยรุ่น ณ สถานบริการสุขภาพต่างๆทั้งภาครัฐ เอกชน ตลอดจนศึกษาถึง รูปแบบบริการสุขภาพที่สอดคล้องกับความต้องการของวัยรุ่น นอกจากนี้ยังศึกษาถึงคุณภาพชีวิตและภาวะซึมเศร้าของวัยรุ่นด้วย

#### A. แบบสอบถามชุดนี้แบ่งออกเป็น 5 ส่วน ประกอบด้วย

ส่วนที่ 1 ข้อมูลทั่วไปของผู้ตอบแบบสอบถาม

ส่วนที่ 2 ข้อมูลด้านความต้องการบริการสุขภาพและสถานบริการสุขภาพของวัยรุ่น

ส่วนที่ 3 การเข้าถึงบริการสุขภาพของวัยรุ่น

ส่วนที่ 4 คุณภาพชีวิตของวัยรุ่น

ส่วนที่ 5 ภาวะซึมเศร้าของวัยรุ่น

#### B. ผู้ตอบแบบสอบถาม

วัยรุ่น

เจ้าหน้าที่สถานและกรอกคำตอบที่วัยรุ่นตอบ

#### C. ผู้ตรวจทานแบบสอบถาม

(ชื่อ - สกุล..... นวสิทธิ์ สุจริตธรรม โนนโศภิต)

ความร่วมมือในการตอบแบบสอบถาม

กรุณาตอบคำถามทุกข้อให้ครบถ้วน และกาเครื่องหมาย (✓) ที่ตรงกับเป็นจริงสำหรับตัวท่านหรือความเห็นของท่านมากที่สุด ข้อมูลนี้ผู้วิจัยจะนำไปวิเคราะห์ในลักษณะกลุ่ม และจะเก็บเป็นความลับ ไม่นำไปเปิดเผยในที่ใดๆ เป็นรายบุคคล และจะนำเสนอผลการศึกษามาในเชิงนโยบายเพื่อส่งเสริมสุขภาพวัยรุ่นของประเทศ หากท่านยินดีให้ความร่วมมือกรุณาเซ็นชื่อ

(ลายเซ็น)

บ้านเลขที่.....

**MALE ONLY**

31 The following items in column (a), I would like to ask you opinion..... in column (b), (c), (d) and (f), please answer as many choices as you think are appropriated

**Column b :** Whether you have had experienced with .....

**Column c :** Whether you have problems ?

**Column d :** Who would you talk to in your family about the problem? ( can answer more than 1 )

**Column e :** Who you would talk to outside of your family about the problem? ( can answer more than 1 )

**Column f :** What health care facility you would like visit about the problem? ( can answer more than 1 )

**Table 3**

No.	(a) Items	(b) Do you ever have experience with.... ?				(c) Do you have the problem?		(d) Who would they talk to?						(e) mark X only one in these columns What health care facility you would like to visit?					
								(d) People in your family			(e) People outside your family			Visit to					
								parent	sibling	peer	youth leader	community leader	community health volunteer	housewife group	gov. hospital	private hospital	health center	private clinic	DPF
<b>1</b>	<b>Sexuality and reproductive health</b>																		
1.1	<i>Development of sexual organs ; e.g. too small or too big of genital organ</i>	Y	N	Y	N														
1.2	<i>Acne</i>	Y	N	Y	N														
1.3	<i>Wet dream</i>	Y	N	Y	N														
1.4	<i>Hygienic care for genital organ</i>	Y	N	Y	N														
1.5	<i>Frequency of arousal</i>	Y	N	Y	N														
1.6	<i>Masturbation</i>	Y	N	Y	N														
1.9	<i>Having sexual Intercourse</i>	Y	N	Y	N														
	-with girl friend	Y	N	Y	N														
	-with sex worker or bar girl	Y	N	Y	N														
	-with acquaintance	Y	N	Y	N														
	- with boy friend	Y	N	Y	N														
1.10	<i>Having unsafe sex</i>	Y	N	Y	N														
1.11	<i>Family planning</i>																		
	-ask sex partner for take contraceptive pill to prevent unintended pregnancy	Y	N	Y	N														
	-condom used	Y	N	Y	N														
	- others specify _____	Y	N	Y	N														
1.12	<i>Unintended pregnancy of sex partner</i>	Y	N	Y	N														
1.13	<i>Abortion of sex partner</i>	Y	N	Y	N														

## แนวแบบสัมภาษณ์เจาะลึก In-depth interview guidelines

เจ้าหน้าที่สาธารณสุข เจ้าหน้าที่มูลนิธิดวงประทีป อาสาสมัครสาธารณสุข ผู้นำชุมชน ผู้นำวัยรุ่น ผู้นำกลุ่มสตรี / แม่บ้าน

Health officers, NGOs staff, Health Volunteer, Comm. Leaders, youth leaders, Chairman of housewife group/women group ( at least 3 persons in each group)

### 1. ข้อมูลประชากร ( Demographic data)

- เพศ gender
- อายุ age
- การศึกษา education
- ระยะเวลาที่ทำงานอยู่ในชุมชน duration of working in community
- ตำแหน่ง/หน้าที่ position
- ประสบการณ์การทำงาน work experiences
- หน้าที่ความรับผิดชอบ job responsibility

### 2. ความคิดเห็นเกี่ยวกับความต้องการด้านสุขภาพและการเข้าถึงบริการของวัยรุ่นในชุมชน

Opinion about adolescents' health needs and accessibility

### 3. ท่านคิดว่าวัยรุ่นในชุมชนมีปัญหาอะไรบ้าง ทำไมถึงคิดเช่นนั้น

Opinion : Problems of adolescents in this community are ....., why is it a problem ?

### 4. ท่านคิดว่าอนามัยเจริญพันธุ์ในกลุ่มวัยรุ่นเป็นปัญหาหรือไม่ เพราะเหตุใด

Are there any problem about adolescents' reproductive health in this community, why you think it is a problem?

### 5. ท่าน/หน่วยงานของท่านมีบทบาทอย่างไรในเรื่องการทำงานด้านสุขภาพของวัยรุ่น

You/Your organization take any role on adolescents' health

### 6. วัยรุ่นที่มาใช้บริการในหน่วยงานของท่านส่วนใหญ่คือวัยรุ่นกลุ่มใด มีปัญหาเรื่องใดมากที่สุด

Which group of adolescents that usually visit/utilize your organization, what are the problem of ado?

### 7. วัยรุ่นที่ไม่มาใช้บริการในหน่วยงานของท่านส่วนใหญ่คือวัยรุ่นกลุ่มใด น่าจะมีปัญหาเรื่องใดบ้าง

Which group of adolescents never been visited/utilize any kind of services at your organization? Can you guess , what are problems that they might be faced ?

### 8. ท่านคิดว่าบริการที่หน่วยงานของท่านมีอยู่ในปัจจุบันนี้สามารถตอบสนองความต้องการด้านสุขภาพของวัยรุ่น มากน้อยเพียงใด ทำไมถึงคิดเช่นนั้น

Are your organization provide the services to server the needs of ado? ( more or less )? And Why?

### 9. การให้บริการด้านสุขภาพแก่วัยรุ่นในชุมชนควรเป็นเช่นใด

Which kind of adolescents' health services should be provide in this community ?

10. ท่านคิดว่าการทำงานร่วมกัน ( Partnerships)จะช่วยแก้ปัญหาเรื่องของคุณภาพวัยรุ่นได้หรือไม่  
เพราะเหตุใด

Is working as "partnerships" can solve Adolescents' health problem? Why?

11. บุคคลหรือหน่วยงานใดที่ท่านอยากร่วมงานด้วย ในเรื่องการแก้ปัญหาสุขภาพวัยรุ่น  
( บอกชื่อ มากกว่า 1 หน่วยงาน )

Suggestion /nomination the organizations for a partnership for working to gather on adolescents'  
health ( more than 1)

## แนวทางการสนทนากลุ่ม FGD guidelines

วัยรุ่นหญิง/ชาย ที่อยู่ในระบบการศึกษาและนอกระบบการศึกษา

Male & Female in school

- Early ado ( 10 – 13 yrs) : Grade 5 - 6
- Intermediate Ado ( 14- 17 yrs) : Grade 7 -9
- Late ado ( 18 –22 yrs.) : Grade 10 upward

Male & female out school

( same as above)

### 1. ข้อมูลประชากร ( Demographic data)

- เพศ gender
- อายุ age , BD
- การศึกษา Edu
- ระยะเวลาที่อยู่ในชุมชน duration of living in community
- ความสัมพันธ์ในครอบครัว Family relationships
  - Parent
  - Sibling

### 2. ท่านคิดว่าท่านมีปัญหาทั่วไปในเรื่องใดบ้าง

Do you have any problems ( general)?

### 3. เมื่อมีปัญหาปรึกษาใคร เพราะอะไร

Whom you turn to when you have problems? Why?

### 4. ท่านมีปัญหาสุขภาพในเรื่องใดบ้าง

Do you have any health problem?

### 5. เมื่อมีปัญหาสุขภาพปรึกษาใครหรือปฏิบัติอย่างไร เพราะอะไร

Whom you turn to when you have problems? Why?

### 6. ท่านทราบหรือไม่ว่าชุมชนของท่านมีบริการด้านสุขภาพของวัยรุ่นหรือไม่ อยู่ที่ใดบ้าง?

Have you ever known that there are any adolescent health services in this community? If yes, what it it and where?

### 7. ท่านเคยไปใช้บริการด้านสุขภาพวัยรุ่นที่ใดบ้าง

Have you ever use /utilize the above services? Where?

### 8. ท่านพอใจกับบริการสุขภาพที่มีอยู่หรือไม่ เพราะอะไร

Are you satisfy with the above services? Why?



## Appendix C

### General Characteristic's Table

#### 1. Gender and stage of adolescence (age)

**Table C.1 :** Gender and stage of adolescence (age)

Stage of adolescence	Gender		Total
	Male	Female	
Early adolescent	73	130	203
(12-13 years)	19.6%	26.1%	23.3%
Mid adolescent	198	222	420
(14-17 years)	53.1%	44.6%	48.2%
Late adolescent	102	146	248
(18-22 years)	27.3%	29.3%	28.5%
Total	373	498	871
	100.0%	100.0%	100.0%
% within gender	42.8%	57.2%	100.0%

Difference Early/Mid/Late

P value = .025

#### 2. Marital status

**Table C.2 :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age), by Gender and by Adolescents' Marital Status

Adolescent marital status	Male				Female				M & F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
Single	73	193	96	362	128	213	117	458	201	406	213	820
	(100.0)	(98.5)	(95.0)	(97.8)	(99.2)	(96.4)	(80.7)	(92.5)	(99.5)	(97.4)	(86.6)	(94.8)
Ever Married	0	3	5	8	1	8	28	37	1	11	33	45
	(0)	(1.0)	(4.0)	(1.7)	(0.8)	(3.2)	(17.9)	(6.9)	(0.5)	(2.4)	(12.6)	(4.8)
Total	73	196	101	370	129	221	145	495	202	417	246	865
	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)
Missing % missing	0	2	1	3	1	1	1	3	1	3	2	6
	0	1	1	0.8	1	0.4	1	0.6	0.5	0.7	1	0.7

Difference Male vs Female

P value < .001

Difference Early/Mid/Late

P value < .001



### 3. Working status

**Table C.3 :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age), by Gender and by Adolescents' Working Status

Adolescents' working status	Male				Female				M & F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
Not working	71 (100.0)	159 (80.7)	46 (45.5)	276 (74.8)	129 (99.2)	204 (92.7)	95 (65.1)	428 (86.3)	200 (99.5)	363 (87.1)	141 (57.1)	704 (81.4)
Working (full time & part time)	0 (0)	38 (19.3)	55 (54.5)	93 (25.2)	1 (0.8)	16 (7.3)	51 (34.9)	68 (13.7)	1 (0.5)	54 (12.9)	106 (42.9)	161 (18.6)
Total	71 (100.0)	197 (100.0)	101 (100.0)	369 (100.0)	130 (100.0)	220 (100.0)	146 (100.0)	496 (100.0)	201 (100.0)	417 (100.0)	247 (100.0)	865 (100.0)
Missing %missing	2 (2.7)	1 (1)	1 (1)	4 (1)	0 (0)	2 (1)	0 (0)	2 (0.4)	2 (1)	3 (0.7)	1 (0.4)	6 (0.7)

Difference Male vs Female P value < .001

Difference Early/Mid/Late P value < .001

### 4. Home town

**Table C.4 :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age), by Home Town

Home town	Male				Female				M & F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
Bangkok	72 (98.6)	181 (91.9)	88 (86.3)	341 (91.7)	122 (93.8)	203 (91.4)	122 (84.7)	447 (90.1)	194 (95.6)	384 (91.6)	210 (85.4)	788 (90.8)
Non-Bangkok	1 (1.4)	16 (8.1)	14 (13.7)	31 (8.3)	8 (6.2)	19 (8.6)	22 (15.3)	49 (9.9)	9 (4.4)	35 (8.4)	36 (14.6)	80 (9.2)
Total	73 (100.0)	197 (100.0)	102 (100.0)	372 (100.0)	130 (100.0)	222 (100.0)	144 (100.0)	496 (100.0)	203 (100.0)	419 (100.0)	246 (100.0)	868 (100.0)
Missing %missing	0 (0)	1 (1)	0 (0)	1 (0.3)	0 (0)	0 (0)	2 (1.5)	2 (0)	0 (0)	1 (0)	2 (1)	3 (0.7)

Difference Early/Mid/Late P value = .001

## 5. Household income

**Table C.5 :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age), by Gender and by Monthly Household Income

Household income	Male				Female				M & F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
Known	37	124	70	231	77	144	115	336	114	268	185	567
	(51.4)	(63.2)	(68.6)	(62.5)	(59.7)	(65.2)	(78.7)	(67.7)	(56.8)	(64.2)	(74.6)	(65.5)
Don't know	35	72	32	139	52	77	31	160	87	149	63	299
	(48.6)	(36.7)	(31.4)	(37.6)	(40.3)	(34.8)	(21.2)	(32.3)	(43.3)	(35.7)	(25.4)	(34.5)
Total	72	196	102	370	129	221	146	496	201	417	248	866
	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)
Missing %missing	1	2	0	3	1	1	0	2	2	3	0	5
	(1)	(1)	(0)	(0.3)	(1)	(0.5)	(0)	(0.4)	(1)	(0.7)	(0)	(0.6)

Difference Male vs Female

P value < .001

Difference Early/Mid/Late

P value < .001

## 6. Household expenses

**Table C.6 :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age), by Gender and by household expenses

Household expenses	Male				Female				M & F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
Left for saving Saving	14	24	11	49	15	28	14	57	29	52	25	106
	(19.2)	(12.1)	(10.8)	(13.1)	(11.5)	(12.6)	(9.6)	(11.4)	(14.3)	(12.4)	(10.1)	(12.2)
Just enough Enough	23	57	38	118	34	41	43	118	57	98	81	236
	(31.5)	(28.8)	(37.3)	(31.6)	(26.2)	(18.5)	(29.5)	(23.7)	(28.1)	(23.3)	(32.7)	(27.1)
Sometime not enough not enough	15	41	27	83	31	68	45	144	46	109	72	227
	(20.5)	(20.7)	(26.5)	(22.3)	(23.8)	(30.6)	(30.8)	(28.9)	(22.7)	(26.0)	(29.0)	(26.1)
Not enough in debt	4	16	5	25	9	26	20	55	13	42	25	80
	(5.5)	(8.1)	(4.9)	(6.7)	(6.9)	(11.7)	(13.7)	(11.0)	(6.4)	(10.0)	(10.1)	(9.2)
Don't know	17	60	21	98	41	59	24	124	58	119	45	222
	(23.3)	(30.3)	(20.6)	(26.3)	(31.5)	(26.6)	(16.4)	(24.9)	(28.6)	(28.3)	(18.1)	(25.5)
Total	73	198	102	373	130	222	146	498	203	420	248	871
	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)

Difference Early/Mid/Late

P value = .007



**C) Consult with parent/relative when they had problems**

**Table C.7C :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age), by Gender and by consult with parent/relative when they has problem

Consult with parent/ relative when having problem	Male				Female				M & F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
Not at all	1	18	17	36	5	14	8	27	6	32	25	63
	(1.4)	(9.2)	(16.8)	(9.8)	(3.8)	(6.3)	(5.5)	(5.4)	(3.0)	(7.7)	(10.1)	(7.3)
Some	29	67	38	134	45	92	47	184	74	159	85	318
	(40.3)	(34.2)	(37.6)	(36.3)	(34.6)	(41.4)	(32.2)	(36.9)	(36.6)	(38.0)	(34.4)	(36.7)
Occasionally	27	62	28	117	40	64	54	158	67	126	82	275
	(37.5)	(31.6)	(27.7)	(31.7)	(30.8)	(28.8)	(37.0)	(31.7)	(33.2)	(30.1)	(33.2)	(31.7)
Most of the time	15	49	18	82	40	52	37	129	55	101	55	211
	(20.8)	(25.0)	(17.8)	(22.2)	(30.8)	(23.4)	(25.3)	(25.9)	(27.2)	(24.2)	(22.3)	(24.3)
Total	72	196	101	369	130	222	146	498	202	418	247	867
	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)
Missing	1	2	1	3	0	0	0	0	1	2	1	4
%missing	(1)	(1)	(1)	(0.8)	(0)	(0)	(0)	(0)	(0.5)	(0.5)	(0.4)	(0.4)

**D) Having things to do with family**

**Table C.7D :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age), by Gender and by having things to do with family

Having things to do with family	Male				Female				M & F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
Not at all	5	23	19	47	16	34	20	70	21	57	39	117
	(6.8)	(11.7)	(19.0)	(12.7)	(12.3)	(15.3)	(13.7)	(14.1)	(10.3)	(13.6)	(15.9)	(13.5)
Some	38	105	64	207	67	109	73	249	105	214	137	456
	(52.1)	(53.3)	(64.0)	(55.9)	(51.5)	(49.1)	(50.0)	(50.0)	(1.7)	(1.1)	(5.7)	(2.5)
Occasionally	21	45	12	78	28	58	32	118	49	103	44	196
	(28.8)	(22.8)	(12.0)	(21.1)	(21.5)	(26.1)	(21.9)	(23.7)	(24.1)	(24.6)	(17.9)	(22.6)
Most of the time	9	24	5	38	19	21	21	61	28	45	26	99
	(12.3)	(12.2)	(5.0)	(10.3)	(14.6)	(9.5)	(14.4)	(12.2)	(13.8)	(10.7)	(10.6)	(11.4)
Total	73	197	100	370	130	222	146	498	203	419	246	868
	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)
Missing	0	1	2	3	0	0	0	0	0	1	2	3
%missing	(0)	(0)	(2)	(0.8)	(0)	(0)	(0)	(0)	(0)	(0.2)	(1)	(0)

**E) Quarrel with family**

**Table C.7E :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age), by Gender and by Quarrel with family

Quarrel with family	Male				Female				M & F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
Not at all	33 (45.8)	56 (29.2)	44 (44.0)	133 (36.5)	42 (33.3)	64 (29.2)	43 (29.5)	149 (30.3)	75 (37.9)	120 (29.2)	87 (35.4)	282 (33.0)
Some	32 (44.4)	107 (55.7)	43 (43.0)	182 (50.0)	59 (46.8)	113 (51.6)	86 (58.9)	258 (52.5)	91 (46.0)	220 (53.5)	129 (52.4)	440 (51.5)
Occasionally	5 (6.9)	24 (12.5)	11 (11.0)	40 (11.0)	22 (17.5)	40 (18.3)	16 (11.0)	78 (15.9)	27 (13.6)	64 (15.6)	27 (11.0)	118 (13.8)
Most of the time	2 (2.8)	5 (2.6)	2 (2.0)	9 (2.5)	3 (2.4)	2 (0.9)	1 (0.7)	6 (1.2)	5 (2.5)	7 (1.7)	3 (1.2)	15 (1.8)
Total	72 (100.0)	192 (100.0)	100 (100.0)	364 (100.0)	126 (100.0)	219 (100.0)	146 (100.0)	491 (100.0)	198 (100.0)	411 (100.0)	246 (100.0)	855 (100.0)
Missing	1	6	2	9	4	3	0	7	5	9	2	16
%missing	(1)	(3)	(2)	(2.5)	(2)	(1.5)	(0)	(1.5)	(2.5)	(2)	(1)	(1.9)

**7. Coefficient of Family Relationships**

**Table C.7F :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age), by Gender and the Level of Family Relationships

Family relationships	Male				Female				M&F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
Poor (2-8)	13 (17.7)	39 (19.6)	31 (30.4)	83 (22.3)	26 (20.0)	35 (15.9)	20 (13.7)	81 (16.2)	39 (19.3)	74 (17.6)	51 (20.4)	164 (18.8)
Moderate (9-13)	49 (67.1)	111 (56.0)	59 (57.8)	219 (58.6)	73 (56.2)	141 (63.6)	89 (61.0)	303 (60.8)	122 (60.1)	252 (60.1)	148 (59.7)	522 (59.9)
High (13 over)	11 (15.0)	48 (24.2)	12 (11.8)	71 (19.0)	31 (23.8)	46 (20.8)	37 (25.3)	114 (22.8)	42 (20.6)	94 (22.5)	49 (19.7)	185 (21.3)
Total	73 (100.0)	198 (100.0)	102 (100.0)	373 (100.0)	130 (100.0)	222 (100.0)	146 (100.0)	498 (100.0)	203 (100.0)	420 (100.0)	248 (100.0)	871 (100.0)

Difference Males vs Females in Late adolescent

P value = .001



**B) Getting Along with Friend**

**Table C.9B :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by Getting Along with Friend

Getting along with friend	Male				Female				M & F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
Not at all	6 (8.2)	5 (2.5)	5 (5.0)	16 (4.3)	8 (6.2)	5 (2.3)	1 (0.7)	14 (2.8)	14 (6.9)	10 (2.4)	6 (2.4)	30 (3.5)
Some	8 (11.0)	15 (7.6)	12 (11.9)	35 (9.4)	8 (6.2)	12 (5.4)	10 (6.9)	30 (6.1)	16 (7.9)	27 (6.4)	22 (9.0)	65 (7.5)
Occasionally	26 (35.6)	75 (38.1)	29 (28.7)	130 (35.0)	47 (36.4)	69 (31.1)	46 (31.9)	162 (32.7)	73 (36.1)	144 (34.4)	75 (30.6)	292 (33.7)
Most of the Time	33 (45.2)	102 (51.8)	55 (54.5)	190 (51.2)	66 (51.2)	136 (61.3)	87 (60.4)	289 (58.4)	99 (49.0)	238 (56.8)	142 (58.0)	479 (55.3)
Total	73 (100.0)	197 (100.0)	101 (100.0)	371 (100.0)	129 (100.0)	222 (100.0)	144 (100.0)	495 (100.0)	202 (100.0)	419 (100.0)	245 (100.0)	866 (100.0)
Missing %missing	0 (0)	1 (0.5)	1 (1)	2 (0.5)	1 (1)	0 (0)	2 (1.5)	3 (0)	1 (0.5)	1 (0.2)	3 (1.5)	5 (0.6)

**C) Consult with friend when they had problem**

**Table C.9C:** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by Consultation with friend when having problem

Consultation with friend when having problem	Male				Female				M & F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
Not at all	7 (9.6)	13 (6.7)	14 (13.7)	34 (9.2)	9 (6.9)	8 (3.6)	13 (9.0)	30 (6.0)	16 (7.9)	21 (5.0)	27 (10.9)	64 (7.4)
Some	42 (57.5)	83 (42.6)	43 (42.2)	168 (45.4)	39 (30.0)	70 (31.5)	50 (34.5)	159 (32.0)	81 (39.9)	153 (36.7)	93 (37.7)	327 (37.7)
Occasionally	17 (23.3)	64 (32.8)	29 (28.4)	110 (29.7)	52 (40.0)	75 (33.8)	56 (38.6)	183 (36.8)	69 (34.0)	139 (33.3)	85 (34.4)	293 (33.8)
Most of the Time	7 (9.6)	35 (17.9)	16 (15.7)	58 (15.7)	30 (23.1)	69 (31.1)	26 (17.9)	125 (25.2)	37 (18.2)	104 (24.9)	42 (17.0)	183 (21.1)
Total	73 (100.0)	195 (100.0)	102 (100.0)	370 (100.0)	130 (100.0)	222 (100.0)	145 (100.0)	497 (100.0)	203 (100.0)	417 (100.0)	247 (100.0)	867 (100.0)
Missing %missing	0 (0)	3 (2)	0 (0)	3 (0.8)	0 (0)	0 (0)	1 (1)	1 (0.2)	0 (0)	3 (0.7)	1 (0.4)	4 (0.5)

**D) Having things to do with friend**

**Table C.9D :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by Having thing to do with friend

Having thing to do with friend	Male				Female				M & F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
Not at all	14	12	13	39	18	25	18	61	32	37	31	100
	(19.2)	(6.1)	(12.7)	(10.5)	(14.1)	(11.3)	(12.3)	(12.3)	(15.9)	(8.8)	(12.5)	(11.5)
Some	32	77	31	140	49	67	55	171	81	144	86	311
	(43.8)	(39.1)	(30.4)	(37.6)	(38.3)	(30.2)	(37.7)	(34.5)	(40.3)	(34.4)	(34.7)	(35.8)
Occasionally	22	75	40	137	34	77	44	155	56	152	84	292
	(30.1)	(38.1)	(39.2)	(36.8)	(26.6)	(34.7)	(30.1)	(31.3)	(27.9)	(36.3)	(33.9)	(33.6)
Most of the Time	5	33	18	56	27	53	29	109	32	86	47	165
	(6.8)	(16.8)	(17.6)	(15.1)	(21.1)	(23.9)	(19.9)	(22.0)	(15.9)	(20.5)	(19.0)	(19.0)
Total	73	197	102	372	128	222	146	496	201	419	248	868
	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)
Missing	0	1	0	1	2	0	0	2	2	1	0	3
%missing	(0)	(0.5)	(0)	(0.3)	(1.5)	(0)	(0)	(0.4)	(1)	(0.2)	(0)	(0.3)

**E) Quarrel with friend**

**Table C.9E :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by Quarrel with friend

Quarrel with friend	Male				Female				M & F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
Not at all	19	63	53	135	50	80	69	199	69	143	122	334
	(26.0)	(32.0)	(52.0)	(36.3)	(38.8)	(36.2)	(47.3)	(40.1)	(34.2)	(34.2)	(49.2)	(38.5)
Some	43	107	42	192	58	119	67	244	101	226	109	436
	(58.9)	(54.3)	(41.2)	(51.6)	(45.0)	(53.8)	(45.9)	(49.2)	(50.0)	(54.1)	(44.0)	(50.2)
Occasionally	9	22	5	36	15	19	9	43	24	41	14	79
	(12.3)	(11.2)	(4.9)	(9.7)	(11.6)	(8.6)	(6.2)	(8.7)	(11.9)	(9.8)	(5.6)	(9.1)
Most of the Time	2	5	2	9	6	3	1	10	8	8	3	19
	(2.7)	(2.5)	(2.0)	(2.4)	(4.7)	(1.4)	(0.7)	(2.0)	(4.0)	(1.9)	(1.2)	(2.2)
Total	73	197	102	372	129	221	146	496	202	418	248	868
	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)
Missing	0	1	0	1	1	1	0	2	1	2	0	3
%missing	(0)	(0.5)	(0)	(0.3)	(1)	(0.5)	(0)	(0.4)	(0.5)	(1)	(0)	(0.3)



**F) Coefficient of Peer Relationships**

**Table C.9F : Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by the Level of Peer Relationships**

Peer relationship	Male				Female				M &F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
Poor	14 (19.1)	28 (14.1)	28 (27.3)	70 (18.7)	20 (15.3)	25 (11.3)	30 (20.6)	75 (15.0)	34 (16.7)	53 (12.7)	58 (23.4)	145 (16.6)
Moderate	57 (78.1)	131 (66.3)	62 (60.8)	250 (67.0)	84 (64.5)	146 (65.9)	97 (66.5)	327 (65.6)	141 (69.4)	277 (66.0)	159 (64.2)	577 (66.2)
High	2 (2.7)	39 (19.7)	12 (11.8)	53 (14.2)	26 (20.0)	51 (23.0)	19 (13.1)	96 (19.2)	28 (13.8)	90 (21.5)	31 (12.4)	149 (17.2)
Total	73 (100.0)	198 (100.0)	102 (100.0)	373 (100.0)	130 (100.0)	222 (100.0)	146 (100.0)	498 (100.0)	203 (100.0)	420 (100.0)	248 (100.0)	871 (100.0)

Difference Male vs Female in Early Adolescents

P value = .003

Difference Early/Mid/Late

P value < .001

## Appendix D Depression's Table

**Table D.1 (5.77) :** Distribution analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by I was bothered by things that usually don't bother me

I was bothered by things that usually Don't bother me	Males				Females				M & F			Total n %
	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	
Rarely (<1 day)	35 (47.9)	79 (40.1)	31 (30.7)	145 (39.1)	30 (23.1)	31 (14.0)	19 (13.0)	80 (16.1)	65 (32.0)	110 (26.3)	50 (20.2)	225 (25.9)
Sometime (2-4 days)	36 (49.3)	113 (57.3)	67 (66.3)	216 (58.2)	89 (68.5)	177 (80.1)	116 (79.5)	382 (76.9)	125 (61.6)	290 (69.4)	183 (74.1)	598 (68.9)
Most of the time (5-7 days)	2 (2.7)	5 (2.5)	3 (3.0)	10 (2.7)	11 (8.5)	13 (5.9)	11 (7.5)	35 (7.0)	13 (6.4)	18 (4.3)	14 (5.7)	45 (5.2)
Total	73 (100.0)	197 (100.0)	101 (100.0)	371 (100.0)	130 (100.0)	221 (100.0)	146 (100.0)	497 (100.0)	203 (100.0)	418 (100.0)	247 (100.0)	868 (100.0)
Missing %missing	0 (0)	1 (0.5)	1 (1)	2 (0.5)	0 (0)	1 (0.5)	0 (0)	1 (0.2)	0 (0)	2 (0.5)	1 (0.4)	3 (0.3)

Difference (Males vs Female)

P-value = .002

Difference (Males vs Female) in Early Adolescents :

P-value = .001

Difference (Males vs Female) in Mid Adolescents :

P-value = <.001

Difference (Males vs Female) in Late Adolescents :

P-value = .002

**Table D.2 (5.78) :** Distribution Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by I did not feel like eating; my appetite was poor

I did not feel like eating; my appetite was poor	Male				Female				M & F			Total n %
	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	
Rarely	34 (47.2)	107 (54.0)	50 (49.0)	191 (51.3)	51 (39.2)	71 (32.1)	54 (37.0)	176 (35.4)	85 (42.1)	178 (42.5)	104 (41.9)	367 (42.2)
Sometime	34 (47.2)	90 (45.5)	49 (48.0)	173 (46.5)	75 (57.7)	145 (65.6)	89 (60.9)	309 (62.2)	109 (54.0)	235 (56.1)	138 (55.6)	482 (55.5)
Most of the time	4 (5.6)	1 (0.5)	3 (2.9)	8 (2.2)	4 (3.1)	5 (2.3)	3 (2.1)	12 (2.4)	8 (4.0)	6 (1.4)	6 (2.4)	20 (2.3)
Total	72 (100.0)	198 (100.0)	102 (100.0)	372 (100.0)	130 (100.0)	221 (100.0)	146 (100.0)	497 (100.0)	202 (100.0)	419 (100.0)	248 (100.0)	869 (100.0)
Missing %missing	1 (1.4)	0 (0)	0 (0)	1 (0.3)	0 (0)	1 (0.5)	0 (0)	1 (0.2)	1 (0.5)	1 (0.2)	0 (0)	2 (0.1)

Difference (Male vs Female)

P-value = .002

Difference (Male vs Female) in Mid Adolescents :

P-value = <.001

**Table D.3 (5.79) :** Destribution Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by I felt that I could not shake off the blues

I felt that I could not shake off the blues	Male				Female				M & F			Total n %
	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	
Rarely	46 (63.0)	107 (55.2)	50 (49.0)	203 (55.0)	62 (48.1)	94 (42.5)	70 (48.3)	226 (45.7)	108 (53.5)	201 (48.4)	120 (48.6)	429 (49.7)
Sometime	23 (31.5)	80 (41.2)	46 (45.1)	149 (40.4)	58 (44.9)	116 (52.5)	69 (47.6)	243 (49.1)	81 (40.1)	196 (47.2)	115 (46.6)	392 (45.4)
Most of the time	4 (5.5)	7 (3.6)	6 (5.9)	17 (4.6)	9 (7.0)	11 (5.0)	6 (4.1)	26 (5.3)	13 (6.4)	18 (4.3)	12 (4.9)	43 (5.0)
Total	73 (100.0)	194 (100.0)	102 (100.0)	369 (100.0)	129 (100.0)	221 (100.0)	145 (100.0)	495 (100.0)	202 (100.0)	415 (100.0)	247 (100.0)	864 (100.0)
Missing %missing	0 (0)	4 (2)	0 (0)	4 (1.1)	1 (1)	1 (0.5)	1 (1)	3 (0.6)	1 (0.5)	5 (1.2)	1 (0.4)	7 (0.8)

**Table D.4 (5.80) :** Distribution Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by I felt that I was just as good as other people

I felt that I was just as good as other people	Male				Female				M & F			Total n %
	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	
Rarely	15 (20.8)	39 (19.8)	8 (7.8)	62 (16.7)	20 (15.4)	26 (11.9)	12 (8.2)	58 (11.7)	35 (17.3)	65 (15.6)	20 (8.1)	120 (13.9)
Sometime	38 (52.7)	99 (50.2)	61 (59.8)	198 (53.4)	77 (59.2)	120 (54.8)	89 (61.0)	286 (57.8)	115 (56.9)	219 (52.6)	150 (60.5)	484 (55.9)
Most of the time	19 (26.4)	59 (29.9)	33 (32.4)	111 (29.9)	33 (25.4)	73 (33.3)	45 (30.8)	151 (30.5)	52 (25.7)	132 (31.7)	78 (31.5)	262 (30.3)
Total	72 (100.0)	197 (100.0)	102 (100.0)	371 (100.0)	130 (100.0)	219 (100.0)	146 (100.0)	495 (100.0)	202 (100.0)	416 (100.0)	248 (100.0)	866 (100.0)
Missing %missing	1 (1.4)	1 (0.5)	0 (0)	2 (0)	0 (0)	3 (1.5)	0 (0)	3 (0.2)	1 (0.2)	4 (1)	0 (0)	5 (0.6)

Difference (Early/Mid/Late Adolescents) :

P-value = .016

**Table D.5 (5.81) :** Destibution Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by I had trouble keeping my mind on what I was doing

I had trouble keeping my mind on what I was doing	Male				Female				M & F			Total
	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	N %
Rarely	22 (30.1)	62 (31.3)	27 (26.7)	111 (29.8)	33 (25.6)	45 (20.3)	32 (22.1)	110 (22.1)	55 (27.2)	107 (25.5)	59 (24.0)	221 (25.5)
Sometime	45 (61.7)	128 (64.7)	70 (69.3)	243 (65.3)	90 (69.8)	165 (74.3)	109 (75.1)	364 (73.4)	135 (66.8)	293 (69.8)	179 (72.8)	607 (69.9)
Most of the time	6 (8.2)	8 (4.0)	4 (4.0)	18 (4.8)	6 (4.7)	12 (5.4)	4 (2.8)	22 (4.5)	12 (5.9)	20 (4.8)	8 (3.3)	40 (4.6)
Total	73 (100.0)	198 (100.0)	101 (100.0)	372 (100.0)	129 (100.0)	222 (100.0)	145 (100.0)	496 (100.0)	202 (100.0)	420 (100.0)	246 (100.0)	868 (100.0)
Missing	0 (0)	0 (0)	1 (1)	1 (0.3)	1 (1)	0 (0)	1 (1)	2 (0.4)	1 (0.2)	0 (0.5)	2 (1)	3 (0.3)

**Table D.6 (5.82) :** Distribution Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by I felt depressed

I felt depressed	Male				Female				M & F			Total n %
	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	
Rarely	39 (53.4)	110 (55.6)	47 (46.1)	196 (52.5)	60 (46.9)	95 (42.8)	64 (43.8)	219 (44.2)	99 (49.3)	205 (48.8)	111 (44.8)	415 (47.8)
Sometime	32 (43.9)	88 (44.5)	53 (51.9)	173 (46.4)	62 (48.4)	121 (54.5)	80 (54.8)	263 (53.0)	94 (46.8)	209 (49.8)	133 (53.6)	436 (50.2)
Most of the time	2 (2.7)	0 (0)	2 (2.0)	4 (1.1)	6 (4.7)	6 (2.7)	2 (1.4)	14 (2.8)	8 (4.0)	6 (1.4)	4 (1.6)	18 (2.1)
Total	73 (100.0)	198 (100.0)	102 (100.0)	373 (100.0)	128 (100.0)	222 (100.0)	146 (100.0)	496 (100.0)	201 (100.0)	420 (100.0)	248 (100.0)	869 (100.0)
Missing	0 (0)	0 (0)	0 (0)	0 (0)	2 (1.5)	0 (0)	0 (0)	2 (0.4)	2 (1)	0 (0)	0 (0)	2 (0.2)

Difference (Male vs Female) :

P-value = .017

**Table D.7 (5.83) :** Destitution Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by I felt that everything I did was an effort

I felt that everything I did was an effort	Male				Female				M & F			Total n %
	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	
Rarely	37 (51.4)	108 (54.8)	48 (47.1)	193 (52.0)	58 (44.6)	106 (47.7)	73 (50.3)	237 (47.7)	95 (47.0)	214 (51.1)	121 (49.0)	430 (49.5)
Sometime	30 (41.7)	84 (42.6)	49 (48.0)	163 (43.9)	68 (52.3)	110 (49.5)	69 (47.5)	247 (49.7)	98 (48.5)	194 (46.3)	118 (47.7)	410 (47.3)
Most of the time	5 (6.9)	5 (2.5)	5 (4.9)	15 (4.0)	4 (3.1)	6 (2.7)	3 (2.1)	13 (2.6)	9 (4.5)	11 (2.6)	8 (3.2)	28 (3.2)
Total	72 (100.0)	197 (100.0)	102 (100.0)	371 (100.0)	130 (100.0)	222 (100.0)	145 (100.0)	497 (100.0)	202 (100.0)	419 (100.0)	247 (100.0)	868 (100.0)
Missing %missing	1 (1.4)	1 (0.5)	0 (0)	2 (0.5)	0 (0)	0 (0)	1 (1)	1 (0.2)	1 (0.5)	1 (0.2)	1 (0.4)	3 (0.3)

**Table D.8 (5.84) :** Distribution Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by I felt hopeful about the future

I felt hopeful about the future	Male				Female				M & F			Total n %
	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	
Rarely	20 (27.8)	33 (16.8)	15 (14.9)	68 (18.4)	17 (13.2)	24 (11.0)	14 (9.7)	55 (11.2)	37 (18.4)	57 (13.7)	29 (11.8)	123 (14.3)
Sometime	36 (50.0)	118 (59.9)	77 (76.2)	231 (62.4)	79 (61.3)	139 (63.5)	103 (71.0)	321 (65.1)	115 (57.2)	257 (61.8)	180 (73.2)	552 (64.0)
Most of the time	16 (22.2)	46 (23.4)	9 (8.9)	71 (19.2)	33 (25.6)	56 (25.6)	28 (19.3)	117 (23.7)	49 (24.4)	102 (24.5)	37 (15.0)	188 (21.8)
Total	72 (100.0)	197 (100.0)	101 (100.0)	370 (100.0)	129 (100.0)	219 (100.0)	145 (100.0)	493 (100.0)	201 (100.0)	416 (100.0)	246 (100.0)	863 (100.0)
Missing %missing	1 (1.4)	1 (0.5)	1 (1)	3 (0.8)	1 (1)	3 (1.5)	1 (1)	5 (1)	2 (1)	4 (1)	2 (1)	8 (0.9)

Difference ( Male vs Female) :

P-value = .007

Difference ( Early/Mid/Late Adolescents) :

P-value = .003

**Table D.9 (5.85) :** Distribution Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by I thought my life had been a failure

I thought my life had been a failure	Male				Female				M&F			Total n %
	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	
Rarely	4 (56.9)	116 (58.6)	57 (56.4)	177 (47.7)	62 (47.7)	113 (50.9)	90 (62.1)	265 (53.3)	66 (32.7)	229 (54.5)	147 (59.8)	442 (50.9)
Sometime	30 (41.7)	77 (38.9)	42 (41.6)	149 (40.2)	65 (50.0)	103 (46.4)	52 (35.9)	220 (44.3)	95 (47.0)	180 (42.9)	94 (38.2)	369 (42.5)
Most of the time	1 (1.4)	5 (2.5)	2 (2.0)	8 (2.2)	3 (2.3)	6 (2.7)	3 (2.1)	12 (2.4)	4 (2.0)	11 (2.6)	5 (2.0)	20 (2.3)
Total	72 (100.0)	198 (100.0)	101 (100.0)	371 (100.0)	130 (100.0)	222 (100.0)	145 (100.0)	497 (100.0)	202 (100.0)	420 (100.0)	246 (100.0)	868 (100.0)
Missing %missing	1 (1.4)	0 (0)	1 (1)	2 (0.5)	0 (0)	0 (0)	1 (1)	1 (0.2)	1 (0.5)	0 (0)	2 (1)	3 (0.3)

**Table D.10 (5.86) :** Distribution Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by I felt fearful

I felt fearful	Male				Female				M&F			Total n %		
	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %			
	Rarely	44.0 (60.3)	128.0 (65.0)	67.0 (66.3)	239.0 (64.4)	63.0 (48.8)	125.0 (56.3)	84.0 (57.5)	272.0 (54.7)	107.0 (53.0)	253.0 (60.4)		151.0 (61.1)	511.0 (58.9)
	Sometime	28.0 (38.3)	66.0 (33.5)	32.0 (31.7)	126.0 (34.0)	63.0 (48.9)	92.0 (41.4)	61.0 (41.8)	216.0 (43.5)	91.0 (45.0)	158.0 (37.7)		93.0 (37.7)	342.0 (39.4)
Most of the time	1.0 (1.4)	3.0 (1.5)	2.0 (2.0)	6.0 (1.6)	3.0 (2.3)	5.0 (2.3)	1.0 (0.7)	9.0 (1.8)	4.0 (2.0)	8.0 (1.9)	3.0 (1.2)	15.0 (1.7)		
Total	73.0 (100.0)	197.0 (100.0)	101.0 (100.0)	371.0 (100.0)	129.0 (100.0)	222.0 (100.0)	146.0 (100.0)	497.0 (100.0)	202.0 (100.0)	419.0 (100.0)	247.0 (100.0)	868.0 (100.0)		
Missing %missing	0.0	1.0	1.0	2.0	1.0	0.0	0.0	1.0	1.0	1.0	1.0	3.0		

Difference (Male vs Female)

P-value = .006

**Table D.11 (5.87) : Distribution Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by my sleep was restless**

My sleep was restless	Male				Female				M&F			
	Early	Mid	Late	Total	Early	Mid	Late	Total	Early	Mid	Late	Total
	n	n	n	n	n	n	n	n	n	n	n	n
	%	%	%	%	%	%	%	%	%	%	%	%
Rarely	40.0 (54.8)	91.0 (46.4)	43.0 (42.2)	174.0 (46.9)	58.0 (45.0)	105.0 (47.3)	60.0 (41.1)	223.0 (44.9)	98.0 (48.5)	196.0 (46.9)	103.0 (41.5)	397.0 (45.7)
Sometime	29.0 (39.8)	98.0 (50.0)	55.0 (53.9)	182.0 (49.1)	69.0 (53.5)	112.0 (50.5)	81.0 (55.4)	262.0 (52.7)	98.0 (48.5)	210.0 (50.2)	136.0 (54.8)	444.0 (51.2)
Most of the time	4.0 (5.5)	7.0 (3.6)	4.0 (3.9)	15.0 (4.0)	2.0 (1.6)	5.0 (2.3)	5.0 (3.4)	12.0 (2.4)	6.0 (3.0)	12.0 (2.9)	9.0 (3.6)	27.0 (3.1)
Total	73.0 (100.0)	196.0 (100.0)	102.0 (100.0)	371.0 (100.0)	129.0 (100.0)	222.0 (100.0)	146.0 (100.0)	497.0 (100.0)	202.0 (100.0)	418.0 (100.0)	248.0 (100.0)	868.0 (100.0)
Missing %missing	0.0 0.0	2.0 (1.0)	0.0 0.0	2.0 (0.5)	1.0 (1.0)	0.0 0.0	0.0 0.0	1.0 (0.2)	1.0 (0.5)	2.0 (0.5)	0.0 0.0	3.0 (0.3)

**Table D.12 (5.88) : Distribution Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by I was happy**

I was happy	Male				Female				M&F			
	Early	Mid	Late	Total	Early	Mid	Late	Total	Early	Mid	Late	Total
	n	n	n	n	n	n	n	n	n	n	n	n
	%	%	%	%	%	%	%	%	%	%	%	%
Rarely	6.0 (8.2)	17.0 (8.7)	4.0 (4.0)	27.0 (7.3)	8.0 (6.2)	14.0 (6.4)	11.0 (7.7)	33.0 (6.7)	14.0 (6.9)	31.0 (7.5)	15.0 (6.1)	60.0 (7.0)
Sometime	56.0 (76.7)	148.0 (75.9)	78.0 (77.3)	282.0 (76.4)	97.0 (74.6)	166.0 (75.4)	109.0 (76.3)	372.0 (75.5)	153.0 (75.4)	314.0 (75.7)	187.0 (76.6)	654.0 (75.9)
Most of the time	11.0 (15.1)	30.0 (15.4)	19.0 (18.8)	60.0 (16.3)	25.0 (19.2)	40.0 (18.2)	23.0 (16.1)	88.0 (17.8)	36.0 (17.7)	70.0 (16.9)	42.0 (17.2)	148.0 (17.2)
Total	73.0 (100.0)	195.0 (100.0)	101.0 (100.0)	369.0 (100.0)	130.0 (100.0)	220.0 (100.0)	143.0 (100.0)	493.0 (100.0)	203.0 (100.0)	415.0 (100.0)	244.0 (100.0)	862.0 (100.0)
Missing %missing	0.0 0.0	3.0 (1.5)	1.0 (1.0)	4.0 (1.0)	0.0 0.0	2.0 (1.0)	3.0 (2.0)	5.0 (1.0)	0.0 0.0	5.0 (1.2)	4.0 (2.0)	9.0 (1.0)

**Table D.13 (5.89) : Distribution Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by I talked less than usual talking less than usual**

I talked less than usual talking less than usual	Male				Female				M&F			Total n %
	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	
Rarely	35 (47.9)	105 (53.0)	50 (49.0)	190 (50.9)	70 (53.8)	106 (47.7)	70 (47.9)	246 (49.4)	105 (51.7)	211 (50.2)	120 (48.4)	436 (50.1)
Sometime	36 (49.3)	89 (45.0)	51 (50.0)	176 (47.2)	52 (40.0)	110 (49.6)	72 (49.3)	234 (47.0)	88 (43.3)	199 (47.4)	123 (49.6)	410 (47.1)
Most of the time	2 (2.7)	4 (2.0)	1 (1.0)	7 (1.9)	8 (6.2)	6 (2.7)	4 (2.7)	18 (3.6)	10 (4.9)	10 (2.4)	5 (2.0)	25 (2.9)
Total	73 (100.0)	198 (100.0)	102 (100.0)	373 (100.0)	130 (100.0)	222 (100.0)	146 (100.0)	498 (100.0)	203 (100.0)	420 (100.0)	248 (100.0)	871 (100.0)

**Table D.14 (5.90) : Distribution Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by I felt lonely**

I felt lonely	Male				Female				M&F			Total n %
	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	
Rarely	41 (56.9)	105 (53.6)	49 (48.5)	195 (52.8)	62 (47.7)	83 (37.4)	62 (42.5)	207 (41.6)	103 (51.0)	188 (45.0)	111 (44.9)	402 (46.4)
Sometime	29 (40.2)	89 (45.4)	47 (46.6)	165 (44.7)	61 (47)	128 (57.6)	82 (56.2)	271 (54.5)	90 (44.5)	217 (51.9)	129 (52.3)	436 (50.3)
Most of the time	2 (2.8)	2 (1.0)	5 (5.0)	9 (2.4)	7 (5.4)	11 (5.0)	2 (1.4)	20 (4.0)	9 (4.5)	13 (3.1)	7 (2.8)	29 (3.3)
Total	72 (100.0)	196 (100.0)	101 (100.0)	369 (100.0)	130 (100.0)	222 (100.0)	146 (100.0)	498 (100.0)	202 (100.0)	418 (100.0)	247 (100.0)	867 (100.0)
Missing %missing	1 (1.4)	2 (1)	1 (1)	4 (1)	0 (0)	0 (0)	0 (0)	0 (0)	1 (0.5)	2 (0.4)	1 (0.4)	4 (0.4)

Difference (Male vs Female)

P-value = .003



**Table D.15 (5.91) : Distribution Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by I felt people were unfriendly**

I felt people were unfriendly	Male				Female				M&F			Total n %
	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	
Rarely	46 (63.9)	122 (61.6)	60 (59.4)	228 (61.5)	79 (61.7)	132 (59.7)	94 (64.8)	305 (61.7)	125 (62.5)	254 (60.6)	154 (62.6)	533 (61.6)
Sometime	23 (31.9)	65 (32.8)	36 (35.6)	124 (33.4)	42 (32.8)	79 (35.7)	44 (30.3)	165 (33.4)	65 (32.5)	144 (34.4)	80 (32.5)	289 (33.4)
Most of the time	3 (4.2)	11 (5.6)	5 (5.0)	19 (5.1)	7 (5.5)	10 (4.5)	7 (4.8)	24 (4.9)	10 (5.0)	21 (5.0)	12 (4.9)	43 (5.0)
Total	72 (100.0)	198 (100.0)	101 (100.0)	371 (100.0)	128 (100.0)	221 (100.0)	145 (100.0)	494 (100.0)	200 (100.0)	419 (100.0)	246 (100.0)	865 (100.0)
Missing %missing	1 (1.4)	0 (0)	1 (1)	2 (0.5)	2 (1.5)	1 (0.5)	1 (1)	4 (0.8)	3 (1.5)	1 (0.2)	2 (1)	6 (0.7)

**Table D.16 (5.92) : Destibution Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by I enjoyed life**

I felt people were unfriendly	Male				Female				M&F			Total n %
	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	
Rarely	46 (63.9)	122 (61.6)	60 (59.4)	228 (61.5)	79 (61.7)	132 (59.7)	94 (64.8)	305 (61.7)	125 (62.5)	254 (60.6)	154 (62.6)	533 (61.6)
Sometime	23 (31.9)	65 (32.8)	36 (35.6)	124 (33.4)	42 (32.8)	79 (35.7)	44 (30.3)	165 (33.4)	65 (32.5)	144 (34.4)	80 (32.5)	289 (33.4)
Most of the time	3 (4.2)	11 (5.6)	5 (5.0)	19 (5.1)	7 (5.5)	10 (4.5)	7 (4.8)	24 (4.9)	10 (5.0)	21 (5.0)	12 (4.9)	43 (5.0)
Total	72 (100.0)	198 (100.0)	101 (100.0)	371 (100.0)	128 (100.0)	221 (100.0)	145 (100.0)	494 (100.0)	200 (100.0)	419 (100.0)	246 (100.0)	865 (100.0)
Missing %missing	1 (1.4)	0 (0)	1 (1)	2 (0.5)	2 (1.5)	1 (0.5)	1 (1)	4 (0.8)	3 (1.5)	1 (0.2)	2 (1)	6 (0.7)

**Table D.17 (5.93) : Distribution Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by I had crying spells**

I had crying spells	Male				Female				M&F			Total n %
	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	
Rarely	33 (45.2)	103 (52.8)	60 (58.8)	196 (53)	46 (35.4)	53 (23.9)	53 (36.3)	152 (30.5)	79 (38.9)	156 (37.4)	113 (45.6)	348 (40.1)
Sometime	39 (53.4)	92 (47.2)	41 (40.2)	172 (46.5)	77 (59.2)	163 (73.4)	90 (61.6)	330 (66.3)	116 (57.1)	255 (61.2)	131 (52.8)	502 (57.8)
Most of the time	1 (1.4)	0 (0)	1 (1)	2 (0.5)	7 (5.4)	6 (2.7)	3 (2.1)	16 (3.2)	8 (3.9)	6 (1.4)	4 (1.6)	18 (2.1)
Total	73 (100.0)	195 (100.0)	102 (100.0)	370 (100.0)	130 (100.0)	222 (100.0)	146 (100.0)	498 (100.0)	203 (100.0)	417 (100.0)	248 (100.0)	868 (100.0)
Missing %missing	0 (0)	3 (1.5)	0 (0)	3 (0.8)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	3 (0.7)	0 (0)	3 (0.3)

Difference (Male vs Female)

P-value = &lt; .001

**Table D.18 (5.94) : Distribution Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by Feeling sad**

Feeling sad	Male				Female				M&F			Total n %
	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	
Rarely	32 (45.7)	108 (54.8)	49 (49.0)	189 (51.5)	52 (40.3)	69 (31.4)	57 (39.0)	178 (36.0)	84 (42.2)	177 (42.4)	106 (43.1)	367 (42.6)
Sometime	37 (52.8)	87 (44.2)	48 (48.0)	172 (46.9)	74 (57.3)	145 (65.9)	87 (59.6)	306 (61.8)	111 (55.8)	232 (55.6)	135 (54.9)	478 (55.5)
Most oft the time	1 (1.4)	2 (1.0)	3 (3.0)	6 (1.6)	3 (2.3)	6 (2.7)	2 (1.4)	11 (2.2)	4 (2.0)	8 (1.9)	5 (2.0)	17 (2.0)
Total	70 (100.0)	197 (100.0)	100 (100.0)	367 (100.0)	129 (100.0)	220 (100.0)	146 (100.0)	495 (100.0)	199 (100.0)	417 (100.0)	246 (100.0)	862 (100.0)
Missing %missing	3 (4)	1 (0.5)	2 (2)	6 (1.8)	1 (1)	2 (1)	0 (0)	3 (0.6)	4 (2)	3 (0.7)	2 (1)	9 (1)

Difference (Male vs Female)

P-value = &lt; .001

**Table D.19 (5.95) :** Distribution Analysis of Adolescents: Distribution by Stage of Adolescence (age) by (and) Gender and by I felt that people dislike them

I felt that people dislike them	Male				Female				M&F			Total n %
	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	
Rarely	41 (56.2)	135 (68.5)	64 (63.4)	240 (64.7)	81 (62.3)	126 (57.0)	92 (63.4)	299 (60.3)	122 (60.1)	261 (62.4)	156 (63.4)	539 (62.2)
Sometime	31 (42.5)	60 (30.5)	33 (32.7)	124 (33.4)	45 (34.6)	92 (41.6)	51 (35.1)	188 (37.9)	76 (37.4)	152 (36.4)	84 (34.1)	312 (36.0)
Most of the time	1 (1.4)	2 (1.0)	4 (4.0)	7 (1.9)	4 (3.1)	3 (1.4)	2 (1.4)	9 (1.8)	5 (2.5)	5 (1.2)	6 (2.4)	16 (1.8)
Total	73 (100.0)	197 (100.0)	101 (100.0)	371 (100.0)	130 (100.0)	221 (100.0)	145 (100.0)	496 (100.0)	203 (100.0)	418 (100.0)	246 (100.0)	867 (100.0)
Missing %missing	0 (0)	1 (0.5)	1 (1)	2 (0.5)	0 (0)	1 (0.5)	1 (1)	2 (0.4)	0 (0)	2 (0.2)	2 (1)	4 (0.5)

**Table D.20 (5.96) :** Distribution Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by I couldn't get "going"

I couldn't get "going"	Male				Female				M&F			Total n %
	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	Total n %	Early n %	Mid n %	Late n %	
Rarely	45 (0.6)	134 (0.7)	61 (0.6)	240 (0.6)	79 (0.6)	129 (0.6)	78 (0.5)	286 (0.6)	124 (0.6)	263 (0.6)	139 (0.6)	526 (60.5)
Sometime	25 (34.8)	60 (30.5)	35 (34.3)	120 (32.3)	48 (36.9)	85 (38.3)	65 (44.5)	198 (39.8)	73 (36.1)	145 (34.6)	100 (40.3)	318 (36.6)
Most of the time	2 (2.8)	3 (1.5)	6 (5.9)	11 (3.0)	3 (2.3)	8 (3.6)	3 (2.1)	14 (2.8)	5 (2.5)	11 (2.6)	9 (3.6)	25 (2.9)
Total	72 (100.0)	197 (100.0)	102 (100.0)	371 (100.0)	130 (100.0)	222 (100.0)	146 (100.0)	498 (100.0)	202 (100.0)	419 (100.0)	248 (100.0)	869 (100.0)
Missing %missing	1 (1.4)	1 (0.5)	0 (0.0)	2 (0.5)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	1.0 (0.5)	1.0 (0.2)	0.0 (0.0)	2.0 (0.2)

## Appendix E Quality of Life's Table

### 1. Overall assessment and general health facet

#### A) Rating of QoL

**Table E.1A :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by Rating of quality of life

Rating of QoL	Male				Female				M&F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
Poor	10 (13.6)	15 (48)	8 (7.8)	33 (8.8)	13 (10)	20 (9.1)	7 (4.8)	40 (8)	23 (11.3)	35 (8.3)	15 (6.1)	73 (8.4)
Neither poor nor good	31 (42.5)	102 (51.5)	71 (69.6)	204 (54.7)	67 (51.5)	139 (62.6)	99 (67.8)	305 (61.2)	98 (48.4)	241 (57.4)	170 (68.5)	509 (58.4)
Good	32 (43.8)	81 (40.9)	23 (22.6)	136 (36.46)	50 (38.5)	63 (28.4)	40 (27.4)	153 (30.3)	82 (40.4)	144 (34.3)	63 (25.4)	289 (33.2)
Total	73 (100)	198 (100)	102 (100)	373 (100)	130 (100)	222 (100)	146 (100)	498 (100)	203 (100)	420 (100)	248 (100)	871 (100)

Difference Early/Mid/Late

P value = .001

#### B) Satisfaction with health status

**Table E.1B :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by Satisfaction with their health status

Satisfaction with health status	Male				Female				M&F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
Dissatisfied	6 (8.4)	16 (8.3)	19 (19.0)	41 (11.2)	15 (11.5)	37 (16.9)	26 (17.9)	78 (15.8)	21 (10.4)	53 (12.8)	45 (18.4)	119 (13.8)
Neither dissatisfied nor satisfied	29 (40.3)	67 (34.5)	37 (37.0)	133 (36.3)	48 (36.9)	84 (38.4)	58 (40.0)	190 (38.5)	77 (38.1)	151 (36.6)	95 (38.8)	323 (37.6)
Satisfied	37 (51.3)	111 (57.2)	44 (44.0)	192 (52.5)	67 (51.5)	98 (44.8)	61 (42.1)	226 (45.7)	104 (51.5)	209 (50.6)	105 (42.9)	418 (48.6)
Total	72 (100.0)	194 (100.0)	100 (100.0)	366 (100.0)	130 (100.0)	219 (100.0)	145 (100.0)	494 (100.0)	202 (100.0)	413 (100.0)	245 (100.0)	860 (100.0)
Missing % missing	1 (1.4)	4 (2.0)	2 (2.0)	7 (2.0)	0 (0)	3 (2.2)	1 (1.0)	4 (0.8)	1 (0.5)	7 (1.8)	3 (2.2)	11 (1.5)

## 2. Components of QoL

### A) Physical domain

**Table E.2A1:** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by Feeling that physical pain prevent them from doing what they need to do

Feeling that physical pain prevent them from doing what they need to do	Male				Female				M&F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
Very much	15 (20.6)	37 (18.7)	16 (15.6)	68 (18.3)	17 (13.3)	55 (25.1)	34 (23.6)	106 (21.6)	32 (15.9)	92 (22.1)	50 (20.3)	174 (20.2)
Moderate amount	27 (37.0)	94 (47.7)	44 (43.1)	165 (44.4)	62 (48.4)	83 (37.9)	65 (45.1)	210 (42.8)	89 (44.3)	177 (42.5)	109 (44.3)	375 (43.5)
A little	31 (42.5)	66 (33.5)	42 (41.1)	139 (37.4)	49 (38.3)	81 (37.0)	45 (31.2)	175 (35.6)	80 (39.8)	147 (35.3)	87 (35.4)	314 (36.4)
Total	73 (100.0)	197 (100.0)	102 (100.0)	372 (100.0)	128 (100.0)	219 (100.0)	144 (100.0)	491 (100.0)	201 (100.0)	416 (100.0)	246 (100.0)	863 (100.0)
Missing % missing	0 (0.0)	1 (0.5)	0 (0.0)	1 (0.3)	2 (1.5)	3 (1.5)	2 (1.5)	7 (1.5)	2 (1.0)	4 (1.0)	2 (1.0)	8 (1.8)

**Table E.2A2 :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by Having enough energy for everyday life

Having enough energy for everyday life	Male				Female				M&F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
A little	16 (21.9)	25 (12.8)	11 (10.8)	52 (14.1)	25 (19.2)	32 (14.4)	15 (10.3)	72 (14.5)	41 (20.2)	57 (13.7)	26 (10.5)	124 (14.3)
Moderate amount	23 (31.5)	78 (40.0)	44 (43.1)	145 (39.2)	48 (36.9)	96 (43.4)	61 (42.1)	205 (41.3)	71 (35.0)	174 (41.8)	105 (42.5)	350 (40.4)
Very much	34 (46.6)	92 (47.1)	47 (46.1)	173 (46.8)	57 (43.8)	93 (42.0)	69 (47.6)	219 (44.2)	91 (44.8)	185 (44.5)	116 (47.0)	392 (45.3)
Total	73 (100.0)	195 (100.0)	102 (100.0)	370 (100.0)	130 (100.0)	221 (100.0)	145 (100.0)	496 (100.0)	203 (100.0)	416 (100.0)	247 (100.0)	866 (100.0)
Missing % missing	0 (0.0)	3 (1.5)	0 (0.0)	3 (0.8)	0 (0.0)	1 (0.5)	1 (1.0)	2 (0.4)	0 (0.0)	4 (1.0)	1 (0.4)	5 (0.6)

**Table E.2A3 :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by Satisfaction with their sleep

Satisfaction with their sleep	Male				Female				M&F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
Dissatisfied	8 (11.1)	22 (11.1)	21 (20.8)	51 (13.7)	28 (21.8)	37 (16.8)	16 (11.1)	81 (16.4)	36 (17.9)	59 (14.1)	37 (15.0)	132 (15.3)
Neither satisfied nor dissatisfied	32 (44.4)	59 (29.8)	27 (26.7)	118 (31.8)	30 (23.3)	81 (36.8)	53 (36.6)	164 (33.2)	62 (30.8)	140 (33.5)	80 (32.5)	282 (32.6)
Satisfied	32 (44.5)	117 (59.1)	53 (52.4)	202 (54.4)	71 (55.1)	102 (46.3)	76 (52.4)	249 (50.4)	103 (51.2)	219 (52.4)	129 (52.4)	451 (52.1)
Satisfied	20 (27.8)	87 (43.9)	36 (35.6)	143 (38.5)	49 (38.0)	72 (32.7)	60 (41.4)	181 (36.6)	69 (34.3)	159 (38.0)	96 (39.0)	324 (37.5)
Total	72 (100.0)	198 (100.0)	101 (100.0)	371 (100.0)	129 (100.0)	220 (100.0)	145 (100.0)	494 (100.0)	201 (100.0)	418 (100.0)	246 (100.0)	865 (100.0)
Missing % missing	1 (1.4)	0 (0.0)	1 (1.0)	2 (0.5)	1 (1.0)	2 (1.0)	1 (1.0)	4 (0.8)	2 (1.0)	2 (0.5)	2 (1.0)	6 (0.7)

Difference Male vs Female in Early Adolescents P value = .014

**Table E.2A4 :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by Satisfaction with their ability to perform their daily living activities

Satisfaction with their ability to perform their daily living activities	Male				Female				M&F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
Dissatisfied	17 (23.6)	28 (14.2)	25 (25.0)	70 (19.0)	25 (19.4)	32 (14.5)	16 (11.1)	73 (14.7)	42 (20.9)	60 (14.4)	41 (16.7)	143 (16.6)
Neither satisfied nor dissatisfied	29 (40.3)	91 (46.2)	41 (41.0)	161 (43.6)	46 (35.7)	112 (50.7)	69 (47.6)	227 (45.9)	75 (37.3)	203 (48.6)	110 (44.9)	388 (44.9)
Satisfied	26 (36.1)	78 (39.6)	34 (34.0)	138 (37.4)	58 (45.0)	77 (34.8)	60 (41.4)	195 (39.4)	84 (41.8)	155 (37.1)	94 (38.4)	333 (38.5)
Total	72 (100.0)	197 (100.0)	100 (100.0)	369 (100.0)	129 (100.0)	221 (100.0)	145 (100.0)	495 (100.0)	201 (100.0)	418 (100.0)	245 (100.0)	864 (100.0)
Missing % missing	1 (1.4)	1 (0.5)	2 (2.0)	4 (1.0)	1 (1.0)	1 (0.5)	1 (1.0)	3 (0.6)	2 (1.0)	2 (0.5)	3 (1.4)	7 (0.8)

Difference Male vs Female in Late Adolescents P value = .016

**Table E.2A5 :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by Need for medical treatment to function their daily life

Need for medical treatment to function their daily life	Male				Female				M&F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
Very much	14 (19.2)	26 (13.2)	15 (14.7)	55 (14.8)	16 (12.5)	23 (10.4)	21 (14.6)	60 (12.2)	30 (14.9)	49 (11.8)	36 (14.6)	115 (13.3)
Moderate amount	22 (30.1)	85 (43.1)	34 (33.3)	141 (37.9)	41 (32.0)	80 (36.4)	47 (32.6)	168 (34.1)	63 (31.3)	165 (39.6)	81 (32.9)	309 (35.8)
A little	37 (50.7)	86 (43.7)	53 (52.0)	176 (47.3)	71 (55.4)	117 (53.2)	76 (52.8)	264 (53.7)	108 (53.7)	203 (48.7)	129 (52.4)	440 (50.9)
Total	73 (100.0)	197 (100.0)	102 (100.0)	372 (100.0)	128 (100.0)	220 (100.0)	144 (100.0)	492 (100.0)	201 (100.0)	417 (100.0)	246 (100.0)	864 (100.0)
Missing % missing	0 (0.0)	1 (0.5)	0 (0.0)	1 (0.3)	2 (1.5)	2 (1.0)	2 (1.5)	6 (1.2)	2 (1.0)	3 (0.7)	2 (1.0)	7 (0.8)

**Table E.2A6 :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by Satisfaction with their capacity for work/study

Satisfaction with their capacity for work/study	Male				Female				M&F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
Dissatisfied	12 (16.4)	31 (15.7)	17 (16.9)	60 (16.2)	18 (13.9)	32 (14.5)	12 (8.3)	62 (12.5)	30 (14.8)	63 (15.1)	29 (11.8)	122 (14.1)
Neither satisfied nor dissatisfied	27 (37.0)	108 (54.8)	43 (42.6)	178 (48.0)	55 (42.3)	115 (52.3)	66 (45.5)	236 (47.7)	82 (40.4)	223 (53.5)	109 (44.3)	414 (47.8)
Satisfied	34 (46.6)	58 (29.4)	41 (40.6)	133 (35.8)	57 (43.8)	73 (33.2)	67 (46.2)	197 (39.8)	91 (44.8)	131 (31.4)	108 (43.9)	330 (38.1)
Total	73 (100.0)	197 (100.0)	101 (100.0)	371 (100.0)	130 (100.0)	220 (100.0)	145 (100.0)	495 (100.0)	203 (100.0)	417 (100.0)	246 (100.0)	866 (100.0)
Missing % missing	0 (0.0)	1 (0.5)	1 (1.0)	2 (0.5)	0 (0.0)	2 (1.0)	1 (1.0)	3 (0.6)	0 (0.0)	3 (0.7)	2 (1.0)	5 (0.6)

Difference Early/Mid/Late

P value < .001

**Table E.2A7 :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by Ability to go around by themselves

Ability to go around by themselves	Male				Female				M&F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
A little	10 (14.0)	28 (14.3)	9 (8.9)	47 (12.8)	25 (19.4)	29 (13.3)	12 (8.3)	66 (13.4)	35 (17.5)	57 (13.7)	21 (8.6)	113 (13.1)
Moderate amount	25 (35.2)	67 (34.2)	39 (38.6)	131 (35.6)	47 (36.4)	91 (41.6)	39 (27.1)	177 (36.0)	72 (36.0)	158 (38.1)	78 (31.8)	308 (35.8)
Very much	36 (50.2)	101 (51.5)	53 (52.5)	190 (51.6)	57 (44.2)	99 (45.2)	93 (64.6)	249 (50.6)	93 (46.5)	200 (48.2)	146 (59.6)	439 (51.1)
Total	71 (100.0)	196 (100.0)	101 (100.0)	368 (100.0)	129 (100.0)	219 (100.0)	144 (100.0)	492 (100.0)	200 (100.0)	415 (100.0)	245 (100.0)	860 (100.0)
Missing % missing	2 (2.0)	2 (1.0)	1 (1.0)	5 (1.2)	1 (1.0)	3 (1.5)	2 (1.5)	6 (1.2)	3 (1.2)	5 (1.2)	3 (1.2)	11 (1.2)

Difference Early/Mid/Late

P value = .009

**B) Psychological domain**

**Table E.2B1 :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by Enjoy life

Enjoy Life	Male				Female				M&F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
A little	12 (16.4)	35 (17.7)	24 (23.7)	71 (19.1)	43 (33.0)	50 (22.6)	25 (17.1)	118 (23.7)	55 (27.1)	85 (20.3)	49 (19.8)	189 (21.8)
Moderate amount	28 (38.4)	81 (41.1)	43 (42.6)	152 (41.0)	44 (33.8)	98 (44.3)	65 (44.5)	207 (41.6)	72 (35.5)	179 (42.8)	108 (43.7)	359 (41.4)
Very much	33 (45.2)	81 (41.1)	34 (33.7)	148 (39.9)	43 (33.0)	73 (33.0)	56 (38.3)	172 (34.6)	76 (37.4)	154 (36.8)	90 (36.4)	320 (36.9)
Total	73 (100.0)	197 (100.0)	101 (100.0)	371 (100.0)	130 (100.0)	221 (100.0)	146 (100.0)	497 (100.0)	203 (100.0)	418 (100.0)	247 (100.0)	868 (100.0)
Missing % missing	0 (0.0)	1 (0.0)	1 (1.0)	2 (0.0)	0 (0.0)	1 (0.0)	0 (0.0)	1 (0.0)	0 (0.0)	2 (0.0)	1 (0.0)	3 (0.0)



**Table E.2B2 :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by Ability on the concentration on their work/study

Ability on the concentration on their work/study	Male				Female				M&F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
A little	17 (23.3)	24 (12.3)	13 (12.7)	54 (14.6)	23 (17.6)	33 (14.9)	20 (13.8)	76 (15.3)	40 (19.7)	57 (13.7)	33 (13.4)	130 (15.0)
Moderate amount	28 (38.4)	105 (53.8)	49 (48.0)	182 (49.2)	63 (48.5)	134 (60.4)	79 (54.5)	276 (55.5)	91 (44.8)	239 (57.3)	128 (51.8)	458 (52.8)
Very much	28 (38.3)	66 (33.9)	40 (39.2)	134 (36.2)	44 (33.8)	55 (24.8)	46 (31.8)	145 (29.2)	72 (35.5)	121 (29.0)	86 (34.8)	279 (32.2)
Total	73 (100.0)	195 (100.0)	102 (100.0)	370 (100.0)	130 (100.0)	222 (100.0)	145 (100.0)	497 (100.0)	203 (100.0)	417 (100.0)	247 (100.0)	867 (100.0)
Missing % missing	0 (0.0)	3 (1.5)	0 (0.0)	3 (0.8)	0 (0.0)	0 (0.0)	1 (1.0)	1 (0.2)	0 (0.0)	3 (0.7)	1 (0.4)	4 (0.5)

Difference Early/Mid/Late

P value < .001

**Table E.2B3 :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by Satisfaction with themselves

Satisfaction with themselves	Male				Female				M&F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
Dissatisfied	4 (5.4)	20 (10.2)	14 (13.7)	38 (10.2)	27 (20.8)	28 (12.7)	23 (15.8)	78 (15.7)	31 (15.3)	48 (11.5)	37 (14.9)	116 (13.3)
Neither satisfied nor dissatisfied	26 (35.6)	66 (33.5)	40 (39.2)	132 (35.5)	34 (26.2)	95 (43.0)	50 (34.2)	179 (36.0)	60 (29.6)	161 (38.5)	90 (36.3)	311 (35.8)
Satisfied	43 (58.9)	111 (56.3)	48 (47.1)	202 (54.3)	69 (53.1)	98 (44.3)	73 (50.0)	240 (48.3)	112 (55.2)	209 (50.0)	121 (48.8)	442 (50.9)
Total	73 (100.0)	197 (100.0)	102 (100.0)	372 (100.0)	130 (100.0)	221 (100.0)	146 (100.0)	497 (100.0)	203 (100.0)	418 (100.0)	248 (100.0)	869 (100.0)
Missing % missing	0 (0.0)	1 (0.5)	0 (0.0)	1 (0.3)	0 (0.0)	1 (0.5)	0 (0.0)	1 (0.2)	0 (0.0)	2 (0.5)	0 (0.0)	2 (0.2)

Difference Male vs Female in Early Adolescents

P value = .012

**Table E.2B4 :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by Acceptance of their bodily appearance

Acceptance of their bodily appearance	Male				Female				M&F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
A little	9 (12.3)	22 (11.1)	14 (14.3)	45 (12.2)	13 (10.3)	29 (13.2)	14 (9.6)	56 (11.4)	22 (11.1)	51 (12.2)	28 (11.5)	101 (11.7)
Moderate amount	31 (42.5)	80 (40.4)	37 (37.8)	148 (40.1)	43 (34.1)	79 (36.1)	68 (46.6)	190 (38.7)	74 (37.2)	159 (38.1)	105 (43.0)	338 (39.3)
Very much	33 (45.2)	96 (48.5)	47 (47.9)	176 (47.7)	70 (55.5)	111 (50.7)	64 (43.9)	245 (49.9)	103 (51.8)	207 (49.6)	111 (45.5)	421 (49.0)
Total	73 (100.0)	198 (100.0)	98 (100.0)	369 (100.0)	126 (100.0)	219 (100.0)	146 (100.0)	491 (100.0)	199 (100.0)	417 (100.0)	244 (100.0)	860 (100.0)
Missing % missing	0 (0.0)	0 (0.0)	4 (4.0)	4 (1.0)	4 (3.0)	3 (1.5)	0 (0.0)	7 (1.5)	4 (2.0)	3 (0.7)	4 (1.5)	11 (1.0)

**Table E.2B5 :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by Having frequently of negative feeling

Having frequently of negative feeling	Male				Female				M&F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
Very much	12 (16.7)	22 (11.1)	9 (8.9)	43 (11.6)	24 (18.5)	45 (20.3)	21 (14.4)	90 (18.1)	36 (17.8)	67 (16.0)	30 (12.1)	133 (15.3)
Moderate amount	26 (36.1)	66 (33.3)	27 (26.7)	119 (32.1)	37 (28.5)	66 (29.9)	41 (28.1)	144 (29.0)	63 (31.2)	132 (31.5)	68 (27.5)	263 (30.3)
A little	34 (47.2)	110 (55.5)	65 (64.4)	209 (56.3)	69 (53.0)	110 (49.7)	84 (57.6)	263 (52.9)	103 (51.0)	220 (52.5)	149 (60.3)	472 (54.4)
Total	72 (100.0)	198 (100.0)	101 (100.0)	371 (100.0)	130 (100.0)	221 (100.0)	146 (100.0)	497 (100.0)	202 (100.0)	419 (100.0)	247 (100.0)	868 (100.0)
Missing % missing	1 (1.0)	0 (0.0)	0 (0.0)	1 (0.3)	0 (0.0)	1 (0.5)	0 (0.0)	1 (0.2)	1 (0.5)	1 (0.2)	0 (0.0)	2 (0.2)

**Table E.2B6 :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by Feeling life is meaningful

Feeling life is meaningful	Male				Female				M&F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
A little	5 (6.8)	20 (10.3)	13 (12.8)	38 (10.4)	18 (14.0)	28 (12.7)	12 (8.3)	58 (11.7)	23 (11.4)	48 (11.6)	25 (10.2)	96 (11.1)
Moderate amount	17 (23.3)	68 (35.2)	36 (35.6)	121 (33.0)	43 (33.6)	78 (35.1)	35 (24.3)	156 (31.6)	60 (29.9)	146 (35.2)	71 (29.0)	277 (32.2)
Very much	51 (69.8)	105 (54.4)	52 (51.5)	208 (56.7)	67 (52.4)	116 (52.3)	97 (67.4)	280 (56.7)	118 (58.7)	221 (53.3)	149 (60.8)	488 (56.7)
Total	73 (100.0)	193 (100.0)	101 (100.0)	367 (100.0)	128 (100.0)	222 (100.0)	144 (100.0)	494 (100.0)	201 (100.0)	415 (100.0)	245 (100.0)	861 (100.0)
Missing % missing	0 (0.0)	5 (2.5)	1 (1.0)	6 (1.8)	2 (1.5)	0 (0.0)	2 (1.5)	4 (0.0)	2 (1.0)	5 (1.2)	3 (1.2)	10 (1.1)

**C) Social domains****Table E.2C1 :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by Satisfaction with their personal relationship

Satisfaction with their personal relationship	Male				Female				M&F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
Dissatisfied	9 (12.5)	21 (10.6)	7 (6.9)	37 (10.0)	12 (9.3)	19 (8.6)	4 (2.8)	35 (7.1)	21 (10.4)	40 (9.6)	11 (4.5)	72 (8.3)
Neither satisfied nor dissatisfied	22 (30.6)	76 (38.6)	43 (42.2)	141 (38.0)	46 (35.7)	75 (33.9)	47 (32.4)	168 (33.9)	68 (33.8)	151 (36.1)	90 (36.4)	309 (35.7)
Satisfied	41 (56.9)	100 (50.8)	52 (51.0)	193 (52.0)	71 (55.1)	127 (57.4)	94 (64.8)	292 (59.0)	112 (55.7)	227 (54.3)	146 (59.1)	485 (56.0)
Total	72 (100.0)	197 (100.0)	102 (100.0)	371 (100.0)	129 (100.0)	221 (100.0)	145 (100.0)	495 (100.0)	201 (100.0)	418 (100.0)	247 (100.0)	866 (100.0)
Missing % missing	1 (1)	1 (0.5)	0 (0)	2 (0.5)	1 (1)	1 (0.5)	1 (0.7)	3 (0.8)	2 (1)	2 (0.5)	1 (0.8)	5 (0.6)

**Table E.2C2 :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by Satisfaction with the support from their friend

Satisfaction with the support from their friend	Male				Female				M&F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
Dissatisfied	5 (7.0)	17 (8.7)	11 (11.0)	33 (9.0)	14 (11.0)	22 (9.9)	13 (8.9)	49 (9.9)	19 (9.5)	39 (9.4)	24 (9.8)	82 (9.5)
Neither satisfied nor dissatisfied	31 (43.1)	78 (40.0)	47 (47.0)	156 (42.5)	50 (39.1)	81 (36.7)	49 (33.6)	180 (36.4)	81 (40.5)	159 (38.2)	96 (39.0)	336 (39.0)
Satisfied	36 (50.0)	100 (51.3)	42 (42.0)	178 (48.5)	64 (50.0)	118 (53.4)	84 (57.5)	266 (53.7)	100 (50.0)	218 (52.4)	126 (51.2)	444 (51.5)
Total	72 (100.0)	195 (100.0)	100 (100.0)	367 (100.0)	128 (100.0)	221 (100.0)	146 (100.0)	495 (100.0)	200 (100.0)	416 (100.0)	246 (100.0)	862 (100.0)
Missing	1	3	2	6	2	1	0	3	3	4	2	9
% missing	(1.4)	(1.5)	(2)	(1.8)	(1.5)	(0.5)	(0)	(1.5)	(1.5)	(1)	(1)	(1)

**Table E.2C3 :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by Satisfaction with their sex life

Satisfaction with their sex life	Male				Female				M&F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
Dissatisfied	27 (38.6)	52 (26.6)	24 (23.5)	103 (28.1)	50 (38.8)	102 (47.2)	42 (29.2)	194 (39.7)	77 (38.7)	154 (37.5)	66 (26.9)	297 (34.7)
Neither satisfied nor dissatisfied	22 (31.4)	84 (43.1)	52 (51.0)	158 (43.1)	41 (31.8)	74 (34.3)	61 (42.4)	176 (36.0)	63 (31.7)	158 (38.4)	113 (45.9)	334 (39.0)
Satisfied	21 (30.0)	59 (30.3)	26 (25.5)	106 (28.8)	38 (29.5)	40 (18.5)	41 (28.5)	119 (24.3)	59 (29.6)	99 (24.1)	67 (27.2)	225 (26.3)
Total	70 (100.0)	195 (100.0)	102 (100.0)	367 (100.0)	129 (100.0)	216 (100.0)	144 (100.0)	489 (100.0)	199 (100.0)	411 (100.0)	246 (100.0)	856 (100.0)
Missing	3	3	0	6	1	6	2	9	4	9	2	15
% missing	(4)	(1.5)	(0)	(1.8)	(1)	(2.5)	(1.4)	(1.8)	(2)	(2.2)	(1)	(1.8)

Difference Male vs Female

P value = .002

Difference Early/Mid/Late

P value = .008

Difference Male vs Female in Mid Adolescents

P value < .001

**D) Environmental domain**

**Table E.2D1 :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by Feeling safely

Feeling safely	Male				Female				M&F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
A little	4 (5.5)	29 (14.8)	25 (24.5)	58 (15.6)	22 (17.2)	34 (15.4)	22 (15.1)	78 (15.7)	26 (12.9)	63 (15.1)	47 (19.0)	136 (15.7)
Moderate amount	34 (46.6)	76 (38.8)	39 (38.2)	149 (40.2)	46 (35.9)	110 (49.5)	56 (38.4)	212 (42.7)	80 (39.8)	186 (44.5)	95 (38.3)	361 (41.6)
Very much	35 (47.9)	91 (46.4)	38 (37.2)	164 (44.2)	60 (46.9)	78 (35.1)	68 (46.5)	206 (41.5)	95 (47.3)	169 (40.4)	106 (42.7)	370 (42.7)
Total	73 (100.0)	196 (100.0)	102 (100.0)	371 (100.0)	128 (100.0)	222 (100.0)	146 (100.0)	496 (100.0)	201 (100.0)	418 (100.0)	248 (100.0)	867 (100.0)
Missing	0	2	0	2	2	0	0	2	2	2	0	4
% missing	(0)	(1)	(0)	(0.5)	(1.5)	(0)	(0)	(0.4)	(1)	(0.2)	(0)	(0.5)

**Table E.2D2 :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by Satisfaction with condition of their living place

Satisfaction with condition\ of their living place	Male				Female				M&F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
Dissatisfied	9 (12.4)	27 (13.7)	26 (25.5)	62 (16.7)	25 (19.2)	41 (18.6)	26 (17.9)	92 (18.5)	34 (16.7)	68 (16.3)	52 (21.1)	154 (17.8)
Neither satisfied nor dissatisfied	24 (32.9)	88 (44.9)	39 (38.2)	151 (40.7)	51 (39.2)	92 (41.6)	67 (46.2)	210 (42.3)	75 (36.9)	180 (43.2)	106 (42.9)	361 (41.6)
Satisfied	40 (54.7)	81 (41.3)	37 (36.3)	158 (42.6)	54 (41.6)	88 (39.8)	52 (35.8)	194 (39.1)	94 (46.4)	169 (40.5)	89 (36.0)	352 (40.6)
Total	73 (100.0)	196 (100.0)	102 (100.0)	371 (100.0)	130 (100.0)	221 (100.0)	145 (100.0)	496 (100.0)	203 (100.0)	417 (100.0)	247 (100.0)	867 (100.0)
Missing	0	2	0	2	0	1	1	2	0	3	1	4
% missing	(0)	(1)	(0)	(0.5)	(0)	(0.5)	(1)	(0.4)	(0)	(0.7)	(0.4)	(0.5)

**Table E.2D3 :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by Money and their need

Money and their need	Male				Female				M&F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
A little	13 (17.8)	34 (17.4)	35 (34.3)	82 (22.1)	33 (25.4)	57 (25.9)	36 (24.7)	126 (25.4)	46 (22.7)	91 (21.9)	71 (28.6)	208 (24.0)
Moderate amount	30 (41.1)	106 (54.1)	50 (49.0)	186 (50.1)	65 (50.0)	108 (49.1)	82 (56.2)	255 (51.4)	95 (46.8)	214 (51.4)	132 (53.2)	441 (50.9)
Very much	30 (41.1)	56 (28.5)	17 (16.7)	103 (27.8)	32 (24.7)	55 (25.0)	28 (19.2)	115 (23.2)	62 (30.5)	111 (26.7)	45 (18.1)	218 (25.1)
Total	73 (100.0)	196 (100.0)	102 (100.0)	371 (100.0)	130 (100.0)	220 (100.0)	146 (100.0)	496 (100.0)	203 (100.0)	416 (100.0)	248 (100.0)	867 (100.0)
Missing % missing	0 (0)	2 (1)	0 (0)	2 (0.5)	0 (0)	2 (1)	0 (0)	2 (0.4)	0 (0)	4 (1)	0 (0)	4 (0.5)

**Table E.2D4 :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by Satisfaction with health services accessibility

Satisfaction with health services accessibility	Male				Female				M&F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
Dissatisfied	13 (17.8)	40 (20.4)	24 (23.5)	77 (20.8)	31 (24.6)	48 (21.7)	28 (19.2)	107 (21.7)	44 (22.1)	88 (21.1)	52 (21.0)	184 (21.3)
Neither satisfied nor dissatisfied	32 (43.8)	97 (49.5)	55 (53.9)	184 (49.6)	57 (45.2)	130 (58.8)	84 (57.5)	271 (55.0)	89 (44.7)	227 (54.4)	139 (56.0)	455 (52.7)
Satisfied	28 (38.3)	59 (30.1)	23 (22.5)	110 (29.6)	38 (30.2)	43 (19.5)	34 (23.3)	115 (23.3)	66 (33.2)	102 (24.5)	57 (23.0)	225 (26.0)
Total	73 (100.0)	196 (100.0)	102 (100.0)	371 (100.0)	126 (100.0)	221 (100.0)	146 (100.0)	493 (100.0)	199 (100.0)	417 (100.0)	248 (100.0)	864 (100.0)
Missing % missing	0 (0)	2 (1)	0 (0)	2 (0.5)	4 (3)	1 (0.5)	0 (0)	5 (1)	4 (2)	3 (0.7)	0 (0)	7 (0.8)

**Table E.2D5 :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by Availability on the day to day information needed

Availability on the day to day information needed	Male				Female				M&F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
A little	17 (23.2)	34 (17.2)	27 (26.4)	78 (20.9)	27 (20.8)	42 (18.9)	27 (18.6)	96 (19.3)	44 (21.7)	76 (18.1)	54 (21.9)	174 (20.0)
Moderate amount	28 (38.4)	101 (51.0)	40 (39.2)	169 (45.3)	61 (46.9)	109 (49.1)	78 (53.8)	248 (49.9)	89 (43.8)	210 (50.0)	118 (47.8)	417 (47.9)
Very much	28 (38.4)	63 (31.8)	35 (34.3)	126 (33.8)	42 (32.3)	71 (32.0)	40 (27.6)	153 (30.8)	70 (34.5)	134 (31.9)	75 (30.4)	279 (32.1)
Total	73 (100.0)	198 (100.0)	102 (100.0)	373 (100.0)	130 (100.0)	222 (100.0)	145 (100.0)	497 (100.0)	203 (100.0)	420 (100.0)	247 (100.0)	870 (100.0)
Missing % missing	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	1 (1)	1 (0.2)	0 (0)	0 (0)	1 (0.4)	1 (0.1)

**Table E.2D6 :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by Having opportunity for leisure activities

Having opportunity for leisure activities	Male				Female				M&F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
A little	12 (16.4)	35 (17.7)	24 (23.7)	71 (19.1)	43 (33.0)	50 (22.6)	25 (17.1)	118 (23.7)	55 (27.1)	85 (20.3)	49 (19.8)	189 (21.8)
Moderate amount	28 (38.4)	81 (41.1)	43 (42.6)	152 (41.0)	44 (33.8)	98 (44.3)	65 (44.5)	207 (41.6)	72 (35.5)	179 (42.8)	108 (43.7)	359 (41.4)
Very much	33 (45.2)	81 (41.1)	34 (33.7)	148 (39.9)	43 (33.0)	73 (33.0)	56 (38.3)	172 (34.6)	76 (37.4)	154 (36.8)	90 (36.4)	320 (36.9)
Total	73 (100.0)	197 (100.0)	101 (100.0)	371 (100.0)	130 (100.0)	221 (100.0)	146 (100.0)	497 (100.0)	203 (100.0)	418 (100.0)	247 (100.0)	868 (100.0)
Missing % missing	0 (0)	1 (0.5)	1 (1)	2 (0.5)	0 (0)	1 (0.5)	0 (0)	1 (0.2)	0 (0)	2 (0.5)	1 (0.4)	3 (0.3)

**Table E.2D7 :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by Feeling about the healthy of their physical environment

Feeling about the healthy of their physical environment	Male				Female				M&F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
A little	14 (19.2)	36 (18.4)	26 (25.7)	76 (20.5)	28 (21.7)	48 (22.1)	37 (25.7)	113 (23.1)	42 (20.8)	84 (20.3)	63 (25.7)	189 (22.0)
Moderate amount	33 (45.2)	99 (50.5)	49 (48.5)	181 (48.9)	65 (50.4)	112 (51.6)	70 (48.6)	247 (50.4)	98 (48.5)	211 (51.1)	119 (48.6)	428 (49.8)
Very much	26 (35.6)	61 (31.1)	26 (25.7)	113 (30.5)	36 (28.0)	57 (26.2)	37 (25.7)	130 (26.5)	62 (30.7)	118 (28.6)	63 (25.7)	243 (28.3)
Total	73 (100.0)	196 (100.0)	101 (100.0)	370 (100.0)	129 (100.0)	217 (100.0)	144 (100.0)	490 (100.0)	202 (100.0)	413 (100.0)	245 (100.0)	860 (100.0)
Missing % missing	0 (0)	2 (1)	1 (1)	3 (0)	1 (1)	5 (2.5)	2 (1.5)	8 (1.5)	1 (0)	7 (1.8)	3 (1.4)	11 (1.3)

**Table E.2D8 :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by Satisfaction with their transportation

Satisfaction with their transportation	Male				Female				M&F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
Dissatisfied	15 (20.9)	39 (19.8)	32 (31.3)	86 (23.2)	24 (18.8)	47 (21.2)	27 (18.5)	98 (19.8)	39 (19.5)	86 (20.5)	59 (23.8)	184 (21.2)
Neither satisfied nor dissatisfied	26 (36.0)	89 (45.2)	44 (43.1)	159 (42.9)	57 (44.5)	116 (52.3)	76 (52.1)	249 (50.2)	83 (41.5)	205 (48.9)	120 (48.4)	408 (47.1)
Satisfied	31 (43.0)	69 (35.0)	26 (25.5)	126 (34.0)	47 (36.7)	59 (26.6)	43 (29.5)	149 (30.0)	78 (39.0)	128 (30.5)	69 (27.8)	275 (31.7)
Total	72 (100.0)	197 (100.0)	102 (100.0)	371 (100.0)	128 (100.0)	222 (100.0)	146 (100.0)	496 (100.0)	200 (100.0)	419 (100.0)	248 (100.0)	867 (100.0)
Missing % missing	1 (1.4)	1 (0.5)	0 (0)	2 (0.5)	2 (1.5)	0 (0)	0 (0)	2 (0.4)	3 (1.5)	1 (0.2)	0 (0)	4 (0.5)





**C) The Psychological Domain of QoL**

**Table E.3C :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by Level of QoL of Psychological Domain

Level of QoL of psychological Domain	Male				Late				Female			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
Poor	2 (2.7)	8 (4.0)	6 (5.9)	16 (4.3)	8 (6.2)	12 (5.4)	4 (2.7)	24 (4.8)	10 (4.9)	20 (4.8)	10 (4.0)	40 (4.6)
Moderate	44 (60.3)	122 (61.6)	55 (54.5)	221 (59.4)	78 (60.0)	136 (61.3)	90 (61.6)	304 (61.0)	122 (60.1)	258 (61.4)	145 (58.7)	525 (60.3)
High	27 (37.0)	68 (34.3)	40 (39.6)	135 (36.3)	44 (33.8)	74 (33.3)	52 (35.6)	170 (34.1)	71 (35.0)	142 (33.8)	92 (37.2)	305 (35.1)
Total	73 (100.0)	198 (100.0)	101 (100.0)	372 (100.0)	130 (100.0)	222 (100.0)	146 (100.0)	498 (100.0)	203 (100.0)	420 (100.0)	247 (100.0)	870 (100.0)
Missing	0	0	1	1	0	0	0	0	0	0	0	1
%missing	(0)	(0)	(1)	(0.3)	(0)	(0)	(0)	(0)	(0)	(0)	(0)	(0.1)

**D) The Social Domain of QoL**

**Table E.3D :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by Level of QoL of Social Domain

Level of QoL of Social Domain	Male				Female				M & F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
Poor	18 (24.7)	25 (12.6)	10 (9.8)	53 (14.2)	19 (14.6)	30 (13.5)	10 (6.8)	59 (11.8)	37 (18.2)	55 (13.1)	20 (8.1)	112 (12.9)
Moderate	31 (42.5)	122 (61.6)	71 (69.6)	224 (60.1)	81 (62.3)	153 (68.9)	98 (67.1)	332 (66.7)	112 (55.2)	275 (65.5)	169 (68.1)	556 (63.8)
High	24 (32.9)	51 (25.8)	21 (20.6)	96 (25.7)	30 (23.1)	39 (17.6)	38 (26.0)	107 (21.5)	54 (26.6)	90 (21.4)	59 (23.8)	203 (23.3)
Total	73 (100.0)	198 (100.0)	102 (100.0)	373 (100.0)	130 (100.0)	222 (100.0)	146 (100.0)	498 (100.0)	203 (100.0)	420 (100.0)	248 (100.0)	871 (100.0)

Difference Male vs Female

P value = .008

Difference Early/Mid/Late in Male

P value = .004

*E) The Environmental Domain of QoL*

**Table E.3E :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by the Environmental Domain of QoL

Environmental Domain of QoL	Male				Female				M & F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
Poor	0 (0)	8 (4.0)	10 (9.8)	18 (4.8)	11 (8.5)	16 (7.2)	7 (4.8)	34 (6.8)	11 (5.4)	24 (5.7)	17 (6.9)	52 (6.0)
Moderate	55 (75.3)	153 (77.3)	78 (76.5)	286 (76.7)	92 (70.8)	179 (80.6)	119 (81.5)	390 (78.3)	147 (72.4)	332 (79.0)	197 (79.4)	676 (77.6)
High	18 (24.7)	37 (18.7)	14 (13.7)	69 (18.5)	27 (20.8)	27 (12.2)	20 (13.7)	74 (14.9)	45 (22.2)	64 (15.2)	34 (13.7)	143 (16.4)
Total	73 (100.0)	198 (100.0)	102 (100.0)	373 (100.0)	130 (100.0)	222 (100.0)	146 (100.0)	498 (100.0)	203 (100.0)	420 (100.0)	248 (100.0)	871 (100.0)

*F) Summation of all domains of QoL*

**Table E.3F :** Descriptive Analysis of Adolescents: Distribution by Stage of Adolescence (age) by Gender and by The summation of all domains of QoL

Level of Quality of Life	Male				Female				M&F			
	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)	Early n (%)	Mid n (%)	Late n (%)	Total n (%)
Poor	2 (2.7)	3 (1.5)	3 (2.9)	8 (2.1)	2 (1.5)	2 (0.9)	1 (0.7)	5 (1.0)	4 (2.0)	5 (1.2)	4 (1.6)	13 (1.5)
Moderate	49 (67.1)	138 (69.7)	74 (72.5)	261 (70.0)	92 (70.8)	165 (74.3)	107 (73.3)	364 (73.1)	141 (69.5)	303 (72.1)	181 (73.0)	625 (71.8)
High	22 (30.1)	57 (28.8)	25 (24.5)	104 (27.9)	36 (27.7)	55 (24.8)	38 (26.0)	129 (25.9)	58 (28.6)	112 (26.7)	63 (25.4)	233 (26.8)
Total	73 (100.0)	198 (100.0)	102 (100.0)	373 (100.0)	130 (100.0)	222 (100.0)	146 (100.0)	498 (100.0)	203 (100.0)	420 (100.0)	248 (100.0)	871 (100.0)

## Curriculum Vitae

**Name** : Ratana Somrongthong

### Education

- 1975-1978 : Dip. of Nursing and Midwifery, Thai Red Cross  
Society Nursing College, Chulalongkorn University, Thailand
- 1981- 1983 : B. Sc. (Social Psychology), Ramkhamhaeng University, Thailand
- 1986-1990 : B. of Home Economic(Community Nutrition)  
Sukhothaimathirath University, Thailand
- 1993 –1996 : MA (Medical and Health Social Sciences) Mahidol  
University, Thailand
- 2000- present : PhD Candidate in Public Health (Health Systems Development),  
College of Public Health, Chulalongkorn University, Thailand

### Publication:

#### *Refereed*

- Pramuanratana, A. Somrongthong,R. Jindasak, K.(1995) Assessment of the potential for spread and control of HIV among cross-border populations along the Thai-Cambodian border. **Institute for populations and Social Research, Mahidol University**
- Pannee, P., Somrongthong, R. .(1999) Sexual Behaviors and the Opinions about Sexuality of Adolescents in a Slum Community. **The Thai AIDS Journal**, 11: 207-216 (Thai Version)
- Ratana, S., Chitr, S. (2000) Existing Health Needs and Related Health Services for Adolescents in Slum Community in Thailand. **International Journal of Adolescent Medicine and Health**. Freund Publishing House, LTD., London, Tel Aviv. 12, NOS, 2-3: 191-201
- Sitthi-amorn, C., Somrongthong, R., (2000) Multi-disciplinary research in public health. **Journal of Research Methodology**. 13: 1-19. (Thai version)



- Sitthi-amorn, C., Somrongthong, R.(2000) Strengthening Health Research Capacity in Developing Countries: A Critical Element for Achieving Health Equity. **BMJ**, No.7246: 813-817
- Sitthi-amorn, C., Somrongthong, R., Wattana S. Janjaroen. (2001) Economic and Health Care Restructuring The Need for Better Governance. **International Journal of Epidemiology**; 30: 717-719.
- Sitthi-amorn, C., Somrongthong, R., Wattana S. Janjaroen. (2001) The Health Effect of Globalization in Thailand. **Bulletin of the World Health Organization**. 79 (9); 889-890.
- Chitr Sitthi-amorn, Sathirakorn Pongpanich, Ratana Somrongthong, Tanawat Likitkirirat and Polachai Likitkirirat. (2002).The Asian Voice in Building Equity in Health of Development- from the Asian Forum for Health Research. **Health Policy and Planning. A journal on health in development** 17, 2:213-217
- Pannee Panuwatsuk, Ratana Somrongthong, Malai Vongsrisunthorn, Koobkul Poolpanyavongse. ( 2002). Health illness and factors influencing the opinion about utilization of Banna primary health care center, Chumporn Province for the next illness. **Region 11 Medical Journal**. 16, 2002:1-6. (in Thai)
- Ratana Somrongthong, Pannee Panuwatsuk, David Amarathithada, Orapin Chaipayom, Chitr Sitthi-amorn. ( 2003). Sexual Behaviours and Opinions on Sexuality of Adolescents in a Slum Community in Bangkok. **Southeast Asian Journal of Tropical Medicine and Public Health. Thailand**. Vol.34, No.2 June 2003. P 443-446.
- Somrongthong R, Suttajit S, Tantipidoke R, Sitthi-amorn, Ross Degnan D. (2004). Using qualitative methods to develop a community-based intervention to improve antibiotic use in treating adult ARI in a slum community in Thailand. Abstract book, the International Conference on Improving Use of Medicines, Chiang Mai, Thailand, 29 March – 2 April 2004.

**Non- Refereed**

- USAID, Office of Khmer Affairs, Thailand. (1993) Expanded Programme on Immunization in Cambodia: Immunization Coverage Survey in Thmar Pouk District & Ampil District, Banteay Meanchai Province, **USAID, Bangkok**, Thailand
  
- Somruedee Sarapirum, Ratana Somrongthong, Chatchai Wangcheu, Aupum Pornmesri, Tsuyoshi Kawakami. ( 1995). Tuberculosis Control in Industrializing & Urbanizing Regions (Situation Analysis): A Case Study at Samutprakan Provincial Hospital: **Proceeding of the International workshop on Improvement of Tuberculosis Control in Developing Countries**, organizing by Institute for Science of Labour and The Research Institute of Tuberculosis, Japan p 185-195.