# DETECTION OF HIGH-RISK PREGNANCIES USING HOME-BASED PRENATAL RECORD THROUGH THE TRADITIONAL BIRTH ATTENDANTS: AN INTERVENTION TO INCREASE ANTENATAL COVERAGE IN NEPAL

Shakuntala Shakya

A Thesis Submitted in Partial Fulfillment of the Requirements for the Degree of Master of Public Health,
Health System Development Programme
College of Public Health
Chulalongkorn University
Academic Year 1998
ISBN: 974-639-230-1

Copyright of College of Public Health, Chulalongkorn University

Thesis Title Detection of High-Risk Pregnancies Using Home-Based Prenatal Record Through the Traditional Birth Attendants: An Intervention to Increase Antenatal Coverage in Nepal. : Shakuntala Shakya By : Master of Public Health (Health Systems Development) Programme College of Public Health. : Nuntavarn Vichit-Vadakan, M.S., M.P.H., Dr. P.H. Thesis Advisor Accepted by the College of Public Health, Chulalongkorn University in Partial Fulfillment of the Requirements for the Master's Degree. Bhit SThi-amwn Dean of the College of Public Health (Professor Chitr Sitthi-amorn, M.D., Ph.D.) THESIS COMMITTEE Wattana S. Tanjara Chairman (Assoc, Prof. Wattana S. Janjaroen, Ph.D.) Vicht-Ordle\_\_\_\_Thesis Advisor (Dr. Nuntavarn Vichit-Vadakan, M.S., M.P.H., Dr.P.H.) ph. King Member (Dr. Stephen King, M. Med. Sci., Dr. P.H.) Redsara Aughet Member

(Assistant Professor Dr. Alisara Chuchat, Ph.D.)

#### **ABSTRACT**

It has been recognized that maternal mortality in Nepal is 539/100,000 live birth which is the highest in the developing countries. Most of the deaths are known to be from the complication of pregnancy and child birth and occur in "at risk" condition which can be prevented by means of antenatal care. Central Bureau of Statistics, National Planning Commission, Kathmandu, Nepal 1994 had shown that out of 370, 890 of total pregnancy 40 percent are in at risk condition. Demographic Health Survey 1995 /1996 shows that National coverage of antenatal service is only 19 percent (Antenatal visits as a percentage of expected pregnancy). Thus more than 80 percent of the pregnant women do not receive any care during pregnancy. Therefore incidence of obstetrical complication and deaths are very high which is really a major Public Health problem in Nepal.

His Majesty's Government of Nepal has set a target to reduce maternal mortality from the present rate to 400 /1000 live birth by the year 2000 AD with the provision to provide antenatal care services by strengthening community based maternity care services by mobilizing the community based health workers which is primarily through Primary Health Care Approach. Antenatal care is a major component of maternity service. Therefore, specific training is required to identify high risk pregnancy in the community level and refer those mothers to the health cente who are in "at risk" for timely management so as to reduce the obstetrical complications and deaths.

This study proposed a seven days training program for Traditional Birth Attendant on Home-based Prenatal Records for the detection of high-risk pregnancies and to increase the antenatal coverage. Training will be carried out in the Kalyanpur Primary Health Center of Siraha District, Nepal in June, 1999 with the support of National Health Training Center which will be funded by Redd Barna /UNICEF. The target population of this study will be all the pregnant women of Siraha District. Impact evaluation will be made after 12 months of program implementation. Method of data collection will be official statistics, Observation, and survey questionnaire. The evaluation team include NHTC and NHECC and representative from funding agency, will take responsibility for data processing and disseminated for the final reports by April, 2000.

### **ACKNOWLEDGEMENTS**

My profound appreciation goes to the Dean, Professor Chitr Sithi-amron for his valuable guidance, encouragement throughout my study. I really appreciate his vast knowledge on scientific research, which has contributed a lot for my study. I extent my sincere regards to Dr. Wattana S.J. Associate Professor and Deputy Dean for her support during my study.

I am very grateful to Ajarn Dr. Stephen King who provided me ideas of logical thinking in viewing the situation as a learning process through out my study period. I will never forget for his great support and assurance in the period of financial problem.

I would like to express my deepest gratitude to my advisor Ajarn Dr. Nuntavarn Vichit-Vadakan who provided me frequent guidance, valuable and constructive suggestion, encouragement, and comments that made this project possible.

I am very grateful to Ajarn Wacharin T. who helped me a lot to learn especially about the use of statistical concept and data exercise. I am very thankful to Ajarn Sathirakorn P. for his kind support, creative advice and sympathetic approach.

I would like to extend my hearty thanks to my co-advisor Ajarn Benjalak Sangsingkeo of Department of Health, Ministry of Public Health, Thailand, who gave me a great assistance, contribution, and ideas to prepare my proposal to this stage.

I would like to express my deepest gratitude to Ajarn Ratana S. and Ajarn Chanawong B. for their courageous, continuous support, and assistance through out the whole year of my study period.

I would like to extend my hearty thanks to Ajarn Tanawat L., Mr Thanapong., Ms Sunanta Wongchalee, and Ms Kingkeau Keawbunroung for their unforgettable help in preparing this study. I would like to give special thanks to Ms Duang and all the staffs of this college for their kind co-operation and helping me directly and indirectly during my study period.

I have no words to express my cordial thanks to my friend Mrs Pawadee Kucita who gave me continuous support, love, and assistance which made my stay in Thailand comfortable and memorable.

My heartfelt gratitude goes to the Director of Nursing Education Department, IOM, Professor Dr. Indira Singh and Medical Educationist Dr. P.T. Jayawickramrajah, WHO, for providing me this opportunity to study MPH in Thailand.

Finally, I have no word to thank my Husband Mr. Krishna Man Shakya for his courageous and dedicated support without which this study would have not been possible.

## LIST OF CONTENTS

	PAGE
ABSTRACT	<b>iii</b>
ACKNOWLEDGEMENTS	iv
LIST OF CONTENTS.	vi
LIST OF TABLES	x
LIST OF FIGURES	xi
ACRONYMS AND ABBREVIATION	xii
CHAPTER I:	
Introduction	1
CHAPTER II: ESSAY	
2.1.Introduction	6
2.2 High Risk Pregnancy	15
2.2.1 Incidences of maternal deaths	15
2.2.2. Risk factors	19
2.2.3. Complications	21
2.2.4. Consequences	24
2.3. Analytical framework	27
2.3.1. Socio-economical impact	28
2.3.2. Cultural influences	28
2.3.3. Bio-physical factors	30

2.3.4. Health service characteristics
2.4. Reduction of risk
2.5. Improving service coverage
2.5.1 Anternatal Coverage
2.5.2. Needs of antenatal services
2.5.3 Effectiveness of services.
2.6. Planned Strategies to increase antenatal coverage
2.6.1. Increasing access to antenatal services
2.6.2. Acceptable health services through female to
female approach
2.6.3. Improvement of existing knowledge, attitudes, and
practices of Traditional Birth Attendants (TBAs)43
2.6.4. Use of low technology
2.6.5.Emphasis on IEC activities
2.6.6. Coordination between health workers
2.7. Constraints of programs
2.8. Financial support 48
2.9. Program sustainability
2.10. Recommendation/ possible alternatives
2.11. Availability of service within the health care system
2.12. Conclusion
References 56

## **CHAPTER III: PROPOSAL**

3.1. Introduction
3.2. Rational of training approach
3.3. Problem statement
3.4. Purpose of the study
3.5. Goal
3.5.1. Objectives of the study69
3.6. Intervention area: Siraha district, Nepal
3.7. Proposed intervention programs
3.7.1. Development of Home-Based Prenatal Record (HBPR)77
3.7.2. Training program for TBAs on HBPR85
3.7.3. Implementation of HBPR94
3.7.4. Provision of IEC in prenatal teaching98
3.8. Supervision of TBAs providing ANC services using HBPR99
3.9. Monitoring ANC services and use of HBPR101
3.10. Evaluation of TBA's activities in ANC services
3.10.1. Training progress and learning outcome evaluation103
3.10.2. Impact evaluation training on TBAs
3.10.3. Impact evaluation of antenatal services on women106
3. 11. Budget for the study
3.12 Potential problems
3.13. Ethical issue in the study
3.14. Limitations of the study

3.15. Program sustainability	121
References	124
CHAPTER IV: DATA EXERCISE	
4.1. Introduction	125
4.2. Objectives of Data Exercise.	126
4.3. Data collection technique	126
4.4. Data collection procedure	127
4.4.1 Field preparation	127
4.4.2 Data exercise population	128
4.5. Limitations of data exercise	129
4.6. Findings	130
4.6.1. Knowledge of antenatal service	130
4.6.2. Use of antenatal service	131
4.6.3. Decision to seek antenatal service	131
4.6.4. Source of information about service	131
4.6.5. Knowledge of risk symptoms during pregnancy	132
4.6.6. Preference of person and place for the service	132
4.6.7. Frequency of visit	133
4.7. Discussion	133
4.8. Conclusion	135
4.9. Lesson learned from FGD.	

CHAPTER V:
Presentation
CHAPTER VI:
Annotated Bibliography
APPENDICES:
Appendix I Curriculum of the training
Appendix II Suggested HBPR, side I and II
Appendix III. Potential member of Project Implementation Team
Appendix IV Pre-test and post-test format and questionnaire
Appendix V Antenatal care checklist
Appendix VI FGD guidelines for women's group
Appendix VII FGD guidelines for TBAs
Appendix VIII Survey questionnaire for impact evaluation
Student's Curriculum Vitae
LIST OF TABLES
TABLE PAGE
2.1: Estimated global incidence and mortality for main obstetric
2.2. Causes of maternal deaths by age group within
10 years time (1993)

2.3. Assessment of risk factors for maternal mortality and morbidity, low birth
weight, and perinatal mortality
2.4. Complications affecting on health of the mother and fetus/baby23
3.1. Reported Antenatal coverage % by Regions of Nepal
3.2. The health facilities available in Siraha district are as follows:
3.3. List of Proposed intervention programs
3.4. Proposed activities with Timetable I, II
3.5. Component of impact evaluation
3.6. Budget for the pilot study
LIST OF FIGURES
FIGURE
2.1. Major cause and consequences of pregnancy complications
2.2. Analytical framework of casual relationship of antenatal service
utilization and reduction of risk
3.1. A flow chart recommended by WHO for showing decision making
process for choosing factors for risk assessment to be included in HBPR80
3.2. Suggested contents to be included in HBPR, side I
3.3. Suggested contents to be included in HBPR, side II
3.4. Suggested Register for consolidating information from HBPR84

## ACRONYMS AND ABBREVIATION

ANC = Antenatal care

ANM = Auxiliary Nurse Midwife

DHS = Demographic and Health Survey

FCHV = Female Community Health Volunteer

FGD = Focus Group Discussion

HBPR = Home-Based Prenatal Record

HMG = His Majesty's Government

HMIS = Health Management Information System

IEC = Information Education Communication

JSI = John Snow Inc.

MCH = Maternal Child Health Worker

MIS = Management Information System

MOH = Ministry Of Health

NFHS = Nepal Family Health Survey

NGO = Non Governmental Organization

NPC = National Planning Commission

PHC = Primary Health Center

SHS = Strengthening Health Services

SMPN = Safe Motherhood Program in Nepal

TB = Tuberculosis

TBA = Traditional Birth Attendant