

## References

- Ananvoranich, S., Likhitwitayawuid, K. and Ruangrungsi, N. 1989. Determination of a New Sesquiterpene Skeleton through Selective INEPT Spectroscopy. Journal of Organic Chemistry. Vol. 54(9), 2253.
- Ansari, S., Jain, P., Tyagi, R. Joshi, B.C. and Barar, F.S.K. 1985. Phytochemical and pharmacological studies of the aerial parts of *Eupatorium adenophorum* L. Chemical Abstracts. Vol. 102.
- Association for International Cooperation of Agriculture & Forestry, Japan (AICAF). 1997. Weed In The Tropics. Sanbi Printing Co.,Ltd. 304.
- Baruah, N.C., Sarma, J.C., Sarma, S. and Sharma, R.P. 1994. Seed Germination and Growth Inhibitory Cadinenes from *Eupatorium adenophorum* Spreng. Journal of Chemistry Ecology. Vol. 20(8), 1885.
- Benson, L.D. 1959. Plant Classification. ASA GRAY. D.C. Health and Company. 295.
- Bergeron, C., Marston, A., Gauthier, R. and Hostettmann, K. 1996. Screening of Plants Used by North American Indians for Antifungal, Bactericidal, Larvicidal, and Molluscicidal Activities. International Journal of Pharmacognosy. Vol. 34(4), 233.
- Bohlmann, F. and Gupta, R.K. 1981. Six Cadinene Derivatives from *Ageratina adenophorua*. Phytochemistry. Vol. 20(6), 1432.
- Bordoloi, M.J., Shukla, V.S. and Sharma, R.P. 1985. Absolute Stereochemistry of the Insect Antifeedant Cadinene from *Eupatorium adenophorum*. Tetrahedron Letter, Vol. 26(4), 509.
- Colegate, S.M. and Molyneux, R.J. 1993. Bioactive Natural Products. CRC Press. 441.
- Cooper, J.W. 1980. Spectroscopic Techniques for Organic Chemistry, 1<sup>st</sup> ed., John-Wiley & Sons, USA. 227.
- Devkota, Govinda Prasad. 1986. Utilization of *Eupatorium* species as an alternative feedstock for producing biogas. Chemical Abstracts, Vol.104, 146.


- Fessenden, R.J. and Fessenden, J.S. 1983. Techniques and Experiments for Organic Chemistry, London : Willard Grant Press. 227.
- Furniss, B.S., Hannaford, A.J., Borgers, V., Smith, P.W.G. and Talchell, A.R. 1989. Vogel's Textbook of Practical Organic chemistry, 5<sup>th</sup> ed., New York : John Wiley and Sons, 430.
- Gonzalez, G.A., Diaz, J.G., Dominguez, B. and Bermejo, B. J. 1987. Triterpenes and steroids from *Eupatorium adenophorum* Spreng. Chemical Abstracts. Vol. 106, 420.
- Harada, J., Paisooksantivatana, Y. and Zungsontiporn, S. 1987. Weeds in the Highlands of Northern Thailand. Bangkok : Mass & Media Co.,Ltd., 126.
- Harborne, J.B. 1984. Phytochemical Method. A Guide to Modern Techniques of Plant Analysis, 2<sup>nd</sup> ed., London : Chapman and Hall. 56.
- Hostettmann, K., Terreux,C., Marston, A. and Potterat, O. 1997. The Role of Planar Chromatography in the Rapid Screening and Isolation of Bioactive Compounds from Medicinal Plants. Journal of Planar Chromatography. Vol. 10, 251.
- Jain, S.K. and Robert A. De Filippis. 1991. Medicinal Plants of India Vol.1. Reference publications.Inc. 649.
- Jiang, Yunhua, Yu, Xiaohua. 1987. Production of marsh gas from *Eupatorium adenophorum* Spreng. Chemical Abstracts. Vol.106, 175.
- Kalsi, P.S., Arora G.S. and Ghulati, R.S. 1979. New Atipodal Sesquiterpene Alcohols from Vertiver Oil. Phytochemistry. Vol. 18, 1223.
- Maneechakr, P. 1994. Chemical Constituents and Biological Activities of *Spaeranthus africanus* Linn. Master's Thesis, Department of Chemistry, Graduate School, Chulalongkorn University.
- Meyer, B.N., Ferrigni, N.R., Putnam, J.E., Jacobson, L.B., Nichols, D.E. and McLaughlin, J.L. 1982. Brine Shrimp : A Convenient General Bioassay for Active Plant Constituents. Planta Medica. Vol. 45, 31.

- Oelrichs, P.B., Calanasan, C.A., MacLeod, J.K., Seawright, A.A. and Ng, J.C. 1996. Isolation of a compound from *Eupatorium adenophorum*(Spreng.)[*Ageratina adenophora* (Spreng.)] causing hepatotoxicity in mice. Chemical Abstracts. Vol. 124, 395.
- Paisooksantivatana, Y. 1994. Medicinal Plant in Asteraceae. Department of Horticulture, Faculty of Agriculture, Kasetsart University. (Mimeographed)
- Parsons, W.T. and Cuthberson, E.G. 1992. Noxious Weed of Australia. Sydney : Inkata Press, 692.
- Proksch, P., Wray, V., Ismant, M.B. and Rahaus, I. 1990. Ontogenetic Variation of Biologically Active Natural Products in *Ageratina adenophora*. Phytochemistry. Vol. 29, 453.
- Proksch, P., Palmer, J. and Hartmann, T. 1986. Metabolism and bioconversion of chromene derivatives in *Ageratina adenophora* (Asteraceae). Chemical Abstracts. Vol. 105, 455.
- Rahalison, L., Hamburger, M., Hostettmann, K., Monod, M. and Frenk, E. 1991. A Bioautographic Agar Overlay Method for the Detection of Antifungal Compounds from Higher Plants. Phytochemical Analysis. Vol. 2, 199.
- Raksilp, T. 1995. Chemical Constituents of the Tuberous Roots of *Butea superba* Roxb. Master's Thesis, Department of Chemistry, Graduate School, Chulalongkorn University.
- Randerath, K. 1996. Thin-Layer Chromatography, nd rev., USA, Academic Press. 264.
- Rice, E.L. 1974. Allelopathy. New York : Academic Press, 331.
- Rice, E.L. 1979. Allelopathy – an uptake. Bot. Rev. 109.
- Rizvi, S.J.H. and Rizvi, V. 1992. Allelopathy : Basic and Applied Aspects. London : Chapman & Hall, 380.
- Sajise, P.E. and Lales, J.S. 1976. Allelopathy in a Mixture of Cogon (*Imperata cylindrica*) and *Stylosanthes guyanensis*. Weed Abstract. Vol. 25, 326.
- Sani, Y., Harper, P.A., Cook, R.L., Seawright, A.A. and Ng, J.C. 1992. The Toxicity of *Eupatorium adenophorum* for the Liver of the Mouse. Weed Abstract. Vol. 43, 837.

- Scott, K.N. 1972. Carbon-13 Nuclear Magnetic Resonance of Biologically Important Aromatic Acids. I. Chemical Shifts of Benzoic Acid and Derivatives. Journal of American Chemistry Society. Vol. 94(24), 8564.
- Shriner, R.P., Fuson, R.C., Curtin, D.Y. and Morril, T.C. 1980. The Systematic Identification of Organic Compounds, 6<sup>th</sup> ed. John Wiley & Sons, New York. 178.
- Shukla, V.S., Barua, N.C., Chowdhury, P.K., Sharma, R.P. and Bordoloi, M. 1986. Absolute Stereochemistry of the Cadinenes from *Eupatorium trapezoideum*. Tetrahedron. Vol. 42(4), 1157.
- Shukla, V.S., Barua, N.C., Chowdhury, P.K., Sharma, R.P. and Baruah, J.N., 1983. Structure of a New and a Degraded (des-isopropyl)-cadinene. Chemistry and Industry. 863.
- Silverstein, R.M., Bassler, G.C. and Morrill, J.C. 1981. Spectrometric Identification of Organic Compounds, 4<sup>th</sup> ed., John Wiley & Sons, USA. 304.
- Smitinand, T. 1980. Thai Plant Name (Botanical names-Vernacular names), 2<sup>nd</sup> ed. Bangkok : Royal Forest Department. 354.
- Solis, N.P., Wright, W.C., Anderson, M.M., Gupta, P.M. and Phillipson, D.J. 1992. A Microwell Cytotoxicity Assay Using *Artemia salina* (Brine Shrimp). Planta Medica. Vol. 59, 250.
- Talwar, K.K., Singh, I.P. and Kalsi, P.S. 1992. A Sesquiterpenoid with Plant Growth Regulatory Activity. Phytochemistry. Vol. 31(1), 336.
- Tripathi, R.S., Singh, R.S. and Rai, J.P.N. 1982. Allelopathic Potential of *Eupatorium adenophorum* a Dominant Ruderal Weed of Meghalaya. Weed Abstract. Vol. 31, 1780.
- Wiboonpun, N. 1996. Chemical Constituents of the Root of *Asparagus racemosus* Willd. Master's Thesis, Department of Chemistry, Graduate School, Chulalongkorn University.
- Xu, Y., Shan, X., Wang, Z. and Wu, M. 1988. A brief report of the chemical constituents from *Eupatorium adenophorum*. Chemical Abstracts. Vol. 109, 428.

Zungsontiporn, S. 1995. Allelopathic Effect of Some Tropical Plants. Ph.D's Thesis,  
Department of Science of Plant and Animal Production, The United Graduate  
School, Tokyo University of Agriculture and Technology.





## **Appendices**

สถาบันวิทยบริการ  
จุฬาลงกรณ์มหาวิทยาลัย

### Anticell Lines Cytotoxicity Test

**Table 1** The results of cytotoxicity test against Leukemia carcinoma (HL-60) cell line of some Compositae weed crude extracts

Plant	Solvent	Concentration ( $\mu\text{g/mL}$ )			Estimation
		1	10	100	
<i>E. adenophorum</i>	Hexane	-44.03	-3.33	77.52	+
	EtOH	-18.30	4.96	42.50	-
	CH <sub>2</sub> Cl <sub>2</sub>	-20.57	5.79	80.86	+
<i>E. odoratum</i>	BuOH	-0.07	1.12	40.21	-
<i>A. conyzoides</i>	Hexane	-17.85	6.39	11.31	-
	EtOH	16.99	14.25	29.47	-
<i>S. africanus</i>	EtOH	3.43	40.10	79.37	+

**Table 2** The results of cytotoxicity test against Gastric carcinoma (BGC-823) cell line of some Compositae weed crude extracts

Plant	Solvent	Concentration ( $\mu\text{g/mL}$ )			Estimation
		1	10	100	
<i>E. adenophorum</i>	Hexane	9.79	9.71	20.27	-
	EtOH	9.53	21.40	74.01	+
	CH <sub>2</sub> Cl <sub>2</sub>	-1.58	13.45	88.41	+
<i>E. odoratum</i>	BuOH	-4.76	0.25	44.35	-
<i>A. conyzoides</i>	EtOH	-2.17	-3.10	65.05	+
<i>S. africanus</i>	EtOH	-0.83	-4.98	-	-

**Table 3** The results of cytotoxicity test against Erythroleukemia carcinoma (K-562) cell line of some Compositae weed crude extracts

Extract	Concentration ( $\mu\text{g/mL}$ )			Estimation
	1	10	100	
<i>E. adenophorum</i> - Hexane	11.62	18.64	80.70	+
EtOH	-20.17	-18.72	-8.77	-
<i>E. odoratum</i> - BuOH	13.59	19.73	30.04	-
<i>A. conyzoides</i> - Hexane	3.50	2.63	48.46	-
EtOH	14.69	34.64	22.14	-

**Table 4** The results of cytotoxicity test against Nasopharyngeal carcinoma (KB) cell line of some Compositae weed crude extracts

Extract	Concentration ( $\mu\text{g/mL}$ )			Estimation
	1	10	100	
<i>E. adenophorum</i> - Hexane	8.93	20.01	23.07	-
EtOH	-2.79	-1.86	54.74	+
<i>E. odoratum</i> - BuOH	-4.46	-6.56	2.35	-
<i>A. conyzoides</i> - EtOH	13.75	47.37	62.56	+
<i>S. africanus</i> - EtOH	1.08	3.23	48.79	-

**Table 5** The results of cytotoxicity test against Bladder carcinoma (BIU) cell line  
*E. adenophorum* crude extract

Extract	Concentration ( $\mu\text{g/mL}$ )			Estimation
	1	10	100	
Hexane	22.16	36.97	41.29	-
EtOH	29.29	36.98	56.48	++
CH <sub>2</sub> Cl <sub>2</sub>	8.45	16.74	34.48	-
EtOAc	2.99	19.86	39.93	-
BuOH	19.38	19.88	22.37	+

**Table 6** The results of cytotoxicity test against Bladder carcinoma (BIU) cell line of some Compositae weed crude extracts

Extract	Concentration ( $\mu\text{g/mL}$ )			Estimation
	1	10	100	
<i>E. adenophorum</i> - Hexane	22.16	36.97	41.29	-
<i>E. odoratum</i> - BuOH	-5.38	-5.13	55.58	+
<i>A. conyzoides</i> - Hexane	33.53	38.17	53.61	+
EtOH	13.75	47.37	62.56	+

**Table 7** The results of cytotoxicity test against Erythroleukemia carcinoma (K-562) cell line of *E. adenophorum* crude extract

Extract	Concentration ( $\mu\text{g/mL}$ )			Estimation
	1	10	100	
Hexane	11.62	18.64	80.70	+
EtOH	-20.17	-18.72	-8.77	-
CH <sub>2</sub> Cl <sub>2</sub>	4.12	-1.31	35.08	-
EtOAc	11.97	16.66	52.26	+
BuOH	8.11	-8.33	21.27	-

**Table 8** The results of cytotoxicity test against Colon carcinoma (HCT-8) cell line of *E. adenophorum* crude extract

Extract	Concentration ( $\mu\text{g/mL}$ )			Estimation
	1	10	100	
Hexane	3.51	2.31	1.66	-
EtOH	5.81	2.08	88.14	+
CH <sub>2</sub> Cl <sub>2</sub>	4.82	3.40	89.13	+

**Table 9** The results of cytotoxicity test against Nasopharyngeal carcinoma (KB) cell line of *E. adenophorum* crude extract

Extract	Concentration ( $\mu\text{g/mL}$ )			Estimation
	1	10	100	
Hexane	8.93	20.01	23.07	-
EtOH	-2.79	-1.86	54.74	+
EtOAc	3.66	5.05	13.03	-

**Table 10** The results of cytotoxicity test against Gastric carcinoma (BGC-823) cell line of *E. adenophorum* crude extract

Extract	Concentration ( $\mu\text{g/mL}$ )			Estimation
	1	10	100	
Hexane	9.79	9.71	20.27	-
EtOH	9.53	21.40	74.01	+
CH <sub>2</sub> Cl <sub>2</sub>	-4.98	-5.87	40.27	-
EtOAc	-14.55	-11.70	50.88	+

**Table 11** The results of cytotoxicity test against Leukemia carcinoma (HL-60) cell line of *E. adenophorum* crude extract

Extract	Concentration ( $\mu\text{g/mL}$ )			Estimation
	1	10	100	
Hexane	-16.23	13.85	53.01	+
EtOH	-9.03	-4.97	16.13	-
CH <sub>2</sub> Cl <sub>2</sub>	34.55	40.84	83.71	++
EtOAc	32.77	39.82	63.16	+
BuOH	19.38	19.88	22.37	-

**Table 12** The results of cytotoxicity test against Hepatocellular carcinoma (Bel-7402) cell line of *E. adenophorum* crude extract

Extract	Concentration ( $\mu\text{g/mL}$ )			Estimation
	1	10	100	
Hexane	4.69	4.07	3.27	-
EtOH	3.78	10.91	67.07	+
CH <sub>2</sub> Cl <sub>2</sub>	7.31	12.06	87.67	+

**Table 13** The results of cytotoxicity test against Leukemia carcinoma (HL-60) cell line of *E. adenophorum* dichloromethane crude extract

Extract	Concentration ( $\mu\text{g/ml}$ )			Estimation
	1	10	100	
IIIA	-5.89	-7.96	-9.82	-
IIIB	-2.17	6.72	43.32	-
IIIC	-8.06	-3.10	77.35	-
IIIF	-12.30	-7.54	64.42	+

**Table 14** The results of cytotoxicity test against Colon carcinoma (HCT-8) cell line of *E. adenophorum* dichloromethane crude extract

Extract	Concentration ( $\mu\text{g/ml}$ )			Estimation
	1	10	100	
IIIA	-13.94	6.69	39.51	-
IIIB	18.11	15.58	92.86	+
IIIC	19.42	17.56	95.86	+
IIIF	11.85	16.02	81.77	+

**Table 15** The results of cytotoxicity test against Hepatocellular carcinoma (Bel-7402) cell line of *E. adenophorum* dichloromethane crude extract

Extract	Concentration ( $\mu\text{g/ml}$ )			Estimation
	1	10	100	
IIIA	8.98	12.32	60.26	+
IIIB	13.79	17.07	86.20	+
IIIC	7.25	20.73	66.30	+
IIIF	7.55	11.26	69.14	+

**Table 16** The results of cytotoxicity test against Gastric carcinoma (BGC-823)  
cell line of *E. adenophorum* dichloromethane crude extract

Extract	Concentration ( $\mu\text{g}/\text{ml}$ )			Estimation
	1	10	100	
III A	8.03	26.26	59.71	+
III B	7.85	21.21	67.66	+
III C	14.61	-15.48	63.03	+
III F	10.79	24.53	66.08	+

**Table 17** The results of cytotoxicity test against Leukemiar carcinoma (HL-60)  
cell line of *E. adenophorum* ethyl acetate crude extract

Extract	Concentration ( $\mu\text{g}/\text{mL}$ )			Estimation
	1	10	100	
IV A	-4.02	10.06	71.58	+
IV B	-10.96	13.64	9.84	-
IV C	-46.53	-15.26	49.76	-
IV D	-9.39	12.52	22.59	-
IV E	-1.56	44.51	41.61	-
IV F	14.59	29.53	64.59	+

**Table 18** The results of cytotoxicity test against Nasopharyngeal carcinoma (KB) cell line of *E. adenophorum* ethyl acetate crude extract

Extract	Concentration ( $\mu\text{g/ml}$ )			Estimation
	1	10	100	
IVA	8.20	27.02	97.42	+
IVB	4.85	19.89	49.61	-
IVC	16.89	16.29	76.45	+
IVD	19.17	17.49	46.97	-
IVE	7.01	17.67	47.99	-
IVF	18.29	20.49	48.41	-

**Table 19** The results of cytotoxicity test against Colon carcinoma (HCT-8) cell line of *E. adenophorum* ethyl acetate crude extract

Extract	Concentration ( $\mu\text{g/ml}$ )			Estimation
	1	10	100	
IVA	0.78	2.21	95.61	+
IVB	2.96	93.81	92.01	++
IVC	0.18	2.92	92.72	+
IVD	-0.93	-1.12	92.49	+
IVE	-1.65	-1.35	91.14	+
IVF	-2.10	-1.01	94.33	+

**Table 20** The results of cytotoxicity test against Hepatocellular carcinoma (Bel-7402) cell line of *E. adenophorum*: ethyl acetate crude extract

Extract	Concentration ( $\mu\text{g/ml}$ )			Estimation
	1	10	100	
IVA	3.77	1.65	93.90	+
IVB	-3.54	59.16	79.97	++
IVC	-2.42	-1.53	85.95	+
IVD	-5.78	-1.33	85.32	+
IVE	-6.68	-2.00	85.12	+
IVF	-6.09	-2.51	84.97	+

**Table 21** The results of cytotoxicity test against Gastric carcinoma (BGC-823) cell line of *E. adenophorum*: ethyl acetate crude extract

Extract	Concentration ( $\mu\text{g/ml}$ )			Estimation
	1	10	100	
IVA	9.02	13.64	71.54	+
IVB	11.64	13.18	36.16	-
IVC	13.95	7.71	66.07	+
IVD	13.56	7.01	40.24	-
IVE	19.19	12.25	34.46	-
IVF	17.11	15.80	38.47	-

Note : The results suggested by Beijing Medical University, Beijing, China

++ good

+ fair

- no activity

## VITA

Miss Suttijit Sriwatcharakul was born on March 17, 1973 in Bangkok, Thailand. She received the Bachelor Degree of Science(Agriculture) at Kasetsart University in 1995. Since 1995, she has been a graduate student studying Biotechnology in Chulalongkorn University. She was supported by a research grant for her Master Degree's thesis from the graduate school, Chulalongkorn University.



สถาบันวิทยบริการ
จุฬาลงกรณ์มหาวิทยาลัย