

REFERENCES

- Allison, A.C. (1958). Genetical and clinical significance of haptoglobins. Proc.Roy.Soc.Med., 51:641.
- Allison, A.C. (1959). Genetic control of human haptoglobin synthesis. Nature, 183 : 1312.
- Allison, A.C. and ap Rees, W. (1957). The binding of haemoglobin by plasma proteins (haptoglobins). Brit.med.J., 2:1137.
- Allison, A.C. and Barnicot, N.A. (1960). Haptoglobins and transferrins in some East African people. Acta Genet. (Basel), 10:17.
- Allison, A.C. and Blumberg, B.S. (1958). The genetically determined serum haptoglobin in rheumatoid arthritis. Arthritis Rheum., 1:239.
- Allison, A.C., Blumberg, B.S. and ap Rees, W. (1958). Haptoglobin types in British, Spanish Basque and Nigerian African populations. Nature, 181:824.
- Alper, C.A., Peters, J.H., Birtch, A.G. and Gardner, F.H. (1965). Haptoglobin synthesis. I In vivo studies of the production of haptoglobin, fibrinogen and  $\alpha_2$ -globulin by the canine liver. J.Clin.Invest., 44:574.
- Anastassea-Vlachou, C., Kattamis, C., Lagos, P., Konstantoulakis, M. and Matsaniotis, N. (1973). Haptoglobin and  $\alpha_2$ -macroglobulin levels in children with rheumatoid arthritis. Clin.Chim.Acta, 44:259.

Areekul, S. and Chantachum, Y. (1972). Serum haptoglobin levels in normal Thais. Southeast Asian J.Trop. Med. Pub.Hlth., 3:286.

Areekul, S., Chantachum, Y. and Matrakul, D. (1974). Serum haptoglobin levels in patients with liver diseases. Southeast Asian J.Trop.Med.Pub.Hlth., 5:601.

Areekul, S., Chantachum, Y., Matrakul, D. and Viravan, C. (1972). Serum haptoglobin levels in malaria. Southeast Asian J.Trop.Med.Pub.Hlth., 3:505 .

Areekul, S., Devakul, K., Chantachum, Y., Boonyananta, C., Egoramaiphol, S. and Viravan, C. (1971). Gastrointestinal protein loss in patients with hookworm infection. J.Med.Ass.Thailand, 54:28.

Barnicot, N.A., Garlick, J.P. and Roberts, D.F. (1960). Haptoglobin and transferrin inheritance in northern Nigerians. Ann.Hum.Genet., 24:171.

Blackwell, R.Q. and Thephusdin, C. (1963). Distribution of haptoglobin among Thais. Nature, 197:503.

Blackwell, R.Q., Chen, H.H. and Chen, H.C. (1964). Haptoglobin distribution in a Filipino population. Nature, 202 : 814.

Blumberg, B.S. and Gentile, Z. (1961). Haptoglobins and transferrins of the two tropical populations. Nature, 189:897.

Blumberg, B.S., Kuvin, S.F., Robinson, J.C., Teitelbaum, J.M. and Contacos, P. (1963). Alterations in haptoglobin levels. J.A.M.A., 184:1021.

Bogdanovich, L. (1968). Serum haptoglobin as an index of the activity of pulmonary tuberculosis. Probl.Tuberk., 46:65.

Borglin, N.E. and Nyman, M. (1961). Effect of estrogens on the haptoglobin level in the blood. Scand.J.Lab.Clin.Invest., 13:107.

Bovornkitti, S. (1962). Serum protein changes in tuberculosis with particular reference to alpha-2-globulin. Amer.Rev.Resp.Dis., 85:56.

Brus, I. and Lewis, M. (1959). The haptoglobin content of serum in haemolytic anaemia. Brit.J.Haemat., 5:348.

Cheftel, R.I., Cloarec, L., Moretti, J., Rafelson, M. and Jayle, M.F. (1960). Etude de l'haptoglobine. III. Composition chimique de la partie glucidique. Bull.Soc.Chim.Biol., 42:993.

Chiancone, E., Alfèn, A., Ioppolo, C., Vecchini, P., Agro, Finazzi, A., Wyman, J. and Antonini, E. (1968). Studies on the reaction of haptoglobin with haemoglobin and haemoglobin chains. I. Stoichiometry and affinity. J.Mol.Biol., 34:347.

Connell, G.E. and Smithies, O. (1959). Human haptoglobins : Estimation and population. Biochem.J., 72:115.

Crockson, R., Payne, J., Ratcliff, A. and Soothill, J.F. (1966).

Time sequence of acute phase reactive proteins  
following surgical trauma. Clin. Chim. Acta, 14:435.

Culliford, B.J. (1963). Haptoglobins and transferrins in forensic  
bloodstain. Nature, 198:797.

Ferris, T.G., Easterling, R.E.; Nelson, K.J. and Budd, R.E.  
(1966). Determination of serum haemoglobin binding  
capacity and haptoglobin type by acrylamide gel  
electrophoresis. Am. J. clin. Path., 46:385.

Galatius-Jensen, F. (1957). Further investigations of the  
genetic mechanism of the haptoglobins. Acta Genet.  
(Basel), 7:549.

Giblett, E.R. (1968). The haptoglobin system. Series Haematol., 1:3.

Giblett, E.R., Motulsky, A.G. and Fraser, G.R. (1966). Population  
genetics in the Congo. IV Haptoglobin and transferrin  
serum groups in the Congo and in other African  
populations. Am. J. Hum. Genet., 18:533.

Gordon, S. and Bearn, A.G. (1966). Haemoglobin binding  
capacity of isolated haptoglobin polypeptide chains.  
Proc. Soc. Exp. Biol. Med., 121:846.

Hall, R. (1964). A simple and rapid method for estimation  
of haptoglobins in serum. J. Med. Lab. Technol., 21:64.

Hommes, F.A. (1959). A new method for the quantitative  
determination of haptoglobin. Clin. Chim. Acta, 4:707.

Ivanyi, J., Sochman, J. and Ivanyi, P.(1961). The haptoglobin level in haemoblastoses. Acta Haematol. (Basel), 25:98.

Javid, J.(1967). Human serum haptoglobin : A brief review. Seminars Hematol., 4:35.

Javid, J. and Horowitz, H.I.(1960). An improved technic for the quantitation of serum haptoglobin. Am.J.clin. Path., 34:35.

Jayle, M.F.(1940). Evaluation oneliclaire de l'haptoglobine par une reaction enzymatique de peroxidation. C.R.Acad.Sci., 211 : 574.

Jayle, M.F. and Boussier, G.(1955). Les seromucoides de sang, leurs relations avec mucoproteines de la substance fondamentale tissu conjonctif. Exposes Ann.Biochem.Med., 17:157.

Jayle, M.F. and Morretti, J.(1962). Haptoglobin : biochemical, genetic and physiopathologic aspects. in Moore, C.V. and Brown, E. B.(eds.) Progress in Haematology Vol.3, p.342. Grune and Stratton, New York.

Kashiwagi, N., Groth, C.G. and Starzl, T.E. (1968). Changes in serum haptoglobin and group specific component after orthotopic liver homotransplantation in humans. Froc.Soc.Exp.Biol.Med., 128:247.

Killander, J.(1964). Separation of heme- and haemoglobin-binding plasma proteins, ceruloplasmin and albumin by gel filtration. Biochim.Biophys.Acta, 93:1.

Kirk, R. L. and Lai L. Y. C. (1961). The distribution of haptoglobins and transferrin in South and Southeast Asia. Acta Genet. (Basel), 11:97.

Kluthe, R., Faul, J. and Hempel, H. (1965). Quantitative estimation of human serum haptoglobin by an immunological method. Nature, 205:93.

Kobiela, J., Turowska, B. and Baniowski, A. (1966). The level of haptoglobins in serum of patients with rheumatoid arthritis. Pol. Med. J., 5:1004.

Krauss, S. and Sarcione, E. J. (1964). Synthesis of serum haptoglobin by the isolated perfused rat liver. Biochim. Biophys. Acta 90:301.

Krauss, S., Schrott, M. and Sarcione, E. J. (1966). Haptoglobin in Hodgkin's disease. Am. J. Med. Sci., 252:185.

Lathem, W. and Jensen, W. N. (1959). Plasma haemoglobin-combining capacity in sickle-cell disease. Blood, 11:493.

Lathem, W. and Worley, W. E. (1959). The distribution of extracorporeal haemoglobin in circulating plasma. J. Clin. Invest., 38:474.

Laurell, C. B. (1959). Purification and properties of different haptoglobin. Clin. Chim. Acta, 4:79.

Laurell, C. and Nyman, M. (1957). Studies on the serum haptoglobin level in haemoglobinaemia and its influence on the renal excretion of haemoglobin. Blood, 12:493.

Lionetti, F.J., Valeri, C.R., Bond, J.C. and Fortier, N.L. (1964).

Measurement of haemoglobin-binding capacity of human serum or plasma by means of dextran gels.

J. Lab. clin. Med., 64:519.

Louderback, A.L. and Shanbrom, E. (1968). Haptoglobin electrophoresis. J.A.M.A., 206:362.

Matsunaga, E. and Marai, K. (1960). Haptoglobin types in a Japanese population. Nature, 186:320.

Mehta, S.R. and Jensen, W.N. (1960). Haptoglobins in haemoglobinopathy: a genetic and clinical study. Brit.J.Haemat., 6:250.

Miale, J.B. (1964). Typing and quantitation of serum haptoglobins. in "Haemoglobin : its precursors and metabolites" Sunderman, F.W. and Sunderman, F.W.Jr. (eds.) Lippincott.

Mueller, W.K., Handschumacher, R. and Wade, M.E. (1971). Serum haptoglobin in patients with ovarian malignancies. Obstet. & Gynec., 38:427.

Murray, R. and Connell, G.E. (1960). Elevation of serum haptoglobin in rabbit in response to experimental inflammation. Nature, 186:86.

Murray, R., Robinson, J., Dublin, T., Pitt, E.L. and Visnich, S. (1966). Haptoglobins and rheumatic fever. Brit.Med.J., 1:762.



Nakajima, H., Takemura, T., Nakajima, O. and Yamaoka, K. (1963).

Studies on heme  $\alpha$ -methenyl oxygenase I The enzymatic conversion of pyridine hemochromogen and haemoglobin-haptoglobin into a possible precursor of biliverdin. J.Biol.Chem., 238:3784.

Neel, J.V., Robinson, A.R., Zuelzer, W.W., Livingstone, F.B. and Sutton, H.E. (1961). The frequency in the  $A_2$  and fetal haemoglobin fractions in the natives of Liberia and adjacent regions, with data on haptoglobin and transferrin types. Am.J.Hum.Genet., 13:262.

Neuhaus, O. (1964). Biochemical significance of serum glycoproteins. 3. Hepatic production of alpha-1 and alpha-2 globulins responding to injury. Proc.Soc.Exp.Biol.Med., 117:244.

Noonan, J.A., Holland, N.H., and Holmes, F.C. (1966). Haptoglobin and rheumatic heart disease. Brit.Med.J., 1:1538.

Nosslin, B. and Nyman, M. (1958). Haptoglobin determination in diagnosis of haemolytic diseases. Lancet, i:1000.

Nyman, M. (1957). Haptoglobin in pernicious anaemia. Scand.J.Clin.Lab.Invest., 9:168.

Nyman, M. (1959). Serum haptoglobin : methodological and clinical studies. Scand.J.Clin.Lab.Invest., 11 : suppl. 39.

Nyman, M., Gydell, K. and Nosslin, B. (1959). Haptoglobin und erythrokinetik. Clin.Chim.Acta, 4:82.

O'Hara, H., Watanabe, K. and Wada, T. (1968). Turnover rate of serum haptoglobin in malignant neoplastic disease. Clin.Chim.Acta, 19:41.

Owen, J.A., Better, F.D. and Hoban, J. (1960). Simple method for determination of serum haptoglobin. J.Clin.Path., 13:163.

Owen, J.A., MacKay, I.R. and Got, C. (1959). Serum haptoglobin in hepatobiliary disease. Brit.Med.J., 2:1454

Owen, J.A., Padanyi, R. and Smith, H. (1961). The effect of surgery on serum haptoglobins. Proc.Ass.Clin.Biochem., 1:89.

Owen, J.A., Smith, R., Padanyi, R. and Martin, J. (1964). Serum haptoglobin in disease. Clin.Sci., 26:1.

Owen, J.A., Careu, J.P., Cowling, D.C., Hoban, J.P. and Smith, H. (1960). Serum haptoglobins in megaloblastic anaemia. Brit.J.Haemat., 6:242.

Polonovski, M. and Jayle, M.F. (1938). Existence dans le plasma sanguin d'une substance activant l'action peroxidesique de l'hémoglobine. C.R.Soc.Biol., 129:457.

Polonovski, M. and Jayle, M.F. (1940). Sur la préparation d'une nouvelle fraction des protéines plasmatiques, l'haptoglobine. C.R.Acad.Sci.(Paris) 211:517.

Poulik, M.D. (1957). Starch-gel electrophoresis in a discontinuous system of buffers. Nature, 180:1477.

Rajendran, G., Dhara, R. and Srinivasan, N.G. (1975). Serum haptoglobins in health and disease. Indian J. Med. Res., 63:525.

Ratcliff, A.P. and Hardwicke, J. (1964). Estimation of serum haemoglobin-binding capacity (haptoglobin) on Sephadex G100. J. Clin. Path., 17:676.

Robert, L., Mombelloni, P. and Crosti, P. (1961). Studies on serum haptoglobin in experimental connective tissue disorders. Proc. Soc. exp. Biol., 107:499.

Rowe, D.S. (1961). A rapid method for the estimations of serum haptoglobin. J. clin. Path., 14:205.

Sanford, R., Grimmo, A.E.P. and Lee, S.K. (1966). The haptoglobin and transferrin types of some Cantonese in Hong Kong. Vox Sang., 11:106.

Schumacher, V.G. and Schlumberger, M.D. (1963). Über veränderungen der alpha-2-globulin des serum. Deutsche Med. Wochens., 13:645.

Shim, B.S. and Bearn, A.G. (1964). Immunological and biochemical studies on serum haptoglobin. J. Exp. Med., 120:611.

Shinton, N.K., Richardson, R.W. and Williams, J.D.F. (1965). Diagnostic value of serum haptoglobins. J. Clin. Path., 18:114.

Smithies, O. (1955). Zone electrophoresis in starch gels : Group variations in the serum proteins of normal human adults. Biochem.J., 61:629.

Smithies, O. (1959). An improved procedure for starch-gel electrophoresis further variations in the serum proteins of normal individuals. Biochem.J., 71:585.

Smithies, O. and Walker, N.F. (1955). Genetic control of some serum proteins in normal humans. Nature, 176:1265.

Smithies, O. and Walker, N.F. (1956). Notation for serum proteins groups and the genes controlling their inheritance. Nature, 178:694.

Smithies, O., Connell, G.E. and Dixon, G.H. (1962). Chromosomal rearrangements and the evaluation of haptoglobin genes, Nature, 196:232.

Studnitz, W.V. and Nyman, M. (1957). Effect of androgen on alpha-2-globulins in women. J.Clin.Endocrinol., 17:910.

Sutton, H.E., Neel, J.V., Livingstone, F.B., Binson, G., Kunstadter, P. and Trombley, L.E. (1959). The frequencies of haptoglobin types in five populations. Ann.hum.Genet., 23:175.

Thephusdin, C., Blackwell, B., Haung, J. and Blackwell, R.Q. (1965). Comparison of the distribution of haptoglobin and haemoglobin types in a Thai population. Israel J.Med.Sci., 1:768.

Turowska, B. (1969). The value of haptoglobin level determination in synovial exudates in the diagnosis of rheumatic disease. Rheumatologica, 7:57.

Tuttle, A.H. (1955). Demonstration of haemoglobin reactive substance in human serum. Science, 121 : 701.

Valeri, C.R., Bond, J.C., Fowler, K. and Sobuck , J. (1965). Quantitation of serum haemoglobin-binding capacity using cellulose acetate membrane electrophoresis. Clin.Chim.Acta, 11:581.

Whitten, C.F. (1962). Studies on serum haptoglobin : a functional inquiry. New Engl.J.Med., 266:529.

William, R., Speyer, B.E. and Billing, B.H. (1961). Serum haptoglobin in liver disease. Gut, 2:297.

Wright, S. (1956). Applied physiology. Oxford University Press, London, 9<sup>th</sup> edition p.1077.

VITA

Name: Mrs.Uma Kittiyanee

Education: Bachelor of Science in Pharmacy  
(Second Class Honour) in 1961,  
University of Medical Sciences,  
Bangkok, Thailand.

Position and Site of the Employer's Office:

Lecturer and Instructor on Pharmacology  
and Toxicology, Department of Pharmacology,  
Faculty of Pharmaceutical Sciences,  
Chulalongkorn University,  
Bangkok, Thailand.

