

# Overweight and obesity and the associated lifestyle factors in Thai adult individuals who came for service at King Chulalongkorn Memorial Hospital during August-September 2004

Suchat Suppapitiporn \*

Jinda Udompanyavit \* Siriluck Suppapitiporn \*\*

**Suppapitiporn S, Udompanyavit J, Suppapitiporn S. Overweight and obesity and the associated lifestyle factors in Thai adult individuals who came for service at King Chulalongkorn Memorial Hospital during August-September 2004. Chula Med J 2006 Sep; 50(9): 623 - 39**

**Objective** : *To study the prevalence of overweight and obesity with the new WPRO (Who Health Organization Western Pacific Region) BMI criteria for overweight and obesity in individuals who came for service at King Chulalongkorn Memorial Hospital and to determine the associated lifestyle in those individual.*

**Method** : *A cross-sectional study using a self-reported questionnaire asking information on general sociodemographic characteristics, underlying disease and lifestyle .Body weight and height were recorded and used for calculation of BMI. The sample consisted of 403 individuals who came for OPD service at King Chulalongkorn Memorial Hospital during August to September 2004.*

\* Department of Outpatient, Thai Red Cross Society, King Chulalongkorn Memorial Hospital

\*\* Department of Psychiatry, Faculty of Medicine, Chulalongkorn University

**Results** : *The prevalence for overweight and obesity according to the new criteria issued by the WPRO was 43.9 %. The study both two groups found to have nearly the same proportion of activity; low work activity level 39.7 % and high work activity level 32.0 %. Most had low leisure time activity level ( 53.1 %), sleep 6-8 hours (71.5 %), watch 1-2.5 hours of TV (43.7 %). About half of them were sitting while watching TV (53.1 %) but 23.3 % were also eating. Most of subjects had never smoke (78.4 %) and never drink (66.5 %). The factors which were significantly associated with overweight and obesity were male, age above 35 years, marital status, lower education, cardiovascular and diabetes underlying disease and moderate or high work activity level. The predictive variables from multiple logistic regression analysis were sex, age and work activity level ( $R^2 = .162$ ).*

**Conclusion** : *Overweight and obesity according to the criteria of the WPRO had increased to 43.9 % almost half in the population studied. Factors associated with overweight and obesity were sex, age above 35 years, marital status, education, underlying disease and work activity level. Recognition and prevention of overweight and obesity particularly in those with associated factors may be beneficial for reducing burden of overweight and obesity including other following health problems.*

**Keywords** : *Overweight, Obesity, Lifestyle.*

Reprint request: Suppavitiporn Su. Department of Outpatient , Thai Red Cross Society,  
King Chulalongkorn Memorial Hospital, Bangkok 10330, Thailand.

Received for publication. October 10, 2005.

สุชาติ ศุภปิติพร, จินดา อุดมปัญญาวิทย์, ศิริลักษณ์ ศุภปิติพร. ภาวะน้ำหนักเกินและอ้วน  
กับปัจจัยด้านการดำเนินชีวิตของผู้ป่วยนอกที่มาใช้บริการที่โรงพยาบาลจุฬาลงกรณ์ ในช่วง  
เดือนสิงหาคมถึงกันยายน พ.ศ.2547. จุฬาลงกรณ์เวชสาร 2549 ก.ย; 50(9): 623 – 39

**วัตถุประสงค์** : เพื่อศึกษาความชุกของภาวะน้ำหนักเกินและอ้วน โดยใช้เกณฑ์การแบ่งระดับ  
ใหม่ของชาวเอเชีย คือดัชนีมวลกายตั้งแต่ 23 กิโลกรัมต่อตารางเมตรขึ้นไป และ  
เพื่อค้นหาความสัมพันธ์ระหว่างปัจจัยด้านการดำเนินชีวิตกับภาวะน้ำหนักเกิน  
และอ้วนในผู้ที่มาใช้บริการที่โรงพยาบาลจุฬาลงกรณ์

**วิธีการวิจัย** : การวิจัยเชิงพรรณนา ณ จุดเวลาหนึ่ง ศึกษาผู้ป่วยอายุ 20 ปีขึ้นไป ซึ่งมาใช้  
บริการที่แผนกผู้ป่วยนอก โรงพยาบาลจุฬาลงกรณ์ ในช่วงเดือนสิงหาคมและ  
กันยายนปี พ.ศ. 2547 โดยใช้แบบสอบถามเกี่ยวกับปัจจัยพื้นฐาน การดำเนิน  
ชีวิตประจำวัน วัดส่วนสูงและน้ำหนักเพื่อคำนวณดัชนีมวลกาย

**ผลการวิจัย** : ความชุกของภาวะน้ำหนักเกินและอ้วน โดยใช้เกณฑ์กำหนดแบ่งระดับความ  
อ้วนของ WPRO (World Health Organization Western Pacific Region) อยู่ที่  
43.9 % กลุ่มผู้เข้าร่วมวิจัยมีระดับการทำงานที่ใช้พลังงานน้อย 39.7 % ใกล้เคียง  
กับกลุ่มที่มีระดับการทำงานที่ต้องใช้พลังงานสูง 32.0 % ส่วนใหญ่มีกิจกรรม  
ยามว่างใช้ระดับพลังงานต่ำ 53.1 % ดูโทรทัศน์ 1-2.5 ชั่วโมงต่อวัน 43.7 %  
โดยที่ส่วนใหญ่จะนั่งอยู่หน้าจอโทรทัศน์และมีการรับประทานอาหารไปด้วย  
23.3 % ส่วนใหญ่นอนหลับ 6 – 8 ชั่วโมงต่อคืน 71.5 % รับประทานอาหาร  
หนักปานกลางถึงหนักมาก (57.8 % , 30.8 % ตามลำดับ) ไม่เคยสูบบุหรี่  
และไม่เคยดื่มสุรา(78.4 % , 66.5 % ตามลำดับ) พบความสัมพันธ์ระหว่างภาวะ  
น้ำหนักเกินและอ้วนกับปัจจัยพื้นฐานอย่างมีนัยสำคัญทางสถิติ ในด้านเพศชาย  
อายุมากกว่า 35 ปี สถานภาพสมรส ระดับการศึกษาต่ำ มีโรคประจำตัวเป็น  
เบาหวานและโรคหลอดเลือดหัวใจ และระดับการทำงานที่ต้องใช้พลังงานระดับ  
กลางถึงสูง จากการวิเคราะห์ถดถอยเชิงพหุ พบว่าตัวแปรที่มีนัยสำคัญทางสถิติ  
คือเพศ อายุ และระดับการทำงาน ซึ่งอธิบายการเปลี่ยนแปลงของภาวะน้ำหนัก  
เกินและอ้วนได้ร้อยละ 16.2

**สรุป** : ความชุกของภาวะน้ำหนักเกินและอ้วนตามเกณฑ์ใหม่ของเอเชีย (WPRO) พบ  
43.9 % เกือบครึ่งหนึ่งของประชากรที่ศึกษา แสดงให้เห็นถึงแนวโน้มการเพิ่มขึ้น  
ของปัญหาสุขภาพที่อาจเกิดตามมา ดังนั้นการตระหนักและวางแผนป้องกัน  
โดยเฉพาะในผู้ที่มีปัจจัยสัมพันธ์ดังกล่าว คงมีส่วนช่วยลดปัญหาภาวะน้ำหนัก  
เกินและอ้วนรวมถึงปัญหาสุขภาพต่าง ๆ

**คำสำคัญ** : ภาวะน้ำหนักเกิน, อ้วน, การดำเนินชีวิต.

In the past decade, the problem of overweight and obesity had been widespread and increasing around the world. Evidence had shown that being overweight and obese can lead to numerous serious disorders and increased mortality. In 1998, WHO had established a BMI (body mass index) classification for overweight and obesity, enlisting those to be overweight when  $BMI \geq 25$  and obese when  $BMI \geq 30$ .<sup>(1,2)</sup> However in the year 2000, researches<sup>(3-12)</sup> had found Asians to develop cardiovascular disorders and metabolic syndromes associated with being overweight and obese at a BMI lower than classified by WHO and thus led to a new, yet unofficial classification by the WPRO (The Regional Office for the Western Pacific, WHO, the International Association for the Study of Obesity and International Obesity Task Force) for overweight and obesity, enlisting those to be overweight when  $BMI \geq 23$  and obese when  $BMI \geq 25$ . In a study of 1513 Hong Kong Chinese, the risk of diabetes, Hypertension, dyslipidaemia and albuminuria starts to increase at a BMI of about  $23 \text{ kg/m}^2$  which is lower than the current WHO BMI cut-off used to define an increase in morbidity among Europeans and also demonstrated in data from the 1992 national health survey in Singapore. The WPRO made it clear risk is a continuum with increasing BMI and that cut-off points are merely a convenience for public health and clinical use.<sup>(13)</sup> With the new criteria, it would be interesting to find out the prevalence of overweight and obesity in Thailand which may be more increased, and thus may demand the public to become more aware of the seriousness of this problem in Thailand.

Obesity is now well recognized as a disease in its own right. The obese have an elevated risk from all cause mortality. Most evidence suggests a J-shaped relationship between BMI and mortality, with the obese having the highest risk. Obesity

is associated with insulin resistance, glucose intolerance, diabetes mellitus, hypertension, dyslipidemia, sleep apnea, arthritis, hyperuricemia, gall bladder disease, certain types of cancer, coronary artery disease, heart failure, cardiac arrhythmia, hemorrhagic stroke, menstrual irregularities and significant deterioration in emotional well-being is found to be controversial.<sup>(14-29)</sup> Overweight and obesity are multifactorial health problems, stemming from two main etiological factors: the genetic factor and the environmental factor. In Thailand, researches remain controversial for whether which factor is more dominant. However, with the environmental factor being within our ability to modify, much research abroad had been done to find out the environmental factors involved.<sup>(30,31)</sup> The environmental factor of focus for this research is the individual's lifestyle, which for people in Asia, has changed progressively in the past. No doubt had it had much influence from the western world, resulting in a more sedentary lifestyle.<sup>(32)</sup> As for Thailand, it would be interesting to document the present lifestyle of Thai people and to find out the association this lifestyle may have on the prevalence of overweight and obesity which may have increased in Thailand.

The problem of overweight and obesity must be attended to and national health programs and campaigns should be implemented to reduce the seriousness of this problem. In this research we have studied only a small sample of individuals who came for service at King Chulalongkorn Memorial Hospital. This research is intended to provide one of the basic information for creating these programs and campaigns, in hopes that in the future, the problem of overweight and obesity may be resolved in Thailand.



## Method

The population studied were individuals over 20 years of age who were legible and came for service in King Chulalongkorn Memorial Hospital from OPD which included medical and surgical department service of the Bhumibol Building and whom were randomly distributed by the nurse of each floor. The tool used for data collection was the self-reported questionnaire. Body weight and height were recorded and then calculated to be body mass index (BMI) as the unit of measurement for overweight and obesity.

The questionnaire consisted of sociodemographic characteristics, work activity, TV watching hours and activity, leisure time activity, sleeping, diet, history of smoking and drinking. The questionnaire was valid and permission was requested from the ethic board committee and the hospital administration to distribute the questionnaire for this study. After receiving permission for distribution of questionnaires, the questionnaires were randomly distributed to the subjects who were waiting to meet the physicians at the OPD, during the months of August and September 2004.

Data analysis was done by using SPSS for windows version 11.0. Descriptive data was analyzed using frequency, percentage, mean, and mode methods. Association between overweight and obesity and the sociodemographic or lifestyle of individuals was found by using the chi-square. The significant factors would further undergone multiple logistic regression analysis.

## Results

During the months of August and September 2004, the 403 completed questionnaires were collected.

Most of individuals who came for service in King Chulalongkorn Memorial Hospital were in the age range of 20-80 years old, the mean being 40.2 years old (SD = 13.95), mode being 25 years old. There were more females than male, female being 61.3 %. About half of the subjects were private employees 56.8 % (n=229). Students were noted to be few due to the selection of sample to have age group over 20 years old. The proportion of married individuals to single individuals was approximately the same, with married individuals being a little more in proportion (50.4 %). Other status, being divorced and widowed, are less than 10 % each. Over 50 % had an education above high school (56.8 %). Most had the income in the range of 10,001-50,000 bahts/month (42.7 %) and had no underlying diseases (56.7 %). Those that did have underlying diseases mostly had noncardiovascular diseases (24.3 %). The data are shown in table 1.

The prevalence for overweight and obesity according to the new criteria issued by the WPRO (BMI  $23 \text{ kg/m}^2$ ) was 43.9 % (n=177). The range of BMI in the present study was 15.61 to 42.15 with the mean 23.07 (SD 4.23).

Regarding the lifestyle, the sample was found to have the same proportion of low work activity level (39.7 %) to high work activity level (32.0 %). Most had low leisure time activity level (53.1 %), watch 1-2.5 hours of TV (43.7 %) of which most were sitting while watching TV (53.1 %), sleep 6-8 hours (71.5 %) of which the proportion of good and bad sleep was approximately the same, had moderate (57.8 %) to heavy (30.8 %) diet, never smoke (78.4 %) and never drink (66.5 %).

**Table 1.** Sociodemographic characteristics of individuals who came for service at King Chulalongkorn Memorial Hospital.

Sociodemographic characteristics	No. of person	%
<b>Sex</b>		
Male	156	38.7
Female	247	61.3
<b>Marital status</b>		
Single	164	40.7
Married	203	50.4
Divorced	15	3.7
Widowed	21	5.2
<b>Education</b>		
Below elementary	6	1.5
Elementary	59	14.6
High school	109	27.1
Above high school	229	56.8
<b>Career</b>		
Private employee	229	56.8
Government official	57	14.1
Self-employed business	49	12.2
Housekeeping/retired	53	13.2
Student	15	3.7
<b>Income (bahts/months)</b>		
0-5000	102	25.3
5001-10000	117	29.0
10001-50000	172	42.7
>50000	12	3.0
<b>Underlying disease</b>		
None	227	56.3
Cardiovascular disease	57	14.2
DM	21	5.2
Noncardiovascular disease	98	24.3

**Table 2.** The lifestyle of individuals who came for service at King Chulalongkorn Memorial Hospital.

Behavior	No. of person	%
<b>Work activity level</b>		
Inactive (0-50 Kcal/d)	2	0.5
Low (50-799 Kcal/d)	160	39.7
Moderate (800-1599 Kcal/d)	112	27.8
High (>1,600 Kcal/d)	129	32.0
<b>TV watching hours</b>		
<1 hour	41	10.2
1-2.5 hours	176	43.7
2.5-4 hours	110	27.3
>4 hours	76	18.8
<b>TV watching activity</b>		
Sitting	214	53.1
Eating	94	23.3
Walking/doing chores	86	21.3
Exercising	6	1.5
others	3	0.7
<b>Leisure time activity level</b>		
Low (0-799 Kcal/d)	214	53.1
Moderate (800-1,599 Kcal/d)	132	32.8
High (>1,600 Kcal/d)	57	14.1
<b>Sleeping hours</b>		
<6 hours	88	21.8
6-8 hours	288	71.5
>8 hours	27	6.7
<b>Quality of sleep</b>		
Good	231	57.3
Bad	172	42.7
<b>Diet</b>		
Light	46	11.4
Moderate	233	57.8
Heavy	124	30.8
<b>Smoking status</b>		
Never smoke	316	78.4
Presently smoking	46	11.4
Quitted smoking	41	10.2
<b>Drinking status</b>		
Never drink	268	66.5
Presently drinking	73	18.1
Quitted drinking	62	15.4

**Table 3.** Association between sociodemographic characteristics and Overweight and Obesity.

Sociodemographic characteristics	BMI group		$\chi^2$	p-value	df
	< 23(n=226)	≥ 23(n=117)			
<b>Sex</b>					
Male	73	83	8.908	0.003**	1
Female	153	94			
<b>Age group</b>					
20 – 35 years	122	47	30.668	0.001**	1
over 35 years	28	29			
<b>Marital status</b>					
Single	114	50	6.62	0.000**	3
Married	92	111			
Divorced	7	8			
Widowed	13	8			
<b>Education</b>					
Below elementary	1	5	8.867	0.031*	3
Elementary	26	33			
High school	61	48			
Above high school	138	91			
<b>Underlying disease</b>					
None	143	84	28.028	0.000**	3
Cardiovascular disease	16	41			
DM	7	14			
Noncardiovascular disease	60	38			

\*p-value<0.05 \*\* p-value<0.01

The association between sociodemographic characteristics and overweight and obesity that were found statistically significant with p-value<0.05 were sex (p=0.003), age above 35 years, marital status (p=0.000), education (p=0.031), and underlying disease (p=0.000). It is worthy to note that the proportion of female that were not overweight was double of those that were overweight. Also, married individuals were more overweight than single

individuals. Those who had education above high school were mostly not overweight, unlike those with lower education. Lastly, it was found that individuals who had no underlying disease or had noncardiovascular disease were not overweight like those who had cardiovascular disease or diabetes mellitus. All of these sociodemographic characteristics which were significant will undergo multivariate analysis—multiple logistic regression.



**Table 4.** Association between lifestyle and overweight and obesity.

Lifestyle	BMI group		$\chi^2$	p-value	df
	<23	≥ 23			
<b>Work activity level</b>					
Inactive or Low (0-799 Kcal/d)	101	61	4.319	0.038*	1
Moderate or High (>800 Kcal/d)	125	116			
<b>TV watching activity</b>					
Sitting and eating	170	141	1.11	0.292	1
Walking/doing chores, exercise	56	36			
<b>Leisure time activity level</b>					
Inactive or Low (0-799 Kcal/d)	124	90	0.644	0.422	
Moderate or High (>800 Kcal/d)	102	87			
<b>Sleeping hours</b>					
<6 hours	38	38	2.539	0.406	2
6-8 hours	146	108			
>8 hours	7	8			
<b>Quality of sleep</b>					
Good	137	94	2.290	0.130	1
Bad	89	83			
<b>Diet</b>					
Light	28	18	1.380	0.502	2
Moderate	125	108			
Heavy	73	51			
<b>Smoking status</b>					
Never smoke	176	140	0.087	0.768	1
Quitted, present smoking	50	37			
<b>Drinking status</b>					
Never drink	159	109	3.429	0.064	1
Quitted, present drinking	67	68			

\*p-value<0.05

**Table 5.** Multiple Logistic Regression Analysis.

Factors	$\beta$	SE	WALD	p-value	ExB	95% CI of OR
Sex	-0.630	.233	7.899	.005	.533	0.344 - 0.824
Age	0.955	.25	15.274	.000	2.657	1.628 – 4.337
Marital status	0.442	.23	3.70	.054	1.557	0.992 – 2.443
Underlying Disease	0.292	.229	1.629	.202	1.339	0.855 – 2.098
Work level	0.442	.222	3.982	.046	1.556	1.082 - 2.402

R<sup>2</sup>=.162

The association between lifestyle and overweight and obesity that was found to have statistical significance with p-value<0.05 was work activity level. Work activity level which was the only significant behavior will undergo multivariate analysis—multiple logistic regression.

From multiple logistic regression analysis, there are 4 significant data. It was found that female had a 0.577 times lesser risk of being overweight compared to male (95 % CI=0.344-0.824, p=0.005). However, those who had over 35 years old had 2.657 times more risk of being overweight (95 % CI=1.628-4.337, p=0.000) and those who had moderate or high work activity level had 1.556 time more risk of being overweight than high work level (95 % CI= 1.082 – 2.402). Interestingly, in the present study underlying disease was not a significant risk for being overweight at all. The model summary according to the Nagelkerke R square was 0.162, meaning the factors listed in this study was significantly involved in being overweight takes up 16.2 % of the total model.

## Discussion

The prevalence of overweight and obesity according to the WPRO (World Health Organization

Western Pacific Region) criteria is 43.9 % in this study. According to the National Nutritional Surveys during the year 1986 and 1995 by the Department of Health, MOPH, the trend of overweight (BMI  $\geq$  25) had increased, being most prominent in ages 40-59 years old. However, the sample in this study consist mostly of people of working age and the prevalence of overweight and obesity might be lower than the general population. Additionally the sample consisted of female more than male. This may caused the prevalence of overweight and obesity to lower than the general population, due to the fact that female are at lesser risk of being overweight than male as seen from the result of this study. The occurrence of gene x and environmental factors interactions makes it more difficult to interpret the specific roles of genetics and lifestyle in obesity risk.<sup>(33)</sup> Nevertheless, the fundamental causes for the obesity epidemic are changing behaviors and lifestyles.<sup>(34)</sup> Women in this study tend to be lower in BMI than men. This may be explained by the self-conscious awareness of body image of women in the working age. Unlike women ages above 35 or women who stays at home as housewives, working women look after their shapes and watch their weight, more so than men. This is

universal for working women around the world. However, in the Thai culture, working men aged above 35 with a slight abdominal obesity are viewed as being financially stable, which men may appreciate being slightly obese. Thus, in this context, men usually have higher BMI than women.

As reviewed previously, individuals who were married tend to have higher BMI than those who are single.<sup>(35, 36)</sup> In this study, the proportion of individuals who were married and who were single are approximately the same. A number of reasons could be behind this. Married individuals are not as self-conscious about their body image as before being married, due to the fact that they no longer more concern to use their image to attract a partner. Also, married individuals can enjoy eating with their partner and children almost every day, more so than singles who may occasionally share meals with friends and family. Married individuals also tend to cook for themselves and their families more than singles, and the quantity could be more than the meals outside. However, it depends whether the married individual was health conscious about his/her weight or not, because cooking could help control weight better than eating outside meals. However, as the data results have shown, most of the individuals in this study had moderate to heavy diet, although it is not specified whether the diet was large in proportion or heavy on the fat and sweets. As for diet it had been expected that those who had heavy diet should have higher BMI than those who had light diet<sup>(37-40)</sup> however it was not found association between diet and overweight and obesity in this study.

However, this study contains mostly of individuals with high educational levels. It is known

that the higher the educational level, the lower the BMI.<sup>(41)</sup> In this study, the prevalence of overweight and obesity may be lower than the general population due to the fact that most individuals in this study are of high educational status. This may be so because individuals with higher education understand the risks involved in being obese, were usually of higher socioeconomic status, and tends to have knowledge of how to control weight more so than those with lower education. It was interesting to note that the sample of this study consisted mainly of individuals who had no underlying disease and those that had underlying diseases mostly had noncardiovascular diseases. As reviewed, overweight and obesity is related with a number of cardiovascular diseases and DM. Had the sample consist mostly of individuals with such comorbidities, it would not have represented the general population and the prevalence of overweight and obesity would be higher than the general population. The sample selected has little proportion of such diseases, reflecting the general population in this aspect. Obesity is associated with various comorbidities. In this study, those who had no underlying disease or had only noncardiovascular disease were not as overweight as those who had cardiovascular diseases or diabetes mellitus. The true nature of this association is yet to be studied, whether obesity causes such diseases or such diseases usually manifest in obesity. However, it could be concluded that these diseases were significantly found together with overweight and obesity.

The overall lifestyle of individuals who attended King Chulalongkorn Memorial Hospital was low working activity, with leisure time activity being low, sitting while watching an average of 1-2.5 hours

of television and taking a moderate to heavy diet. However, most individuals had healthy practices such as sleeping 6-8 hours, no smoking, and no drinking. Work activity level varies widely from low to high, with the proportion being the same for both.

It was found that work activity level was the only one factor in the aspect of lifestyle that associated with overweight and obesity. However, this work activity level was measured in terms of energy expenditure which resulted from the calculation of the type of work activity measured in METs and the duration of that activity. Focus only on the type of work activity may help increase the level of association to overweight and obesity, thus determining whether the type of work activity can affect BMI significantly. Light work-related physical activity was the strong predictors of weight gain.<sup>(42-46)</sup> In the present study those who had moderate or high work activity were found more overweight and obese than lower work activity, it may due to they consumed more food for energy but the final net energy was excess.

Many studies have shown diet and sedentary lifestyle to affect BMI and have significant association with overweight and obesity.<sup>(47-49)</sup> TV watching has been a controversial issue, whether it affect BMI or not and why.<sup>(50-55)</sup> Some consider it to be a sedentary activity and replace time which could be for moderate levels of physical activities such as exercise. In this study, it was interesting to note that most Thai individuals in this study watch only 1-2.5 hours of television which might be too short a duration to cause any significance to overweight and obesity. Also, most who watch television tend to sit but because the duration may be short, it did not matter what activity was done in front of the television. However, it had

been expected that those who exercise while watching television will have significantly lower BMI than those who did not exercise. However, the sample group of those who exercise in front of the television was too small to analyze accurately. Also, no data was collected to determine the duration or intensity of exercise which may have effect on BMI because the more longer more vigorous the exercise, the more energy was expended and should have effect on BMI.

While the proportion of good and bad sleep may be the same, it was found that less of those in the good sleep group were overweight while in the bad sleep group, the proportion of those overweight and not overweight were the same. Although review has shown that sleep duration affect BMI<sup>(56-58)</sup>, but in this study, sleep duration had no significant association to BMI. However, sleep quality does. It may be possible that the sample in this subgroup was too little to find the true proportion that may occur in the general population. Another inference from the data on sleep was that Thai people sleep adequately in terms of hours but half of the samples have sleep problems. It would be worthy to research more on this topic, accurately defining the term "bad sleep", listing out the sleep problems involved, and analyzing the effect the quality of sleep might have on other chronic diseases.

Regarding drinking and smoking status, it was not found the association with overweight. Although some studies noted that those who currently smoke have lower BMI than nonsmoker<sup>(59-65)</sup> but moderately amount consumption of alcohol has no effect on BMI.<sup>(66-67)</sup> However in the present study the amount of cigarettes or alcohol consumed had not been analyzed.

However, this study was only meant to provide approximate initial data to give the overall idea of lifestyle found in Thai people and the association it may have on overweight and obesity. Its generalization to the general population may be difficult because it had only been done in one specialized setting which is the hospital. To generalize this type of data, samples must be taken from a larger, heterogeneous population. As for the recording of lifestyle, an interview would provide a more comprehensive picture of the different behaviors involved and new information on the variety of behavior may be found for further studies. Nevertheless, overweight and obesity according to the criteria of the WPRO had increased to almost half in the population studied. Factors associated with overweight and obesity were sex, age above 35 years, marital status, education, underlying disease and work activity level. Recognition and prevention of overweight and obesity particularly in those with associated factors may be beneficial for reducing burden of overweight and obesity including other following health problems.

## References

1. World Health Organization. Obesity: preventing and managing the global epidemic. Report of a WHO Consultation. Geneva: World Health Organization, 2000. (WHO Technical Report Series, No. 894)
2. Clinical guidelines on the identification and treatment of overweight and obesity in adults—The evidence report. National Institutes of Health. *Obes Res* 1998 Sep;6 Suppl 2:51S–209S
3. Deurenberg P, Yap M, van Staveren WA. Body mass index and percent body fat: a meta analysis among different ethnic groups. *Int J Obes Relat Metab Disord* 1998 Dec; 22(12): 1164-71
4. Zhou BF. Predictive values of body mass index and waist circumference for risk factors of certain related diseases in Chinese adults—study on optimal cut-off points of body mass index and waist circumference in Chinese adults. *Biomed Environ Sci* 2002 Mar;15(1): 83-96
5. Ko GT, Tang J, Chan JC, Sung R, Wu MM, Wai HP, Chen R. Lower BMI cut-off value to define obesity in Hong Kong Chinese: an analysis based on body fat assessment by bioelectrical impedance. *Br J Nutr* 2001 Feb;85(2):239-42
6. Deurenberg-Yap M, Schmidt G, van Staveren WA, Deurenberg P. The paradox of low body mass index and high body fat percentage among Chinese, Malays and Indians in Singapore. *Int J Obes Relat Metab Disord* 2000 Aug;24(8): 1011-7
7. Pan WH, Flegal KM, Chang HY, Yeh WT, Yeh CJ, Lee WC. Body mass index and obesity-related metabolic disorders in Taiwanese and US whites and blacks: implications for definitions of overweight and obesity for Asians. *Am J Clin Nutr* 2004 Jan;79(1):31-9
8. Chang CJ, Wu CH, Chang CS, Yao WJ, Yang YC, Wu JS, Lu FH. Low body mass index but high percent body fat in Taiwanese subjects: implications of obesity cutoffs. *Int J Obes Relat Metab Disord* 2003 Feb;27(2):253-9
9. Anuurad E, Shiwaku K, Nogi A, Kitajima K, Enkhmaa B, Shimono K, Yamane Y. The new BMI



- criteria for asians by the regional office for the western pacific region of WHO are suitable for screening of overweight to prevent metabolic syndrome in elder Japanese workers. *J Occup Health* 2003 Nov;45(6): 335-43
10. Shiwaku K, Anuurad E, Enkhmaa B, Nogi A, Kitajima K, Shimono K, Yamane Y, Oyunsuren T. Overweight Japanese with body mass indexes of 23.0-24.9 have higher risks for obesity-associated disorders: a comparison of Japanese and Mongolians. *Int J Obes Relat Metab Disord* 2004 Jan;28(1):152-8
  11. Gurrici S, Hartriyanti Y, Hautvast JG, Deurenberg P. Relationship between body fat and body mass index: differences between Indonesians and Dutch Caucasians. *Eur J Clin Nut* 1998 Nov;52(11):779-83
  12. WHO/IASO/IOTF. The Asia-Pacific perspective: redefining obesity and its treatment. Health Communications Australia Pty Ltd, 2000
  13. WHO expert consultation. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *Lancet* 2004 Jan 10;363(9403): 157-63
  14. Pi-Sunyer FX. Comorbidities of overweight and obesity: current evidence and research issues. *Med Sci Sports Exerc* 1999 Nov;31 (11 Suppl):S602-8
  15. Doll HA, Petersen SE, Stewart-Brown SL. Obesity and physical and emotional well-being: associations between body mass index, chronic illness, and the physical and mental components of the SF-36 questionnaire. *Obes Res* 2000 Mar;8(2):160-70
  16. Rohrer JE, Rush Pierce JR Jr, Blackburn C. Lifestyle and mental health. *Prev Med* 2005 Apr;40(4):438-43
  17. Jousilahti P, Tuomilehto J, Vartiainen E, Pekkanen J, Puska P. Body weight, cardiovascular risk factors, and coronary mortality. 15 year follow-up of middle-aged men and women in eastern Finland. *Circulation* 1996 Apr 1; 93(7):1372-9
  18. LaBarthe DR. *Epidemiology and Prevention of Cardiovascular Diseases: A Global Challenge*. Gaithersburg, Md: Aspen Publishers, 1998
  19. Clark JM, Brancati FL, Diehl AM. Nonalcoholic fatty liver disease. *Gastroenterology* 2002 May;122(6):1649-57
  20. Angulo P. Nonalcoholic fatty liver disease. *N Engl J Med* 2002 Apr 28;346(16):1221-31
  21. Sheth SG, Gordon FD, Chopra S. Nonalcoholic steatohepatitis. *Ann Intern Med* 1997 Jan 15; 126(2):137-45
  22. Leung RS, Bradley TD. Sleep apnea and cardiovascular disease. *Am J Respir Crit Care Med* 2001 Dec 15;164(12):2147-65
  23. Peker Y, Hedner J, Kraiczi H, Loth S. Respiratory disturbance index: an independent predictor of mortality in coronary artery disease. *Am J Respir Crit Care Med* 2000 Jul;162(1):81-6
  24. Peppard PE, Young T, Palta M, Skatrud J. Prospective study of the association between sleep-disordered breathing and hypertension. *N Engl J Med* 2000 May 11;342(19):1378-84
  25. Grunstein RR, Stenlof K, Hedner J, Sjöström L. Impact of obstructive sleep apnea and sleepiness on metabolic and cardiovascular

- risk factors in the Swedish Obese Subjects (SOS) Study. *Int J Obes Relat Metab Disord* 1995 Jun;19(6):410-8
26. International Agency for Research on Cancer. *Weight Control and Physical Activity*. Lyon, France: IARC Press, 2002
27. Hill AJ, Williams J. Psychological health in a non-clinical sample of obese women. *Int J Obes Relat Metab Disord* 1998 Jun;22(6):578-83
28. Willett WC, Dietz WH, Colditz GA. Guideline for healthy weight. *N Engl J Med* 1999 Aug 5; 341(6):427-34
29. Thompson D, Edelsberg J, Colditz GA, Bird AP, Oster G. Lifetime health and economic consequences of obesity. *Arch Intern Med* 1999 Oct 11;159(18):2177-83
30. Swinburn BA, Caterson I, Seidell JC, James WP. Diet, nutrition and the prevention of excess weight gain and obesity. *Public Health Nutr* 2004 Feb;7(1A):123-46
31. Mullis RM, Blair SN, Aronne LJ, Bier DM, Denke MA, Dietz W, Donato KA, Drewnowski A, French SA, Howard BV, et al. Prevention Conference VII: Obesity, a worldwide epidemic related to heart disease and stroke: Group IV: prevention/treatment. *Circulation* 2004 Nov 2;110(18):e484-8
32. Popkin BM, Gordon-Larsen P. The nutrition transition: worldwide obesity dynamics and their determinants. *Int J Obes Relat Metab Disord* 2004 Nov;28 Suppl 3:S2-9
33. Marti A, Moreno-Aliaga MJ, Hebebrand J, Martinez JA. Genes, lifestyles and obesity. *Int J Obes Relat Metab Disord* 2004 Nov;28 Suppl 3: S29-36
34. Press Release WHO/46. Obesity Epidemic puts Millions at risk from Related Diseases. 12 June 1997
35. Jeffery RW, Rick AM. Cross-sectional and longitudinal associations between body mass index and marriage-related factors. *Obes Res* 2002 Aug;10(8):809-15
36. Lipowicz A, Gronkiewicz S, Malina RM. Body mass index, overweight and obesity in married and never married men and women in Poland. *Am J Hum Biol* 2002 Jul-Aug;14(4): 468-75
37. Astrup A. The role of dietary fat in the prevention and treatment of obesity. Efficacy and safety of low-fat diets. *Int J Obes Relat Metab Disord* 2001 May;25 Suppl 1:S46-50
38. Astrup A. Macronutrient balances and obesity: the role of diet and physical activity. *Public Health Nutr* 1999 Sep;2(3A):341-7
39. Jebb SA, Moore MS. Contribution of a sedentary lifestyle and inactivity to the etiology of overweight and obesity: current evidence and research issues. *Med Sci Sports Exerc* 1999 Nov;31 (11 Suppl):S534-41
40. Ruidavets JB, Bongard V, Bataille V, Gourdy P, Ferrieres J. Eating frequency and body fatness in middle-aged men. *Int J Obes Relat Metab Disord* 2002 Nov;26(11):1476-83
41. Maddah M, Eshraghian MR, Djazayeri A, Mirdamadi R. Association of body mass index with educational level in Iranian men and women. *Eur J Clin Nutr* 2003 Jul;57(7):819-23
42. Gutierrez-Fisac JL, Guallar-Castillon P, Diez-Ganan L, Lopez Garcia E, Banegas Banegas JR, Rodriguez Artalejo F. Work-related physical

- activity is not associated with body mass index and obesity. *Obes Res* 2002 Apr;10(4):270-6
43. McCarthy SN, Gibney MJ, Flynn A. Overweight, obesity and physical activity levels in Irish adults: evidence from the North/South Ireland food consumption survey. *Proc Nutr Soc* 2002 Feb;61(1):3-7
44. Bell AC, Ge K, Popkin BM. Weight gain and its predictors in Chinese adults. *Int J Obes Relat Metab Disord* 2001 Jul;25(7):1079-86
45. Brown WJ, Miller YD, Miller R. Sitting time and work patterns as indicators of overweight and obesity in Australian adults. *Int J Obes Relat Metab Disord* 2003 Nov;27(11):1340-6
46. Martinez-Gonzalez MA, Martinez JA, Hu FB, Gibney MJ, Kearney J. Physical inactivity, sedentary lifestyle and obesity in the European Union. *Int J Obes Relat Metab Disord* 1999 Nov;23(11):1192-201
47. Buchowski MS, Sun M. Energy expenditure, television viewing and obesity. *Int J Obes Relat Metab Disord* 1996 Mar;20(3):236-44
48. Tucker LA, Bagwell M. Television viewing and obesity in adult females. *Am J Public Health* 1991 Jul;81(7):908-11
49. Hu FB, Li TY, Colditz GA, Willett WC, Manson JE. Television watching and other sedentary behaviors in relation to risk of obesity and type 2 diabetes mellitus in women. *JAMA* 2003 Apr 9;289(14):1785-91
50. Kronenberg F, Pereira MA, Schmitz MK, Arnett DK, Evenson KR, Crapo RO, Jensen RL, Burke GL, Sholinsky P, Ellison RC, et al. Influence of leisure time physical activity and television watching on atherosclerosis risk factors in the NHLBI Family Heart Study. *Atherosclerosis* 2000 Dec;153(2):433-43
51. Salmon J, Bauman A, Crawford D, Timperio A, Owen N. The association between television viewing and overweight among Australian adults participating in varying levels of leisure-time physical activity. *Int J Obes Relat Metab Disord* 2000 May;24(5):600-6
52. Jakes RW, Day NE, Khaw KT, Luben R, Oakes S, Welch A, Bingham S, Wareham NJ. Television viewing and low participation in vigorous recreation are independently associated with obesity and markers of cardiovascular disease risk: EPIC-Norfolk population-based study. *Eur J Clin Nutr* 2003 Sep;57(9):1089-96
53. Crawford DA, Jeffery RW, French SA. Television viewing, physical inactivity and obesity. *Int J Obes Relat Metab Disord* 1999 Apr;23(4):437-40
54. Jeffery RW, French SA. Epidemic obesity in the United States: are fast foods and television viewing contributing? *Am J Public Health* 1998 Feb; 88(2):277-80
55. Gore SA, Foster JA, DiLillo VG, Kirk K, Smith West D. Television viewing and snacking. *Eat Behav* 2003 Nov;4(4):399-405
56. Spiegel K, Tasali E, Penev P, Van Cauter E. Brief communication: Sleep curtailment in healthy young men is associated with decreased leptin levels, elevated ghrelin levels, and increased hunger and appetite. *Ann Intern Med* 2004 Dec 7;141(11):846-50
57. Spiegel K, Leproult R, Van Cauter E. Impact of

- sleep debt on metabolic and endocrine function. *Lancet* 1999 Oct 23;354(9188):1435-9
58. Hasler G, Buysse DJ, Klaghofer R, Gamma A, Ajdacic V, Eich D, Rossler W, Angst J. The association between short sleep duration and obesity in young adults: a 13-year prospective study. *Sleep* 2004 Jun 15;27(4):661-6
59. Akbartabartoori M, Lean ME, Hankey CR. Relationships between cigarette smoking, body size and body shape. *Int J Obes (Lond)* 2005 Feb;29(2):236-43
60. Barrett-Connor E, Khaw KT. Cigarette smoking and increased central adiposity. *Ann Intern Med* 1989 Nov 15; 111(10):783-7
61. Bamia C, Trichopoulou A, Lenas D, Trichopoulos D. Tobacco smoking in relation to body fat mass and distribution in a general population sample. *Int J Obes Relat Metab Disord* 2004 Aug;28(8):1091-6
62. Visser M, Launer LJ, Deurenberg P, Deeg DJ. Past and current smoking in relation to body fat distribution in older men and women. *J Gerontol A Biol Sci Med Sci* 1999 Jun;54(6):M293-8
63. Troisi RJ, Heinold JW, Vokonas PS, Weiss ST. Cigarette smoking, dietary intake, and physical activity: effects on body fat distribution—the Normative Aging Study. *Am J Clin Nutr* 1991 May; 53(5):1104-11
64. Rasmussen F, Tynelius P, Kark M. Importance of smoking habits for longitudinal and age-matched changes in body mass index: a cohort study of Swedish men and women. *Prev Med* 2003 Jul;37(1):1-9
65. Molarius A, Seidell JC. Differences in the association between smoking and relative body weight by level of education. *Int J Obes Relat Metab Disord* 1997 Mar;21(3):189-96
66. Cordain L, Bryan ED, Melby CL, Smith MJ. Influence of moderate daily wine consumption on body weight regulation and metabolism in healthy free-living males. *J Am Coll Nutr* 1997 Apr;16(2):134-9
67. Flechtner-Mors M, Biesalski HK, Jenkinson CP, Adler G, Ditschuneit HH. Effects of moderate consumption of white wine on weight loss in overweight and obese subjects. *Int J Obes Relat Metab Disord* 2004 Nov;28(11):1420-6